

Total and regional Runoff to the Baltic Sea

Author: Pia Andersson, SMHI

Key message

During the last 8 years the total inflow has decreased from a top flow rate of over 17500 m³/s in 1998 to less than 11000 in 2003. Since 2003, the runoff has slightly increased and in 2005 the runoff was just below the mean values reaching 13876 m³/s.

All sub basins but the Gulf of Finland, had lower runoff values in the year 2005 compared to the mean values. The Gulf of Bothnia, the Gulf of Finland and the Gulf of Riga, the difference to the mean value was less than 1%. In the Baltic Proper, the difference to the mean value was more than 9%. When looking at the entire Baltic Sea, the difference to the mean value was 2% hence the general runoff for the entire area was slightly lower compared to the yearly mean runoff value.

During the period 1950 – 2005, the total runoff to the Baltic Sea area shows no long-term trends. On the other hand this time period is characterised by dry and wet periods lasting for a couple of years to a decade generally following the NAO index.




Figure 1: Total runoff deviation during the years 1950 - 2005 to the Baltic sub basins based on annual mean values. The mean runoff value for each sub basin is written within each figure. The black line is the running mean over 5 years.

Results and Assessment

Relevance of the indicator for describing developments in the environment

Runoff is a quantitative background indicator on the freshwater discharge, carrying the nutrients from the drainage areas to the coast.

Runoff is an important parameter on the change of pressure on nutrient supply due to varying climate and climate change. Also change in land-use can influence runoff. To evaluate the change of pressure on nutrient supply to the Baltic region it is necessary to know the variability of runoff and normalise for this natural variability. Dry periods, like the one during the 1970s, can mask the marine eutrophication since the runoff was lower than average and hence also the total load of nutrients. Extended dry periods should also lead to a slight increase in surface layer salinity. During wet periods, the total nutrient load (pressure) increase, making marine eutrophication (effects) even worse.

The indicator shows the annual runoff from drainage areas but integrated over the Baltic sub-regions. Runoff is governed by the precipitation - evaporation on land areas and is also influenced by air temperature. It is the sum of direct river and diffusive runoff. In all sub-regions a strong seasonal, annual

and decadal variability can be distinguished. Especially wet and dry periods are characterising the runoff. The 1970s were a fairly dry period compared with the 1980s and the later part of the 1990s. Geographically, the runoff is of about the same size in the Gulf of Finland and the Baltic Proper, whereas the Gulf of Riga contributes to a lesser extent and the Gulf of Bothnia to a larger extent to the total runoff.

Assessment

Four different sub basins are described as deviation from their mean value based on runoff during 1950 to 2005. The mean value is included as a textbox inside each sub basin figure (figure 1). The years with higher runoff compared to the mean value is displayed as a red bar in the positive direction and a lower value with a blue bar in the negative direction. A running mean over five years is displayed as a black line overlaying the bars in the figure. The sub basins are displayed in the centre of figure 1 and the sub basins described are the Baltic Proper, the Gulf of Riga, the Gulf of Finland and the Gulf of Bothnia. A figure with the sum of the Baltic Sea sub basins is also included, partly to give an overview of the entire Baltic Sea and partly to compare the annual changes to the NAO index.

During the period 1950 – 2005, there is no obvious trend in the annual runoff during the last 50 years, neither in the total runoff to the Baltic Sea area, nor in the sub-regions. On the other hand this time period is characterised by dry and wet periods lasting for a couple of years to a decade. 2005 was a year with runoff under the normal values, see table 1. All sub basins but the Gulf of Finland, had lower runoff values in the year 2005 compared to the mean values from 1950 to 2005. The Gulf of Bothnia, the Gulf of Finland and the Gulf of Riga, the difference to the mean value was less than 1%. In the Baltic Proper, the difference to the mean value was more than 9%. When looking at the entire Baltic Sea, the difference to the mean value was 2% hence the general runoff for the entire area was slightly lower compared to the yearly mean runoff value.

Table 1: Mean of the annual mean runoff values [m³/s] for the sub basins in the Baltic Sea and the Kattegat compared to the 2005 values and the difference in %.

	Mean runoff 1950-2005 [m ³ /s]	Runoff 2005 [m ³ /s]	Difference from mean [%]
Gulf of Bothnia	6052.2	6041.9	-0.170186048
Gulf of Finland	3567.5	3592.2	0.692361598
Gulf of Riga	1019	1013.5	-0.539744848
Baltic Proper	3554.4	3228.7	-9.163290569
Total Baltic Sea	14193	13876.2	-2.23208624

At times, there have been similar features in the changes of the runoff values for all the sub basins. Other time periods, the changes are similar only in some of the sub basins. All the sub basins had low runoff

values in the early to the mid 70's and higher in the end of the 90's. In the Baltic Proper, the Gulf of Riga and the Gulf of Finland, there were high values from the mid 50's to the start of the 60's. In the Gulf of Bothnia, the Gulf of Riga and the Gulf of Finland, there was a feature of increasing values during the 80's while in the Baltic Proper, there was a feature of decreasing values. There were low values in the Baltic Proper in the early 90's while there were high values in the end of the 80's and the start of the 90's in the Gulf of Riga, the Gulf of Finland and the Gulf of Bothnia.

The total runoff to the Baltic Sea is mostly influenced by the sub basins with the largest contributions, obviously. The highest contribution is from the Gulf of Bothnia followed by the Gulf of Finland and the Baltic Proper. By comparing the Gulf of Bothnia to the Gulf of Finland, there is a rather good correlation in the features of the running mean values. By comparing the Gulf of Bothnia to the Baltic Proper, there are some correlations but also some deviations in the features. By making a generalization, the figure displaying the total runoff to the Baltic Sea, represents the general features of the different sub basins rather well. By making this generalisation, only the figure displaying the total runoff to the Baltic Sea is compared to the NAO index.




Figure 2: Total runoff deviation during 1950 - 2005 to the Baltic Sea and the NAO index during the years 1864 - 2005 based on winter mean values of the NAO index. Positive index indicates stronger westerly winds bringing warmer and wetter winters to Scandinavia. The black line is the running mean over 5 years.

Figure 2 is displaying the total runoff deviation during 1950 - 2005 to the Baltic Sea both as integrated difference (sum of abnormalities, starting and ending with 0 km³ 1950 and 2005) and with bars displaying the year to year deviation from the mean. The NAO index during the years 1864 - 2005 based on winter mean values of the NAO index is displayed in the bottom of the figure. The integrated difference gives an idea of the total amount of runoff in the Baltic Sea. The black line is the running mean over 5 years. By comparing the running mean of the two lower figures between 1952 and 2003, the features correspond rather well with each other. Since the correlation is good, the NAO indices may be used to indicate general runoff to the Baltic Sea back in time. This is the motivation of including the longer time series of the NAO indices as well as looking for possible oscillation features extending the 55 year time scale. There is a slight tendency to high NAO indices around 1910 and 1990 and low values around 1880 and 1960. This forms an oscillation over 80 years. By looking at this 1.5 oscillation, it is not possible to draw conclusions, but comments can be made. To go from generally high values to generally low values takes approximately 50 years while going from generally low values to generally high values takes approximately 30 years. The recent peak of the NAO index occurred in 1990. If the comment concerning the 80 year oscillation is true, we are 15 years downhill the NAO index slope and we will continue that ride for approximately 35 years, with weaker westerly winds resulting in colder and dryer winters in Scandinavia.

References

Bergström, S. And B. Carlsson 1994. River runoff to the Baltic Sea 1950 - 1990. AMBIO Vol. 23, No. 4-5, 280 - 287.

Graham, Phil 1999. Modelling runoff to the Baltic Sea. AMBIO Vol. 28, No. 4, 328-334.

<http://www.cgd.ucar.edu/cas/jhurrell/indices.data.html#naostatdjfm>

Data

Observations are collected at the BALTEX Hydrological Data Centre (<http://www.smhi.se/sgn0102/bhdc/bhdc.htm>), whereas modelled data is done at SMHI using the HBV-model (Graham-99). Gulf of Riga runoff is based observations up through 2001, while simulations are used for 2002. Gulf of Finland runoff is based on observations up through 1997, while simulations are used for 1998-2002. Baltic Proper runoff is based on observations up through 1996, while simulations are used for 1997-2003. For 2003 to 2005 all data is based on model simulations carried out by Dr. Phil Graham, SMHI. The NAO indices are collected from <http://www.cgd.ucar.edu/cas/jhurrell/indices.data.html#naostatdjfm>.

Table 2: NAO index and annual mean values of runoff [m³/s] to the sub basins in the Baltic Sea during 1950 to 2005.

Year	NAO index	Bothnian Bay [m ³ /s]	Bothnian Sea [m ³ /s]	Gulf of Finland [m ³ /s]	Gulf of Riga [m ³ /s]	Baltic Proper [m ³ /s]	Total Baltic Sea [m ³ /s]	% Notes:
1950	1.40	2 867.7	3 225.7	3 409.3	1 081.9	3 501.7	14 086.3	%BB & BS are observations
1951	-1.26	2 904.1	2 667.8	3 264.0	1 322.5	3 336.7	13 495.1	%GOR is observations up

HELCOM Baltic Sea Environment Fact Sheet 2006

								through 2001
1952	0.83	3 625.1	2 614.6	3 001.8	1 216.7	3 106.0	13 564.3	%GOR uses HBV simulations for 2002
1953	0.18	3 785.6	3 363.7	3 969.4	1 282.4	3 255.4	15 656.5	%GOF is observations up through 1997
1954	0.13	3 512.3	2 810.4	3 536.1	825.7	2 782.8	13 467.3	%GOF uses HBV simulations for 1998-2002
1955	-2.52	3 251.4	2 448.4	4 209.0	1 082.9	3 489.8	14 481.4	%BP is observations up through 1993
1956	-1.73	2 563.8	2 569.5	3 784.0	1 258.2	3 725.2	13 900.7	%BP uses replacement stations for 1994-1996
1957	1.52	3 468.6	3 223.0	4 231.5	1 264.2	3 607.6	15 794.9	%BP uses HBV simulations for 1997-2002
1958	-1.02	3 025.8	2 451.4	4 395.9	1 244.8	4 552.2	15 670.1	
1959	-0.37	2 433.1	2 304.4	3 785.3	1 194.0	2 771.3	12 488.2	
1960	-1.54	2 329.2	2 940.5	2 868.0	1 107.4	3 704.6	12 949.6	
1961	1.80	3 352.8	3 458.4	3 296.4	892.5	3 329.4	14 329.4	
1962	-2.38	3 217.2	2 982.8	4 521.3	1 507.7	4 254.7	16 483.7	
1963	-3.60	2 577.2	2 282.3	3 867.8	707.1	2 814.5	12 248.8	
1964	-2.86	3 109.3	2 487.2	2 807.6	824.5	2 511.9	11 740.6	

HELCOM Baltic Sea Environment Fact Sheet 2006

1965	-2.88	3 566.1	2 734.0	2 679.8	790.4	3 479.7	13 250.1	
1966	-1.69	2 976.7	3 206.1	3 560.0	973.1	4 126.8	14 842.6	
1967	1.28	3 946.0	3 499.1	3 826.6	924.6	4 323.2	16 519.5	
1968	-1.04	2 838.1	2 593.1	3 667.5	859.2	3 757.2	13 715.1	
1969	-4.89	2 510.1	2 185.6	3 304.5	743.3	2 669.0	11 412.5	
1970	-1.89	2 473.5	2 492.3	3 185.0	983.3	4 223.2	13 357.4	
1971	-0.96	2 674.7	2 674.3	3 255.2	751.1	3 272.3	12 627.6	
1972	0.34	2 829.6	2 657.2	2 826.7	749.4	2 966.9	12 029.9	
1973	2.52	3 303.5	2 585.6	2 268.5	644.1	2 783.3	11 585.0	
1974	1.23	3 581.6	3 085.2	3 156.8	783.3	4 209.5	14 816.4	
1975	1.63	3 167.1	2 612.5	3 538.2	1 342.0	3 698.2	14 358.1	
1976	1.37	2 413.4	1 993.0	2 855.7	855.0	2 678.5	10 795.7	
1977	-2.14	3 484.4	2 844.5	3 256.7	855.5	4 180.1	14 621.2	
1978	0.17	2 642.0	2 514.1	3 498.4	1 299.9	4 038.7	13 993.0	
1979	-2.25	3 069.3	2 828.7	3 315.3	890.4	4 331.8	14 435.4	
1980	0.56	2 549.7	2 617.0	3 078.4	999.9	4 974.8	14 219.8	
1981	2.05	3 678.8	3 367.9	3 977.1	1 212.6	5 045.6	17 281.8	
1982	0.80	3 305.4	2 749.0	4 327.9	937.8	3 816.1	15 136.2	
1983	3.42	3 513.1	3 053.8	3 711.3	930.5	3 477.5	14 686.1	
1984	1.60	3 421.0	3 284.5	3 918.5	721.6	3 371.0	14 716.7	

HELCOM Baltic Sea Environment Fact Sheet 2006

1985	-0.63	3 360.8	3 611.7	3 476.7	988.9	3 886.7	15 324.7	
1986	0.50	3 131.2	3 129.3	3 700.3	1 024.4	3 691.3	14 676.4	
1987	-0.75	3 492.0	3 711.2	4 092.6	1 210.5	3 883.0	16 389.3	
1988	0.72	3 140.2	3 306.3	4 153.9	1 088.7	4 088.5	15 777.5	
1989	5.08	3 691.6	3 028.3	4 109.4	1 237.6	3 100.4	15 167.3	
1990	3.96	2 840.6	3 101.3	4 194.2	1 603.5	3 124.4	14 864.0	
1991	1.03	3 353.6	2 791.9	3 971.7	1 127.8	2 809.2	14 054.3	
1992	3.28	3 972.9	3 215.8	4 018.8	834.5	2 675.8	14 717.7	
1993	2.67	3 765.1	3 522.8	3 373.1	955.0	2 840.1	14 456.0	
1994	3.03	2 786.7	2 504.2	3 699.4	1 184.5	3 990.6	14 165.4	%replacement stations used for BP!
1995	3.96	3 249.8	3 204.2	4 047.0	1 029.5	3 761.0	15 291.5	%replacement stations used for BP!
1996	-3.78	2 795.7	2 013.4	2 950.6	651.5	3 239.3	11 650.4	%replacement stations used for BP!
1997	-0.20	3 056.3	2 739.6	3 147.8	975.7	3 420.7	13 340.1	%HBV simulations used for BP!
1998	0.72	4 173.6	3 883.9	4 009.2	1 437.0	4 040.9	17 544.5	%HBV simulations used for BP & GOF!
1999	1.70	3 390.0	2 916.5	3 679.1	1 015.1	3 908.6	14 909.3	%HBV simulations used for BP & GOF!

2000	2.80	4 204.0	4 099.0	3 487.6	967.3	3 567.0	16 324.9	%HBV simulations used for BP & GOF!
2001	-1.89	3 541.0	3 803.7	3 554.4	953.4	3 696.5	15 548.9	%HBV simulations used for BP & GOF!
2002	0.76	2 800.6	2 532.9	3 344.4	1 018.0	3 864.2	13 560.0	%HBV simulations used for BP, GOF & GOR!
2003	0.20	2 354.0	1 856.0	3 160.0	677.0	2 706.0	10 752.0	%HBV simulations used for all subbasins
2004	-0.07	3 190.0	2 245.0	3 864.0	1 011.0	3 354.0	13 665.0	%HBV simulations used for all subbasins
2005	0.12	3 357.5	2 684.4	3 592.2	1 013.5	3 228.7	13 876.2	%HBV simulations used for all subbasins

For reference purposes, please cite this Baltic Sea environment fact sheet as follows:

[Author's name(s)], [Year]. [Baltic Sea environment fact sheet title]. HELCOM Baltic Sea Environment Fact Sheets. Online. [Date Viewed], <http://www.helcom.fi/baltic-sea-trends/environment-fact-sheets/>.