


Baltic Sea Environment Proceedings No. 148



Baltic Marine Environment Protection Commission

Ecological coherence assessment of the
Marine Protected Area
network in the Baltic Sea



Ecological coherence assessment of the Marine Protected Area network in the Baltic Sea



HELCOM

This report has been developed under the State and Conservation Working Group and with the support of the HELCOM MPA Task Group. The work was funded by the Marine Group (HAV) at the Nordic Council of Ministers (HELCOM ECONET project). The Swedish Agency for Marine and Water Management (SwaM) is acknowledged for funding on the HELCOM MPA database, which enabled data collection for this assessment.

Published by:

Helsinki Commission

Katajanokanlaituri 6 B

FI-00160 Helsinki

Finland

<http://www.helcom.fi>

Authors

Janica Borg, Petra Kääriä, Ulla Li Zweifel

GIS analyses

Iiris Kokkonen, Joni Kaitaranta

Contributors

Dieter Boedeker, Penina Blankett, Jan Ekebom, Manuel Frias Vega, Andrzej Ginalski, Michael Haldin, Norbert Häubner, Anna Karlsson, Samuli Korpinen, Lasse Kurvinen, Florent Nicolas, Małgorzata Opechowska, Hanna Paulomäki

For bibliographic purposes, this document should be cited as:

HELCOM 2016. Ecological coherence assessment of the Marine Protected Area network in the Baltic. Balt. Sea Environ. Proc. No. 148

Information included in this publication or extracts thereof are free for citing on the condition that the complete reference of the publication is given as above.

Editorial corrections have been made to the electronic copy on pages 2 and 31.

Copyright 2016 by the Baltic Marine Environment Protection Commission – HELCOM

Layout: Bitdesign, Vantaa, Finland

Photo credits: Helmi Mentula, Metsähallitus 2014

Number of pages: 69

Printed by: Erweko Painotuote Oy, Finland

ISSN 0357-2994

Table of contents

Executive summary	5
1 Introduction	7
1.1 Purpose of assessing marine protected areas in the Baltic Sea	7
1.2 The concept of coastal and marine Baltic Sea protected areas (HELCOM MPAs ²) ...	7
2 Status of the network	9
2.1 The Natura 2000 network in the HELCOM area	10
3 Assessment of ecological coherence of the MPA network in the Baltic Sea	13
3.1 Representativity	14
3.1.1 Method and results of the representativity assessment of the HELCOM MPA network	15
3.1.2 Method and results of the representativity assessment of the combined network of HELCOM MPAs and marine Natura 2000 sites	20
3.2 Replication	22
3.2.1 Method and results of the replication assessment of the HELCOM MPA network	22
3.2.2 Method and results of the replication assessment of the combined network of HELCOM MPAs and marine Natura 2000 sites	26
3.3 Adequacy	27
3.3.1. Method and results of the adequacy assessment of the HELCOM MPA network	28
3.3.2 Method and results of the adequacy assessment of the combined network of HELCOM MPAs and marine Natura 2000 sites	33
3.4 Connectivity	33
3.4.1 Method and results of connectivity assessment of the HELCOM MPA network.	33
3.4.2 Method and results of connectivity of the combined network of HELCOM MPAs and marine Natura 2000 sites	36
3.5 Conclusions of the ecological coherence assessment	38
3.5.1. Current data gaps and proposal for improvements to the assessment.	39
4 Testing a new model for aggregating the results of subcriteria analyses of the ecological coherence assessment into a single outcome	40
4.1 Integration tables of the results of subcriteria evaluations of the HELCOM MPAs. .	40
4.2 Final outcome of the ecological coherence assessment for the HELCOM MPA network, using the integration tables for aggregating the subcriteria results	47
5 Recommendation 35/1 on coastal and marine Baltic Sea protected areas (HELCOM MPAs⁶)	48
5.1 Current status of implementation	48
5.1.1 Accomplishment of Recommendation 35/1	48
5.1.2 Summary on Recommendation 35/1 follow-up	53
5.1.3 Progress of additional commitments.	54
6 Next steps for improving the network and assessments of HELCOM MPAs.	55
6.1 Measures by countries.	55
6.2 Joint measures	56

References	58
Annex 1.	
Coverage of the HELCOM MPA network in 2013.....	60
Annex 2.	
Overview of the assessment design of the HELCOM ecological coherence assessment in 2016.....	61
Annex 3.	
Overview of the ecological coherence assessment carried out for (1) the HELCOM MPA network and (2) the combined network of the HELCOM MPAs and the marine Natura 2000 sites in the Baltic Sea.....	62
Overview table of the data used for the HELCOM ecological coherence assessment in 2016.....	63
Annex 4.	
Overview table of the data used for the HELCOM ecological coherence assessment in 2016.....	63
Annex 5.	
Fishing effort within HELCOM MPAs.....	64
Annex 6.	
Rationale for uncertainty ranking and values in the integration tables used for aggregating the subcriteria results (chapter 4.1).....	66
Annex 7.	
Follow up of HELCOM Recommendation 35/1 on the system of coastal and marine Baltic Sea protected areas (HELCOM MPAs)	68

Executive summary

The Baltic Sea Action Plan sets out, as one of its main goals, to achieve ‘a favourable status of Baltic Sea biodiversity’. Designation of marine protected areas (MPAs) has been an instrument for protection in the Baltic Sea for more than 30 years and serves as an important measure to meet this commitment of the Contracting Parties to the Helsinki Convention. To reach an ecologically coherent network of MPAs, i.e. a network of protected sites which deliver more benefits than individual MPAs, has been on the agenda of HELCOM since the first joint Ministerial Meeting of the Helsinki and OSPAR Commissions in Germany 2003, when Member States of these two conventions and the European Commission agreed upon a Joint Work Programme (JWP). The fundament of the HELCOM Recommendation 35/1 on the system of coastal and marine Baltic Sea protected areas (HELCOM MPAs¹), adopted in 2014, is to establish an ecologically coherent and effectively managed network of coastal and marine Baltic Sea protected areas.

This report presents the recent development of the HELCOM MPA network, assesses the ecological coherence of the network, and follows up on other commitments made through HELCOM Recommendation 35/1. The assessment shows a continued positive development of HELCOM work on MPAs, but also room for improvement and the need for implementing the made agreements.

Since the designation of the first HELCOM MPAs in 1994, there has been a substantial increase in the areal coverage of MPAs: in 2004, the protected marine area of the Baltic Sea was 3.9%, in 2010 it was 10.3%, and today, in 2016, it is 11.8%. Thus, the target of conserving at least 10% of coastal and marine areas, set by the UN Convention on Biological Diversity, was reached already in 2010 in the Baltic Sea. Through Recommendation 35/1, HELCOM has furthermore agreed to reach the 10% target for each sub-basin, when scientifically justified. This target has been met in 11 out of 17 sub-basins, the exceptions being the Eastern and Western Gotland Basins, Northern Baltic Proper, Åland Sea, Bothnian Sea and Bothnian Bay.

At this time, the initial steep increase in the area of HELCOM MPAs has slowed down somewhat. However, recently 11 new Finnish sites were nominated as HELCOM MPAs, covering a total of 715 km² of marine area. 83% of this area is situated in the exclusive economic zone, which takes the network one step closer to the HELCOM target to include more offshore areas under the MPA protection regime.

The assessment of ecological coherence carried out for this report considered four aspects; representativity, replication, adequacy and connectivity. Two of these aspects were evaluated to be at an acceptable level for supporting a coherent MPA network: the areal representation of different types of geographical features and broad scale habitats, and the replication of a set of indicative species and biotope complexes, as well the broad scale habitats. However, evaluations of adequacy, which considers the quality of the network, and connectivity, which measures how well the network supports the migration and dispersal of species, indicate that the network is not yet ecologically coherent. Improving connectivity requires joint efforts from all HELCOM countries when planning and nominating new sites to the HELCOM MPA network, as connectivity cannot be improved on the level of single sites.

This report includes a first attempt at a quantitative approach for aggregating the results of the ecological coherence assessment. The quantitative aggregation indicates that it

¹ Former BSPAs.

is highly unlikely that the network of HELCOM MPAs is ecologically coherent. It proved to be a straightforward and transparent method and is recommended to be used in future HELCOM MPA assessments, alongside descriptive information on the status of the network. For the assessment of ecological coherence at large, further development is needed. Important basic information needed for appropriate analyses, for example on the spatial distribution of species and biotopes, is still missing, as well as science based targets for assessment criteria, such as replication.

Actions for reaching the objectives of Recommendation 35/1 are at this time in progress, but yet only partly accomplished. For example the goal to ensure that HELCOM MPAs provide specific protection to HELCOM Red Listed species, habitats, biotopes and biotope complexes has not been reached, since many threatened features are not protected in any of the HELCOM MPAs, at least not according to information reported by the Contracting Parties.

HELCOM Recommendation 35/1 also emphasizes the development and implementation of management plans for MPAs, as well as assessing the effectiveness of management plans, or other measures, to ensure protection. One of the commitments is to develop and apply management plans, or measures, for all existing HELCOM MPAs by 2015, and to establish a management plan, or measures, for every new MPA within five years after its designation. This agreement has not been met; currently only 67% of the HELCOM MPAs have management plans. Regarding the assessment of the effectiveness of the plans, this has not yet taken place and joint guidelines still remain to be developed on how to carry out such assessments. At present, monitoring within MPAs, a prerequisite for the assessment of effectiveness, occurs in 64% of HELCOM MPAs.

1 Introduction

1.1 Purpose of assessing marine protected areas in the Baltic Sea

The purpose of assessing marine protected areas (MPAs) is to follow up on the development of the MPA network in the Baltic Sea, to identify where further development of the network is needed, and to evaluate commitments made in HELCOM with regard to MPAs. The overarching target is to achieve a coherent and effectively managed network of MPAs in the Baltic Sea, including not only the network of HELCOM MPAs, but also other protection programmes, such as Natura 2000 and Ramsar sites.

Additional specific targets include, as agreed through Recommendation 35/1 on the system of coastal and marine Baltic Sea protected areas (HELCOM MPAs²) to;

- protect at least 10% of the marine area of each Baltic Sea sub-basin, when scientifically justified,
- designate new sites as HELCOM MPAs, where ecologically meaningful, especially in offshore areas beyond territorial waters,
- ensure that HELCOM MPAs provide specific protection to those species, habitats, biotopes and biotope complexes included in the HELCOM Red Lists,
- develop and apply management plans or measures for all existing HELCOM MPAs by 2015, and establish a management plan or measures for every new MPA within five years after its designation,
- assess the effectiveness of the management plans or measures of HELCOM MPAs by conducting monitoring, and, where feasible, scientific research programmes, which are directly connected to the conservation interests of HELCOM MPAs, including the placement of monitoring stations inside the MPAs,
- modernize the HELCOM MPA database, taking into account and harmonizing with other similar databases.

This report includes an assessment of the ecological coherence of the HELCOM MPA network (chapter 3), a proposal for a new aggregation method for summarizing the outcome of the ecological coherence (chapter 4) and a follow-up of the commitments of HELCOM Recommendation 35/1 (chapter 5). The basis of the assessment is data, information and shapefiles reported to the HELCOM MPA database, which was modernized and updated in 2015. The assessment methodology is based on the previous HELCOM assessment of ecological coherence (HELCOM 2010) and discussions within the HELCOM MPA Task Group and State and Conservation working group.

1.2 The concept of coastal and marine Baltic Sea protected areas (HELCOM MPAs²)

The main goal of the coastal and marine Baltic Sea protected areas (HELCOM MPAs) is to protect valuable marine and coastal habitats in the Baltic Sea. This is done by designating suitable areas which have particular nature values as protected areas, and by managing human activities within those areas (HELCOM 2003).

The first HELCOM MPAs were established in 1994, following the adoption of the 1992 Helsinki Convention, specifically its Article 15 on Nature conservation and biodiversity. Article 15 of the Helsinki Convention requires the Contracting Parties to take all appropri-

² Former BSPAs.

ate measures to conserve natural habitats and biological diversity in the Baltic Sea. To further implement Article 15, HELCOM Recommendation 15/5 on the system of coastal and marine Baltic Sea protected areas was adopted in 1994. Furthermore, HELCOM agreed upon guidelines and criteria for HELCOM MPAs, as presented in Box 1 (HELCOM 1994). At a later stage, the Baltic Sea Action Plan (HELCOM 2007a) and HELCOM 2010 and 2013 Ministerial Meetings agreed upon objectives for the network of protected areas, encouraging the Contracting Parties to nominate new areas (HELCOM 2007a). Recommendation 15/5 was reviewed and updated in 2014 and is now superseded by Recommendation 35/1 on the system of coastal and marine Baltic Sea protected areas (HELCOM MPAs). The new recommendation consolidates previous commitments and its main focus is on improving the management and ecological coherence of the HELCOM MPA network in the Baltic Sea (HELCOM 2014).

Initially, 62 sites from all nine riparian states were nominated as HELCOM MPAs. Today the number of designated sites is 174. HELCOM MPAs cover both marine and coastal areas, including islands. The status of the HELCOM MPA network has been reviewed on several occasions (HELCOM 2006a, Al-Hamdani & Reker 2007, Andersson et al. 2007, HELCOM 2007b, Piekäinen & Korpinen 2008, HELCOM 2009 and 2010), and this publication provides an overview of the status and ecological coherence of the network as reported in November 2015.

BOX 1. HELCOM objectives and criteria for the HELCOM MPA network ^{a)}

- | | |
|--|--|
| <ol style="list-style-type: none"> 1. A HELCOM MPA should give particular protection to the species, natural habitats and nature types in order to conserve biological and genetic diversity. 2. It should protect ecological processes and ensure ecological function. 3. It should enable the natural habitat types and the habitats of the species to be maintained at, or where appropriate, restored to a favourable conservation status in their natural range. 4. The minimum marine size of a HELCOM MPA should preferably be 30 km² for marine/lagoon parts. 5. The system should be enlarged stepwise by additional areas, preferably purely marine areas. | <ol style="list-style-type: none"> 6. The network should protect areas with: <ul style="list-style-type: none"> • threatened and/or declining species and habitats, • important species and habitats, • high natural biodiversity, • rare, unique, or representative geological or geomorphological structures or processes, • high sensitivity, • ecological significance: <ul style="list-style-type: none"> – a high proportion of habitats of migratory species, – important feeding, breeding, moulting, wintering or resting sites, – important nursery, juvenile or spawning areas, – a high natural biological productivity of the species or features being represented. |
|--|--|

^{a)} The objectives and criteria are based on the Joint HELCOM/ OSPAR Work Programme on Marine Protected Areas (Bremen 2003), HELCOM Recommendation 15/5 on the system of coastal and marine Baltic Sea protected areas and on the Minutes of the Eight Meeting of Nature Protection and Biodiversity Group (HELCOM HABITAT 8/2006).

2 Status of the network

Today there are 174 designated HELCOM MPAs (status in March 2016). They cover a total of 54 367 km², of which 90% (49 107 km²) is marine area (Table 1). The marine area of all HELCOM MPAs equals 11.8% of the total surface area of the Baltic Sea (Figure 1). Denmark protects the largest marine area measured in square kilometres (10 411 km²), while Germany protects the largest marine area measured as the percentage of the total area of the nation (36%). Sweden protects the largest area in the exclusive economic zone (2 750 km²).

The last overview of the HELCOM MPA network was carried out in 2013 (HELCOM 2013a). Since then, Finland has nominated 11 new MPAs and the total number of sites has grown from 163 to 174. This increased the total area of HELCOM MPAs by 725 km².

Table 1. Coverage of the HELCOM MPA network in 2016. Total number, total area and marine fraction of HELCOM MPAs, as well as total marine area and protected marine area per each country. The data is based on HELCOM MPA shapefiles from March 2016.

Country and number of HELCOM MPAs		Total area of HELCOM MPAs km ²	Marine fraction of HELCOM MPAs km ² (%)		Total marine area per country TW km ² EEZ km ² Total km ²			Size of protected marine area (km ²) and fraction (%) of the national territorial waters (TW), exclusive economic zone (EEZ) and total area					
								TW km ²	(%)	EEZ km ²	(%)	Total km ²	(%)
Denmark	66	11 223	10 436	(93%)	33 032	13 075	46 107	9 344	(28%)	1 092	(8%)	10 436	(23%)
Estonia	7	7 237	5 997	(83%)	25 084	11 840	36 923	5 954	(24%)	43	(0%)	5 997	(16%)
Finland	33	6 825	6 367	(93%)	52 622	28 696	81 318	6 285	(12%)	83	(0%)	6 367	(8%)
Germany	12	4 853	4 627	(95%)	11 041	4 504	15 544	2 158	(20%)	2 469	(55%)	4 627	(30%)
Latvia	7	4 367	4 363	(100%)	12 642	16 123	28 765	4 199	(33%)	164	(1%)	4 363	(15%)
Lithuania	6	1 387	992	(72%)	2 275	4 253	6 527	992	(44%)	0	(0%)	992	(15%)
Poland	9	7 939	7 230	(91%)	10 091	19 492	29 583	5 562	(55%)	1 668	(9%)	7 230	(24%)
Russia	6	1 339	894	(67%)	16 303	7 372	23 675	894	(5%)	0	(0%)	894	(4%)
Sweden	28	8 386	7 398	(88%)	77 281	71 402	148 683	4 649	(6%)	2 750	(4%)	7398	(5%)
Total	174	53 556	48 305	(90.2%)	240 370	176 755	417 125	40 037	(16.7%)	8 269	(4.7%)	48 305	(11.6%)

The current overview numbers (Table 1) are calculated from shapefiles, while the numbers published in 2013 were based both on shapefiles and background data reported by the countries. During the last years, the shapefiles have become more detailed, which directly affects the surface area calculated from them. Therefore the current numbers are not directly comparable with the ones published in 2013. To enable better comparison, the 2013 values have been recalculated, based on current shapefiles and adapted for the number of HELCOM MPAs in 2013 (Annex 1). In 2013, the protected marine area of the Baltic Sea was reported as 11.7% (HELCOM 2013a), but, according to calculations on the new shapefiles, it was 11.6%. Today the corresponding number is still 11.6% (Table 1).




Figure 1. HELCOM MPAs in the Baltic Sea, as reported by the HELCOM countries (status in March 2016).

2.1 The Natura 2000 network in the HELCOM area

The HELCOM MPA network overlaps with sites established under other frameworks, foremost the Natura 2000 network established under EU legislation (Figure 2). The Natura 2000 network aims to support the EU member states in achieving or maintaining a favourable conservation status for European biodiversity features in both terrestrial and marine habitats. It is based on the Birds Directive adopted in 1979 (Anonymous 1979, 2009) and the Habitats Directive adopted in 1992 (Anonymous 1992) and provides legal protection to the sites. Many Natura 2000 sites in the Baltic Sea have also been designated as HELCOM MPAs, and some smaller sites have been merged together under one large HELCOM MPA. Overlapping Natura 2000 sites and HELCOM MPAs often have

different shapes as the Natura 2000 sites may also include inland areas, while the HELCOM MPAs are restricted to the coastal zone and marine area. The Natura 2000 network protects certain natural habitats and species in the EU countries, whereas the HELCOM MPA network targets marine and coastal habitats and species specific to the Baltic Sea. In addition, the HELCOM MPA network also includes Russian waters in the Baltic Sea, while the Natura 2000 network is restricted to marine areas under EU jurisdiction. Today the HELCOM MPA network covers 48 184 km² of the Baltic Sea, while the Natura 2000 sites cover 45 688 km². According to the recently published European report on protected area coverage, the Baltic Sea has the highest protection of all European marine regions (EEA 2015, Technical Report No 17/2015).




Figure 2. Overlap of the marine Natura 2000 sites and the HELCOM MPAs in the Baltic Sea.

OSPAR (Convention for the Protection of the Marine Environment of the North-East Atlantic) is a regional seas convention for protecting and conserving the North-East Atlantic and its resources. HELCOM and OSPAR areas overlap in Kattegat, and the protected sites in this area are protected by both regional seas conventions at the same time. A joint target and work programme for these two networks was agreed upon at the HELCOM/OSPAR Bremen Ministerial Meeting in 2003.

3 Assessment of ecological coherence of the MPA network in the Baltic Sea

Ecological coherence describes how well a collection of MPAs provide protection to certain features, such as species, habitats, landscapes and ecological processes, both individually and as a network. When well planned and managed as a network, a collection of sites can deliver more benefits than unconnected individual MPAs can provide on their own (IUCN-WCPA 2008, UNEP-WCMC 2008, and Catchpole 2012). Ecological coherence is assessed by criteria, which describe different characteristics of the network, such as how well certain features are represented within the MPAs and how these MPAs are connected to each other.

This assessment is based on the same four main criteria as the previous HELCOM ecological coherence assessment (HELCOM 2010): representativity, replication, adequacy and connectivity. Each criterion was further divided into subcriteria, which were evaluated through spatial analysis (GIS) against specific targets (Table 2 and Annex 2).

This ecological coherence assessment was carried out in parallel for (1) the HELCOM MPA network, and (2) the combined network of the HELCOM MPAs and the marine Natura 2000 sites in the Baltic Sea. The HELCOM MPA network was assessed by all criteria and subcriteria, while the combined Baltic Sea network of HELCOM MPAs and marine Natura 2000 sites was only assessed by those criteria and subcriteria which were deemed applicable, as there were limitations due to fewer data available for the Natura 2000 sites (detailed information in Annex 3).

Table 2. Targets of the subcriteria of the ecological coherence assessment.

Criteria and subcriteria	Target	Reasoning or source
I. Representativity		
Benthic marine landscapes	<20% coverage = inadequate protection 20-60% coverage = adequate protection of common habitats (>60% coverage = adequate protection of rare habitats ³)	Piekäinen & Korpinen 2008, HELCOM 2010
Geographical representation	≥10% of the total Baltic Sea shall be protected, as well as each sub-basin and the coastal sea, outer coastal sea and open sea zones ⁴	CBD 2010 (Aichi 11 target), HELCOM 2010
II. Replication		
Marker species and biotope complexes	A minimum of 3 replicates within the HELCOM MPA network	Piekäinen & Korpinen 2008, HELCOM 2010
Benthic marine landscapes	A minimum of 3 replicates within the HELCOM MPA network	Piekäinen & Korpinen 2008, HELCOM 2010
III. Adequacy		
Marine size of MPAs	80% of marine sites ≥ 30 km ²	Recommended size for HELCOM MPAs, and decision taken by HELCOM STATE AND CONSERVATION 3-2015
Terrestrial size of MPAs	80% of terrestrial sites ≥ 10 km ²	Recommended size for HELCOM MPAs, and decision taken by HELCOM STATE AND CONSERVATION 3-2015
IV. Connectivity		
Theoretical connectivity of benthic marine landscapes	50% of landscape patches have ≥20 connections at the given dispersal distance	HELCOM 2010, Wolters et al. 2015
Species-specific connectivity	50% of landscape patches representing habitats for the species have ≥20 connections at the given dispersal distance	HELCOM 2010, Wolters et al. 2015


³ The 60% target originates from the previous HELCOM assessment (HELCOM 2010), which was inspired by the European guidelines for the application of the Habitats Directive in the marine environment (EC 2007).

⁴ The target of a minimum 10% coverage for geographical representativity is based on political decisions and not scientific studies, which usually refer to much higher percentages (25-30%) for protection goals (IUCN-WCPA 2008, UNEP-WCMC 2008).

3.1 Representativity

An assessment of representativity considers in broad terms different types of areal coverage, in order to determine whether the MPA network covers enough different features (for example species or biotopes) or factors linked to them (for example, suitable landscapes or areas for the species or biotopes). The basic assessment includes coverage of MPAs in the Baltic Sea and its sub-basins. In addition, representativity can be assessed for the coverage of conservation features, such as species, biotopes and landscapes.

In this assessment, representativity was evaluated by two subcriteria; (1) geographical representativity of MPAs (consisting of sub-basins and zonation categories of the Baltic Sea), and (2) benthic marine landscapes (Figures 3, 4, 5 and 6). The target for geographical representativity (1) was a minimum of 10% coverage of MPAs in the whole Baltic Sea, the sub-basins, the coastal sea, outer coastal sea and open sea zones. The target for benthic marine landscapes (2) was 20% coverage of each landscape within the MPA network (of the total landscape area in the Baltic Sea). For rare landscapes, it is proposed to use a stricter target of 60% coverage. This target was used in the previous HELCOM assessment (HELCOM 2010), which was inspired by the European guidelines for application of the Habitats Directive in the marine environment (EC 2007). The marine landscapes cover the entire marine area and requiring a 60% MPA coverage for all of them is not realistic. In this assessment, the measure of landscape scarcity was not included in the representativity analyses, but the stricter target is included in Figures 9 and 12, in order to compare benthic marine landscapes to this level. However, 20% remains the main target for representativity of benthic marine landscapes in this assessment (Table 2).



* excluded due to lack of comprehensive distribution data of species and biotope complexes in the entire Baltic Sea.

**the 60% target originates from discussions under the Habitats Directive, and was originally intended only for rare habitats.

Figure 3. Overview of the assessment design of the representativity criterion of the ecological coherence assessment. The third subcriterion for marker species and biotope complexes was not used in the current assessment, due to a lack of data.

In an ideal case, the representativity of the Baltic Sea MPAs should also be assessed for species and biotopes complexes. However, the representativity assessment requires spatial data on the distribution of species and biotope complexes in the entire Baltic Sea, both inside and outside the MPAs, and such data is currently not available (please see chapter 3.5.1 for further details).

3.1.1 Method and results of the representativity assessment of the HELCOM MPA network

Representativity for both subcriteria was calculated as the percentage of the total area of the protected feature in the Baltic Sea occurring within the MPA network. The calculations were limited to the natural distribution range of the features. Data used for these analyses included shapefiles of the Baltic Sea sub-basins (Figure 4), coastal zones, benthic marine landscapes and the HELCOM MPAs (see Annex 4 for an overview table of all data).




Figure 4. The 17 sub-basins of the Baltic Sea according to the HELCOM Monitoring and Assessment Strategy (HELCOM 2013d).

The coastal zonation was created for this analysis at fixed distances from the countries' baselines ((1) coastal sea: <1 nm, (2) outer coastal sea: 1-12 nm, and (3) open sea: >12 nm) (Figure 5). The benthic marine landscape categories were derived from maps developed in the EUSeaMap project, based on light, bottom substrate and salinity in the Baltic Sea (EUSeaMap 2015). For this assessment, the original 60 categories were combined into 30 categories, based on salinity values and bottom substrata (Table 3). Figure 6 depicts the distribution of these categories in the Baltic Sea and their relative size. In this assessment the benthic marine landscapes have been used as a proxy for diversity, and it should be noted that some of these landscapes include more biological features than others.




Figure 5. The coastal zones of the Baltic Sea: 1) coastal sea (light green): from the coastline to 1 nm beyond the baseline, 2) outer coastal sea (light blue): 1-12 nm beyond the baseline, and 3) open sea (blue): >12 nm beyond the baseline.




Figure 6. The benthic marine landscapes of the Baltic Sea. The 30 landscape categories were derived from maps developed by the EUSeaMap project based on light, bottom substrate and salinity in the Baltic Sea. The diagram shows the area of the different landscapes in square kilometres.

Table 3. The basis for categorization of benthic marine landscapes. All possible combinations of the light, substrate and salinity classes below produce the 30 benthic marine landscape categories used in this assessment (please see Figure 6 for their distribution).

Light	Bottom substrate	Salinity
Photic	Sand	< 5
Aphotic	Rock and other hard substrata	5-18
	Mud	>18
	Mixed sediment	
	Coarse sediment	

The representativity target for the coverage of MPAs within the Baltic Sea, its sub-basins and the coastal zones (subcriteria 1) was a minimum 10% of the total area of each category. The target was reached for the Baltic Sea (12% coverage of MPAs), and for 11 of the 17 sub-basins (Figure 7). The target was also reached in two out of three zonation categories; coastal and outer coastal sea areas (Figure 8). The 20% protection target for landscapes (subcriteria 2) was reached for 18 of 30 landscapes (60%). The stricter target of 60% protection (intended for rare landscapes) was not reached by any landscape (Figure 9). Please note that the area of the different landscape varies between 14 and 126 395 km² (Figure 6).




Figure 7. Coverage of HELCOM MPAs in each sub-basin of the Baltic Sea. The HELCOM MPAs cover 12% of the entire Baltic Sea. The values were calculated as the area covered by HELCOM MPAs of the total area of the sub-basin, based on shapefiles of the MPAs provided by the HELCOM countries. The target (red line) is 10% coverage in each sub-basin.




Figure 8. Coverage of HELCOM MPAs in the Baltic Sea zones. The zones are 1) coastal sea: from the coastline to 1 nm beyond the baseline, 2) outer coastal sea: 1-12 nm beyond the baseline, and 3) open sea: >12 nm beyond the baseline (see Figure 5). The values were calculated as the area covered by HELCOM MPAs of the total area of the zone, based on shapefiles of MPAs provided by the HELCOM countries. The target (red line) is 10% coverage in each zone.




Figure 9. Coverage of benthic marine landscapes within the HELCOM MPA network. The values were calculated as the percentage protected by the HELCOM MPA network of the total landscape area, based on shapefiles of MPAs provided by the HELCOM countries. The percentage of the landscape in the Baltic Sea is given in brackets at the end of each bar (also see the diagram in figure 6). The target is 20% coverage for protecting common landscapes (solid red line) and 60% coverage for protecting rare landscapes (dotted red line).

SUMMARY OF THE RESULTS OF THE REPRESENTATIVITY ASSESSMENT OF THE HELCOM MPA NETWORK

The network of HELCOM MPAs meets some of the targets set for representativity in this assessment. The minimum target of 10% is met for the Baltic Sea as a whole, but not for all sub-basins or the open sea area (subcriteria 1). Representation does not reach 10% in the Eastern and Western Gotland basin, the northern Baltic Proper, the Åland Sea, the Bothnian Sea and the Bothnian Bay. The minimum target for protecting landscapes (20% coverage) is not met for 40% of the landscapes, which mainly includes aphotic landscapes occurring largely in the open sea area. The 60% protection target intended for rare landscapes is not met by any landscape (subcriteria 2).

3.1.2 Method and results of the representativity assessment of the combined network of HELCOM MPAs and marine Natura 2000 sites

Representativity was also assessed for the combined network of HELCOM MPAs and marine Natura 2000 sites by both subcriteria (geographical representation and benthic marine landscapes, Figure 3). For this analysis, a shapefile consisting of marine HELCOM MPAs and marine Natura 2000 areas in the Baltic Sea was constructed. The combined area of the Natura 2000 sites and HELCOM MPAs is 8127 km² larger than the HELCOM MPA network. Apart from this, the method and data were identical to the assessment of the HELCOM MPA network (chapter 3.1.1).

The representativity target for the coverage of MPAs within the Baltic Sea, its sub-basins and the coastal zones (subcriteria 1) was a minimum 10% of the total area of each category. The target was reached for the Baltic Sea with 13.5% coverage of the combined area of marine Natura 2000 sites and HELCOM MPAs, and for 11 of the 17 sub-basins (Figure 10). The target was also reached in two out of three zonation categories; coastal and outer coastal sea areas (Figure 11).

The 20% protection target for landscapes (subcriteria 2) was reached for 22 out of 30 landscapes, i.e. 73%. The stricter target of 60% protection (intended for rare landscapes) was




Figure 10. Coverage of marine Natura 2000 sites and HELCOM MPAs in each sub-basin of the Baltic Sea. In the whole Baltic Sea, these MPAs cover 13.5% of the marine area. The values were calculated as the area covered by marine Natura 2000 sites and HELCOM MPAs of the total area of the sub-basin, based on shapefiles of the HELCOM MPAs provided by the HELCOM countries and shapefiles of the Natura 2000 sites downloaded from the EEA Data & Map service. The target (red line) is 10% coverage in each sub-basin.




Figure 11. Coverage of marine Natura 2000 sites and HELCOM MPAs in the Baltic Sea zones. The zones are 1) coastal sea: from the coastline to 1 nm beyond the baseline, 2) outer coastal sea: 1-12 nm beyond the baseline, and 3) open sea: >12 nm beyond the baseline (see Figure 5). The values were calculated as the area covered by marine Natura 2000 sites and HELCOM MPAs of the total area of the zone, based on shapefiles of HELCOM MPAs provided by the HELCOM countries and shapefiles of the Natura 2000 sites downloaded from the EEA Data & Map service. The target (red line) is 10% coverage in each zone.

reached for nine landscapes. These are all scarce landscapes with a coverage less than 3.5% in the whole Baltic Sea (Figure 12). Almost all of the landscape *photic sand 5-18 psu* in the Baltic Sea is covered by marine Natura 2000 sites and HELCOM MPAs (99%). Please note that the areas of the different landscapes vary between 14 and 126 395 km² (Figure 6).




Figure 12. Coverage of benthic marine landscapes within the combined area of the marine Natura 2000 sites and HELCOM MPA network. The values were calculated as the area covered by the combined network of the total area of the landscape within the Baltic Sea, based on shapefiles of MPAs provided by the HELCOM countries and shapefiles of the Natura 2000 sites downloaded from the EEA Data & Map service. The total area of the landscape in the Baltic Sea is given as a separate percentage at the end of each bar (also see the diagram in figure 6). The target is 20% coverage for protecting common landscapes (solid red line) and 60% coverage for protecting rare landscapes (dotted red line).

SUMMARY OF THE RESULTS OF THE REPRESENTATIVITY ASSESSMENT OF THE COMBINED NETWORK OF HELCOM MPAS AND MARINE NATURA 2000 IN THE BALTIC SEA

The combined network of the HELCOM MPAs and the marine Natura 2000 sites gives better protection than the HELCOM MPA network alone, but the difference for the geographical distribution is small. The minimum target of 10% coverage is met for the Baltic Sea, but not for all sub-basins or the open sea area (subcriteria 1). The minimum target for protecting landscapes (20% coverage) is not met for 23% of the landscapes (mainly aphotic landscapes occurring largely in the open sea area), compared to 40% when analysing the HELCOM MPA network alone (subcriteria 2). However, the protection of certain landscapes, such as *photic sand 5-18 psu*, improves remarkably when the Natura 2000 network and the HELCOM MPAs are evaluated together, and nine landscapes reach the stricter 60% protection target. The HELCOM MPA network covers 53% of this landscape, while the combined network covers 99% of the landscape. This is mainly explained by the fragmented distribution and coastal location of the Natura 2000 network, as the *photic sand 5-18 psu* is a typical coastal landscape.

3.2 Replication

An assessment of replication considers the number of replicas of a conservation feature in the assessment area, for example certain species, and aims to ensure that the provided protection does not depend on only one site in the network.

In this assessment, replication was assessed by two subcriteria; (1) marker species and biotope complexes and (2) benthic marine landscapes (Figure 13). The target was set at a theoretical minimum of three replicas, based on the previous HELCOM assessment (HELCOM 2010 and references therein). The replication number equals the total number of occurrences minus one, which means that in total four occurrences are needed to reach the target of three replications. It should be noted that the chosen target is static and does not work equally well for all species and biotope complexes. In an ideal situation, the target would be adjusted for each species, biotope complex and landscape, based on scientific background information and on how rare or common they are.




Figure 13. Overview of the assessment design of replication criterion of the ecological coherence assessment.

3.2.1 Method and results of the replication assessment of the HELCOM MPA network

Replication values for marker species and biotope complexes were calculated as the number of HELCOM MPAs where the protected feature occurred, and calculations were limited to the natural distribution range of the features. Replication values for benthic marine landscapes were calculated as the total number of landscape patches within the HELCOM MPA network. Data used for these analyses included shapefiles of species distribution (reported per sub-basin), biotope complexes distribution (reported on a 100x100 km grid), benthic marine landscapes and the MPAs, as well as data on the presence of species and biotope complexes reported in the HELCOM MPA database (Annex 4). The benthic marine landscape classes were derived from maps developed in the EuSeaMap project, which were further developed by combining the existing categories to create the 30 categories as described in Table 3 and Figure 6 (EuSeaMap 2015).

Twelve marker species and eleven marker biotope complexes were chosen, based on the species and biotope complexes used in the previous assessment (HELCOM 2010) and discussions by the HELCOM MPA Task Group and the third meeting of the State and Conservation working group (STATE & CONSERVATION 3-2015) (Tables 4 and 5). The list of marker species used in the previous assessment was modified to ensure that all parts of the Baltic Sea were covered by the natural distributions of at least some of the chosen species. To achieve this, functionally similar species were paired together to form a marker

species pair with a distribution covering the whole Baltic Sea. This was done for four bird species and two mammal species, in order to form three species pairs.

The list of marker biotope complexes used in the previous assessment was expanded to include all marine biotope complexes of Annex I of the Habitat Directive which have been included in the MPA database (Table 5). One biotope complex ('macrophytes') was removed as the HELCOM MPA database no longer contains data on this biotope.

Table 4. The twelve marker species or species pairs used in the replication analysis.

Species group	Species/Species pair	English name of species or species pair	Distribution
Macrophytes	<i>Chara</i> spp.	Stoneworts	Whole Baltic Sea (HELCOM 2012).
Fish	<i>Anguilla anguilla</i>	European eel	Whole Baltic Sea, except the Bothnian Bay and the Quark (where the occurrence is temporary) (HELCOM 2012, 2013c).
Fish	<i>Lampetra fluviatilis</i>	River lamprey	Whole Baltic Sea, except the Belt Seas (where the occurrence is temporary) (HELCOM 2012, 2013c).
Fish	<i>Salmo salar</i>	Atlantic salmon	Whole Baltic Sea (HELCOM 2012, 2013c).
Birds	<i>Gavia arctica</i> & <i>Gavia stellata</i>	Black-throated diver & Red-throated diver	Combined wintering distribution of these species covers the whole Baltic Sea (HELCOM 2013c, BirdLife International 2015). ⁱ
Birds	<i>Mergus serrator</i> & <i>Mergus merganser</i>	Red breasted merganser & Goosander	Combined distribution of these species covers the whole Baltic Sea (HELCOM 2012).
Birds	<i>Sternula albifrons</i>	Little tern	Whole Baltic Sea (HELCOM 2012, 2013c).
Birds	<i>Tadorna tadorna</i>	Common Shelduck	Whole Baltic Sea (HELCOM 2012, 2013c).
Birds	<i>Somateria mollissima</i>	Common Eider	Whole Baltic Sea (combined wintering and breeding) (HELCOM 2012, 2013c, BirdLife International 2015).
Mammals	<i>Halichoerus grypus</i>	Grey seal	Whole Baltic Sea, except the Belt Seas (HELCOM 2012).
Mammals	<i>Phoca vitulina</i> & <i>Phoca hispida botnica</i>	Harbour seal & Ringed seal	Combined distribution of these species covers the whole Baltic Sea (HELCOM 2013c).
Mammals	<i>Phocoena phocoena</i>	Harbour porpoise	Southern and western Baltic Sea (i.e. Western Gotland basin, Bornholm basin, Arkona Basin, Bay of Mecklenburg basin, Kiel Bay, the Belts, the Sound and Kattegat) (HELCOM 2013c).

ⁱ Expert opinion.

Table 5. The eleven marker biotope complexes used in the replication analysis.

Biotope complex code	Biotope complex name	Distribution
1110	Sandbanks which are slightly covered by sea water all the time	Whole Baltic Sea (HELCOM 2013c).
1130	Estuaries	Whole Baltic Sea (HELCOM 2013c).
1140	Mudflats and sandflats not covered by seawater at low tide	Whole Baltic Sea (HELCOM 2013c).
1150	Coastal lagoons	Whole Baltic Sea (HELCOM 2013c).
1160	Large shallow inlets and bays	Whole Baltic Sea (HELCOM 2013c).
1170	Reefs	Whole Baltic Sea (HELCOM 2013c).
1180	Submarine structures made by leaking gases	Kattegat (HELCOM 2013c).
1610	Baltic esker islands with sandy, rocky and shingle beach vegetation and sublittoral vegetation	Bothnian Bay, Quark, Bothnian Sea, Åland Sea, Archipelago Sea, Gulf of Finland, Northern Baltic Proper, Western Gotland Basin (HELCOM 2013c).
1620	Boreal Baltic islets and small islands	Bothnian Bay, Quark, Bothnian Sea, Åland Sea, Archipelago Sea, Gulf of Finland, Northern Baltic Proper, Western Gotland Basin, Bornholm basin (HELCOM 2013c).
1650	Boreal Baltic narrow inlets	Bothnian Bay, Bothnian Sea, Archipelago Sea, Gulf of Finland, Northern Baltic Proper, Western Gotland Basin (HELCOM 2013c).
8330	Submerged or partially submerged sea caves	No distribution data available.

The replication target for marker species and biotope complexes (subcriteria 1), as well as the landscapes (subcriteria 2), within the HELCOM MPA network was set at a minimum of 3 replications. The target was reached for all marker species (Figure 14) and landscapes (Figure 16), and for 9 of 11 (82%) marker biotope complexes (Figure 15).




Figure 14. Replication of marker species within the HELCOM MPA network. The values were calculated as the number of HELCOM MPAs in which the species is present, within its natural distribution range, based on data reported in the HELCOM MPA database and shapefiles on species distribution from the HELCOM Species Information Sheets (HELCOM 2013e). The target of at least three replicates is reached when the species is present in at least four HELCOM MPAs (red line).




Figure 15. Replication of marker biotope complexes within the MPA network. The values were calculated as the number of MPAs in which the biotope complex occurs within its natural distribution range, based on data reported to the HELCOM MPA database and shapefiles on biotope complex distribution from the HELCOM Biotope Complex Information Sheets (HELCOM 2013f). The target of at least three replicates is reached when the biotope complex occurs in a minimum of four MPAs (red line).




Figure 16. Replication of landscapes within the HELCOM MPA network. The values were calculated as the total number of landscape patches within the HELCOM MPA network, based on the shapefile of landscapes (Figure 6) and shapefiles of MPAs. The target of at least three replicates is reached when the landscape occurs in a minimum of four MPAs (red line).

SUMMARY OF THE RESULTS OF THE REPLICATION ASSESSMENT OF THE HELCOM MPA NETWORK

The network of HELCOM MPAs meets almost all of the targets set for replication in this assessment. Only two biotope complexes (1650 Boreal Baltic narrow inlets and 8830 Submerged or partially submerged sea caves) did not reach the target of 3 replications within the HELCOM MPA network.

3.2.2 Method and results of the replication assessment of the combined network of HELCOM MPAs and marine Natura 2000 sites

Replication was also assessed for the combined network of HELCOM MPAs and marine Natura 2000 sites, but only by the second subcriteria: benthic marine landscapes (Figure 13), since data for subcriteria 1 (marker species and biotope complexes) was not readily available for the Natura 2000 sites. For this analysis, a shapefile consisting of marine HELCOM MPAs and marine Natura 2000 areas in the Baltic Sea was constructed. The combined area of the Natura 2000 sites and HELCOM MPAs is 8127 km² larger than the HELCOM MPA network. Apart from this, the method and data were identical with the assessment of the HELCOM MPA network alone (chapter 3.1.2).

The replication target for the landscapes (subcriteria 2) within the combined network of marine Natura 2000 sites and HELCOM MPAs was set at a minimum of 3 replicas of each landscape, and it was reached for all landscapes (Figure 17).




Figure 17. Replication of landscapes within the marine Natura 2000 sites and HELCOM MPA network. The values were calculated as the total number of landscape patches within the combined area of the marine Natura 2000 sites and the HELCOM MPA network, based on the shapefile of landscapes (Figure 6), shapefiles of MPAs provided by the HELCOM countries and shapefiles of the Natura 2000 sites downloaded from the EEA Data & Map service. The target of at least three replicates is reached when the landscape occurs in a minimum of four MPAs (red line).

SUMMARY OF THE RESULTS OF THE REPLICATION ASSESSMENT OF THE COMBINED MARINE NATURA 2000 SITES AND HELCOM MPA NETWORK IN THE BALTIC SEA

The combined network of the HELCOM MPAs and the marine Natura 2000 sites ensures better replication of the landscapes than the HELCOM MPA network alone.

3.3 Adequacy

Adequacy is a concept which describes the quality aspects of single MPAs. While the other three main criteria used in this assessment evaluate the MPAs as a network, adequacy focuses on evaluating whether the single MPAs are sufficient as building blocks for the network. Adequacy has commonly been assessed based on the size, shape or location of the MPAs, but also based on the pressures affecting the MPAs and the level of protection provided by the MPAs (HELCOM 2010, Wolters et al. 2015, CBD 2008). According to the CBD adequacy definition, MPA networks should include a core system of no-take areas, a larger system of multiple-use MPAs, as well as areas of sustainable use (CBD, 2008).

In this assessment, the evaluation of adequacy was restricted to one subcriterion where clear targets could be set: MPA size (Figure 18). The targets were based on the recommended size of 30 km² (3000 ha) for marine areas and 10 km² (1000 ha) for terrestrial areas of HELCOM MPAs (HELCOM 2003). This target was defined to apply when 80% of the MPAs have reached the recommended size, thus also acknowledging that smaller MPAs may be relevant components of the network (Table 2).




Figure 18. Overview of the assessment design of adequacy criterion of the ecological coherence assessment.

In addition to the one subcriterion, data on pressures and protection level were analysed as supporting information. No targets have been set within HELCOM for either pressures or protection level of the MPAs, and therefore the results of these analyses were only used to provide descriptive information on the MPAs. Pressures were analysed based on the spatial distribution of fishing activity and ship traffic. Fishing activity was represented by the total hours of fishing intensity (Vessel Monitoring System (VMS) data on fishing effort, all gear types) within C-square cells during 2013, and ship traffic (all ship types) during 2014 in a 500 m grid based on HELCOM Automatic Identification System (AIS) within the MPAs. The level of protection was analysed as the percentage of MPAs which have been assigned the most strictly protected levels of the IUCN protection categories, i.e. categories I-II (reflecting the CBD definition) (Table 6).

3.1.1. Method and results of the adequacy assessment of the HELCOM MPA network

The size of the marine and terrestrial areas of each MPA were calculated from MPA shapefiles submitted by the HELCOM countries. Neither marine nor terrestrial areas reached the 80% target, as 68% of marine parts of MPAs were at least 30 km² large and 40% of the terrestrial parts of the MPAs were at least 10 km² large (Figure 19). It can be noted that a majority of MPAs are sized between 100-1000 km² (Figure 20).




Figure 19. The percent of MPAs reaching the given size (30 km² for marine areas and 10 km² for terrestrial areas of MPAs). The values were calculated based on the shapefiles of MPAs provided by the HELCOM countries. The target is reached when 80% of the MPAs reach the recommended size (red line).




Figure 20. Distribution of MPAs into size classes, according to the size of their marine area. The values were calculated based on the shapefiles of MPAs provided by the HELCOM countries.

Supporting analyses of pressures and protection level of the HELCOM MPAs

Supporting analyses related to fishing, which was carried out by ICES, showed that intensive fishing activities are occurring in some HELCOM MPAs, especially in the southern and western Baltic Sea (Figure 21). In total, 11 630 hours of fishing took place within 69 HELCOM MPAs in 2013, over half of this (70%) by bottom contacting gear and about a fourth (26%) by midwater trawls. Only a small part of fishing (4%) was done by longlines and only in 7 MPAs (ICES 2015 and Annex 5).

Pressures caused by ship traffic, such as input of litter and contaminants, disturbance of species and introduction or spread of non-indigenous species occur within many HELCOM MPAs, and in some areas major ship lines go through or very close to an MPA (Figure 22).

In total, 41 (24%) of all 174 HELCOM MPAs are protected under at least one of the IUCN categories for strict protection (Ia Strict Nature Reserve, Ib Wilderness Area and II National Park) (Figure 23 and Table 6). This lies within the range of 10-30% strict protection, which was recommended by the Fifth World Parks Congress (2003) for each habitat type. This recommendation is however based on areal coverage, which is not possible to analyse with the current data, as information on the areal coverage of the different categories is lacking. The most common IUCN category assigned to an MPA (or a part of it) is the category IV Habitat/Species Management Area.




Figure 21. HELCOM MPAs and spatial distribution of total hours of fishing effort (Vessel Monitoring System data on fishing effort, all gear types) within C-square cells during the year 2013 (ICES 2015) in the Baltic Sea (excluding Skagerrak). Please note that currently no data is available for the white areas on the map.





Figure 22. HELCOM MPAs and ship traffic (all ship types) during 2014, reported on a 1,000 m grid based on the HELCOM Automatic Identification System (AIS).



IUCN categories for protected areas

Figure 23. Number of HELCOM MPAs assigned to the different IUCN categories. The IUCN category is provided for 118 of all 174 (68%) HELCOM MPAs, and in many cases, one MPA has been assigned several IUCN categories. In total, 41 HELCOM MPAs have been assigned to one or several of the strict protection IUCN categories (Ia-II).

Table 6. Categorization and definitions of protected areas according to IUCN.

Code	Name	Definition
Ia	Strict Nature Reserve	Category Ia are strictly protected areas set aside to protect biodiversity and also possibly geological/geomorphical features, where human visitation, use and impacts are strictly controlled and limited to ensure protection of the conservation values. Such protected areas can serve as indispensable reference areas for scientific research and monitoring.
Ib	Wilderness Area	Category Ib protected areas are usually large unmodified or slightly modified areas, retaining their natural character and influence without permanent or significant human habitation, which are protected and managed so as to preserve their natural condition.
II	National Park	Category II protected areas are large natural or near natural areas set aside to protect large-scale ecological processes, along with the complement of species and ecosystems characteristic of the area, which also provide a foundation for environmentally and culturally compatible, spiritual, scientific, educational, recreational, and visitor opportunities.
III	Natural Monument or Feature	Category III protected areas are set aside to protect a specific natural monument, which can be a landform, sea mount, submarine cavern, geological feature such as a cave or even a living feature such as an ancient grove. They are generally quite small protected areas and often have high visitor value.
IV	Habitat/Species Management Area	Category IV protected areas aim to protect particular species or habitats and management reflects this priority. Many Category IV protected areas will need regular, active interventions to address the requirements of particular species or to maintain habitats, but this is not a requirement of the category.
V	Protected Landscape/Seascape	A protected area where the interaction of people and nature over time has produced an area of distinct character with significant, ecological, biological, cultural and scenic value: and where safeguarding the integrity of this interaction is vital to protecting and sustaining the area and its associated nature conservation and other values.
VI	Protected area with sustainable use of natural resources	Category VI protected areas conserve ecosystems and habitats together with associated cultural values and traditional natural resource management systems. They are generally large, with most of the area in a natural condition, where a proportion is under sustainable natural resource management and where low-level non-industrial use of natural resources compatible with nature conservation is seen as one of the main aims of the area.

SUMMARY OF THE RESULTS OF THE ADEQUACY ASSESSMENT OF THE HELCOM MPA NETWORK

The network of HELCOM MPAs does not meet the targets set for adequacy in this assessment.

The supporting information reveals that fishing and shipping occurs in many HELCOM MPAs, but further analyses are needed to see if this contradicts with the protection targets of the specific HELCOM MPAs. The number of HELCOM MPAs protected under the stricter IUCN protection categories seems to be sufficient, but to verify this result analyses should be done based on areal coverage instead of the number of MPAs.

3.3.2 Method and results of the adequacy assessment of the combined network of HELCOM MPAs and marine Natura 2000 sites

This analysis was not carried out for the Natura 2000 sites, as there is no minimum size recommendations for these sites. In general, Natura 2000 sites tend to be smaller than HELCOM MPAs.

3.4 Connectivity

Connectivity of MPAs has been called the ‘glue of the network’, and it measures whether a group of MPAs function as a network. Connectivity aims to ensure that species’ migrations and dispersals during different life stages is supported by the MPA network.

Connectivity was assessed by two subcriteria; (1) theoretical connectivity and (2) species-specific connectivity (Figure 24). The target for both subcriteria was that 50% of landscape patches would have ≥ 20 connections.




Figure 24. Overview of the assessment design of the connectivity criterion of the ecological coherence assessment.

3.4.1 Method and results of connectivity assessment of the HELCOM MPA network

Connectivity for both subcriteria was calculated as the number of connections within the given dispersal distance between specific landscape patches in the MPAs. Calculations were limited to the natural distribution range of the features. Data used for the analyses included shapefiles on benthic marine landscapes and the MPAs (see Annex 4 for an overview table of all data). The benthic marine landscape categories are described in chapter 3.1.

Theoretical connectivity (subcriteria 1) for (i) 25 km and (ii) 50 km dispersal distance was analysed for five landscapes chosen to correspond to the ones used in the previous HELCOM ecological coherence assessment (HELCOM 2010): photic mud <5 psu, photic sand 5-18 psu, aphotic rock and other hard substrata <5 psu, aphotic sand 5-18 psu, and aphotic mixed substrate >18 psu. The patches of each landscape (min size 0.24 km^2) were buffered with half the dispersal distance, and the number of buffers touching or overlapping equalled the number of connections. The target for theoretical connectivity was set at 50% of the landscape patches having ≥ 20 connections at the given dispersal distance (i) 25 and (ii) 50 km).

Species-specific connectivity (subcriteria 2) was analysed based on landscapes representing five species with differing dispersal strategies and distances (Table 7). The same five species were used in the previous HELCOM ecological coherence assessment (HELCOM 2010). The analysis was carried out as for subcriteria 1, this time choosing several landscapes into each analysis to cover all probable habitats occupied by the species, as reported in literature. The target for species-specific connectivity was that 50% of the landscape patches representing habitats for the species should have ≥ 20 connections at the species-specific dispersal distance.

Table 7. The species used in the analysis of species-specific connectivity.

Species	Substrate	Salinity	Photic depth	Dispersal distance	Notes and references
<i>Macoma baltica</i>	Sand and mud ⁱ	> 5 psu	Non-photic and photic	100 km ⁱⁱ	Tolerates salinity of 4 psu ⁱⁱⁱ . Distribution whole Baltic Sea, except the Bothnian Bay.
<i>Psetta maxima</i> spawning and nursery grounds	Bedrock, hard bottom complex and sand ^{iv}	> 5 psu	Photic	25 km ^v	Spawning and nursery grounds are not found north of the Finnish south coast.
<i>Furcellaria lumbricalis</i>	Bedrock, hard bottom complex and sand	> 5 psu	Photic	25 km ^{vi}	Distribution whole Baltic Sea, except the Bothnian Bay.
<i>Idotea baltica</i>	Bedrock, hard bottom complex and sand	> 5 psu	Photic	25 km ^{vii}	Distribution whole Baltic Sea, except the Bothnian Bay.
<i>Fucus vesiculosus</i>	Bedrock, hard bottom complex and sand	> 5 psu	Photic	1 km ^{viii}	Distribution whole Baltic Sea, except the Bothnian Bay.

i MarLIN,

ii larval settling time 1-6 months, Marlin,

iii Laine & Seppänen 2001,

iv Iglesias et al. 2003, Sparrevohn & Sottrup 2003, Stankus 2006,

v based on genetical studies, Florin & Höglund 2006,

vi Fletcher & Callow 1992, Norton 1992,

vii based on measurements by Alexander & Chen 1990,

viii according to Gaylord et al. 2002: a fraction of algal propagules can drift distances of several kilometres.

The theoretical connectivity (1) target for 50% of the landscape patches having ≥ 20 connections was not reached for the 25 km dispersal distance, nor for the 50 km dispersal distance (Figure 25). The species-specific connectivity (2) target for 50% of the landscape patches having ≥ 20 connections was reached for the dispersal distance of *Macoma baltica*, *Psetta maxima*, *Furcellaria lumbricalis* and *Idotea baltica*, but not for *Fucus vesiculosus* (Figure 26). This is explained by the very short dispersal distance of *F. vesiculosus* (only 1 km). The results correlate with the dispersal distance of the species; *M. baltica* with the longest dispersal distance (100 km) has the highest connectivity value (96% of species-specific habitat patches with a minimum of 20 connections).




Figure 25. Theoretical connectivity of landscape patches within the HELCOM MPA network. The values were calculated as the number of connections between landscape patches of the same type within the HELCOM MPAs for five landscapes, based on the shapefile of landscapes (Figure 5) and shapefiles of MPAs provided by the HELCOM countries. The target of at least 50% of all patches having 20 or more connections lies outside of the range of this figure.




Figure 26. Species-specific connectivity of landscape patches representing habitats for five species within the MPA network. The chart shows the percentage of patches with 20 or more connections. The values were calculated as the number of connections between landscape patches chosen for a species within the HELCOM MPAs for five species, based on the shapefile of landscapes (Figure 5) and shapefiles of MPAs provided by the HELCOM countries. The target is reached when 50% of the patches have at least 20 connections (red line).

SUMMARY OF THE RESULTS OF THE CONNECTIVITY ASSESSMENT OF THE HELCOM MPA NETWORK

The network of HELCOM MPAs meets the targets set for species-specific connectivity (subcriteria 2) in this assessment in four of five cases, but in none for theoretical connectivity (subcriteria 1). The minimum target of 20 connections for at least 50% of the landscape patches representing habitats for a chosen species is met for all species in the analysis except *Fucus vesiculosus* (subcriteria 2). This is explained by the very short dispersal distance of *F. vesiculosus* (only 1 km).

3.4.2 Method and results of connectivity of the combined network of HELCOM MPAs and marine Natura 2000 sites

Connectivity was also assessed for the combined network of HELCOM MPAs and marine Natura 2000 sites by both subcriteria (theoretical and species-specific connectivity) (Figure 24). For this analysis, a shapefile consisting of marine HELCOM MPAs and marine Natura 2000 areas in the Baltic Sea was constructed. The combined area of the Natura 2000 sites and HELCOM MPAs is 8127 km² larger than the HELCOM MPA network. Apart from this, the method and data were identical with the assessment of the HELCOM MPA network (chapter 3.4.1).

The same targets were used; 50% of the landscape patches should have ≥20 connections at the given dispersal distance, which for theoretical connectivity was (i) 25 and (ii) 50 km, and for species-specific connectivity followed the species-specific dispersal distance (Table 7).

The theoretical connectivity target of 50% of the landscape patches having ≥20 connections was not reached for the 25 km dispersal distance, nor for the 50 km dispersal distance (Figure 27) in the combined area of the HELCOM MPAs and marine Natura 2000 sites. The values are, however, higher than for the HELCOM MPA network alone. The species-specific connectivity target for 50% of the landscape patches having ≥20 connections was reached for the dispersal distance of *Macoma baltica*, *Psetta maxima*, *Furcellaria lumbricalis* and *Idotea baltica*, but not for *Fucus vesiculosus* (Figure 28). This is explained by the very short dispersal distance of *F. vesiculosus* (only 1 km). Also in this analysis, the results correlate with the dispersal distance of the species; *M. baltica* with the longest




Figure 27. Theoretical connectivity of landscape patches within the marine Natura 2000 sites and HELCOM MPA network. The values were calculated as the number of connections between landscape patches of the same type within the combined area of marine Natura 2000 sites and HELCOM MPAs for five landscapes, based on the shapefile of landscapes (Figure 6), shapefiles of MPAs provided by the HELCOM countries and shapefiles of the Natura 2000 sites downloaded from the EEA Data & Map service. The target of at least 50% of all patches having 20 or more connections lies outside of the range of this figure.

dispersal distance (100 km) has the highest connectivity value (95% of species-specific habitat patches with a minimum of 20 connections). Interestingly, the overall results for this analysis are slightly lower than for the HELCOM MPA network alone, despite the increased area. This is most likely due to the analysis design, where increasing the area enables several smaller patches to combine into one large one, losing the connections between the smaller patches. The analysis design would benefit from defining a specific size range for the patches.




Figure 28. Species-specific connectivity of landscape patches representing habitats for five species within the marine Natura 2000 sites and HELCOM MPA network. The chart shows the percentage of patches with 20 or more connections. The values were calculated as the number of connections between landscape patches chosen for a species within the combined area of marine Natura 2000 sites and HELCOM MPAs for five species, based on the shapefile of landscapes (Figure 6), shapefiles of MPAs provided by the HELCOM countries and shapefiles of the Natura 2000 sites downloaded from the EEA Data & Map service. The target is reached when 50% of the patches have at least 20 connections (red line).

SUMMARY OF THE RESULTS OF THE CONNECTIVITY ASSESSMENT OF THE COMBINED NETWORK OF HELCOM MPAS AND MARINE NATURA 2000 SITES IN THE BALTIC SEA

The combined area of the HELCOM MPAs and the marine Natura 2000 sites meets the targets set for species-specific connectivity (subcriteria 2) in this assessment in four of five cases, but in none for theoretical connectivity (subcriteria 1). The minimum target of 20 connections for at least 50% of the landscape patches representing habitats for a chosen species is met for all species in the analysis, except *Fucus vesiculosus* (subcriteria 2). This is explained by the very short dispersal distance of *F. vesiculosus* (only 1 km).

3.5 Conclusions of the ecological coherence assessment

HELCOM has agreed that in order to conclude that the MPA network is ecologically coherent, all main criteria need to meet their targets. As only the replication criterion meets its target fully, the Baltic Sea MPA network is not yet ecologically coherent (however, see chapter 4 for a test on a quantitative integrated approach of ecological coherence).

The targets set for the **representativity** criterion were partly met in this assessment. With 12% areal coverage, the representativity of HELCOM MPAs in the Baltic Sea exceeds the 10% target. However, improved protection is needed in the open sea and for most of the aphotic landscapes (commonly found in the open sea), as well as in the Bothnian Bay, Bothnian Sea, Åland Sea, Northern Baltic Proper, Eastern and Western Gotland Basins. The results described in chapter 3.1 revealed that 40% of the landscapes (mainly aphotic landscapes) are in need of improved protection. However, the open sea aphotic zone also includes areas which are suffering from severe and long-term oxygen depletion, and are therefore not meaningful to be designated as HELCOM MPAs.

The targets set for the **replication** criterion were met in this assessment for all but two biotope complexes. All other features used to analyse the two subcriteria had at least three replicates within the HELCOM MPA network. It should be noted that the chosen target is static and rather low, and does not work equally well for all species and biotope complexes. In an ideal situation, the target would be adjusted for each species, biotope complex and landscape, based on scientific background information.

The targets set for the **adequacy** criterion were not met in this assessment. The MPA network fell 12 percentage points short from the 80% target for the marine areas (68% reached).

The targets set for the **connectivity** criterion were partly met in this assessment. Theoretical connectivity (subcriteria 1) showed a very low number of connections for all landscapes, but regarding species-specific connectivity (subcriteria 2), the target was met for all chosen species, except *Fucus vesiculosus*. This correlates directly to the dispersal distances of the chosen species, where *F. vesiculosus* has the shortest dispersion distance of 1 km, in comparison to the other species (25-100 km).

The combined network of the HELCOM MPAs and the marine Natura 2000 sites is 8127 km² larger than the HELCOM MPA network alone, and analysing these networks together reveals improved representativity, replication and connectivity. This is a result of the larger size of the network, along with the typical coastal location of the Natura 2000 sites. For example, the typical coastal landscape *photic sand in 5-18 psu*, which covers 3.5% of all landscapes in the Baltic Sea, doubles its representativity value when the Natura 2000 network is combined to the HELCOM MPA network. The fourth criteria, adequacy, was not analysed for this combined network. While the marine Natura 2000 sites and the HELCOM MPA networks have a large overlap in the Baltic Sea, they target slightly different features in their protection; the Natura 2000 sites protect certain natural habitats and species in the EU countries, while the HELCOM MPAs aim to protect habitats, species and processes specific for the Baltic Sea. Together these protected areas cover a larger area and ensure better protection to the habitats and species in that area.

The results of the current ecological coherence assessment do not differ distinctively from the previous assessment (HELCOM 2010). The main difference can be seen in the replication of species, where the previous assessment concluded that some of the assessed species, namely *Fucus serratus*, *Zostera marina*, *Alosa fallax*, *Anguilla anguilla* and *Gadus morhua*, are in need of enhanced replication. Of these species, only *Anguilla*

anguilla was included in the current assessment, and this assessment indicates a better connectivity for this species than the previous HELCOM assessment.

3.5.1. Current data gaps and proposal for improvements to the assessment

The main shortcomings of the assessment are related to a lack of relevant data for the desired analyses, and partial lack of scientifically based targets and methodology.

Improving data availability

The main data gap for this assessment was the lack of adequate information on the distribution of species and biotope complexes, which lead to the exclusion of this sub-criterion from the assessment of representativity. For a more appropriate assessment of representativity, information on the spatial distribution is needed on these features, throughout the Baltic Sea. Currently only presence-data based on point wise observations is available. The previous assessment (HELCOM 2010) used the presence data as a proxy for distribution in the representativity assessment, but that is in fact a repetition of the replication assessment. In addition to improving the data availability, it is important to focus the data collection on the right type of data (for example, by continuously updating the data in the HELCOM MPA database).

Setting science-based targets for the subcriteria

Setting targets for the subcriteria analyses is a central question, as it determines the outcome of the ecological coherence assessment. The targets should be set with the prevailing environmental conditions in mind (for example, accounting for species abundance when setting the targets for adequate replication values for species) and revised to fit the scale of the used data. For the next ecological coherence assessment, all subcriteria targets should be reviewed, keeping in mind the scale of new and possibly more detailed maps. In particular, the representativity assessment of benthic marine landscapes could be improved by accounting for their relative abundance in the Baltic Sea, in order to distinguish the rare landscapes from the common ones. This would enable more precise targets to be set, for example, 60% coverage for rare habitats and 20% coverage for common habitats. The replication assessment would benefit from targets set according to each marker species and biotope complex, instead of the currently used fixed target.

Scrutinizing the methods of the analyses

The approach to evaluate connectivity in this assessment is rudimentary, and the results of species-specific connectivity clearly correspond to the different dispersal distances of the species. An appropriate analysis would, for example, be based on the direction and strength of major Baltic Sea currents and information on stepping stone habitats which facilitate species dispersal from one suitable habitat patch to another. Some of the required data and methods for a more elaborate analysis are however lacking at this time.

If data is available, replication could also be analysed within the MPAs, as currently it is only analysed between the MPAs. In addition, the reliability of the method for assessing adequacy would be improved by using more than one subcriterion.

4 Testing a new model for aggregating the results of subcriteria analyses of the ecological coherence assessment into a single outcome

Ecological coherence assessments previously carried out in HELCOM (HABITAT 2006, Piekäinen & Korpinen 2007, HELCOM 2010) have not attempted to aggregate numerical results of the subcriteria analyses into one single outcome of ecological coherence. Instead, most assessments have focused on describing how far the MPA network is from reaching ecological coherence. Here a method is tested for aggregating the results, using weighted averaging of the subcriteria and the one-out-all-out principle for the main criteria, while accounting for uncertainty of the data, target and method. A set of five integration tables were used to carry out the aggregation, according to the steps described below (Tables 9-13) (Wolters et al. 2015), based on the following criteria:

Subcriteria A ratio of the result of the subcriteria analysis and its target is calculated. Uncertainty in the data, target and method is defined, and averaged for each subcriteria. A weighted average is then calculated for each subcriteria (average uncertainty × subcriteria ratio). This is repeated for all subcriteria.

Main criteria An average of all subcriteria under one main criterion is calculated separately for all four main criteria. The likelihood of reaching the target is given for each criterion, based on the weighted average of the subcriteria.

Whole assessment Finally, the assessment of ecological coherence is based on the one-out-all-out principle, where the criterion with the lowest score (or likelihood) determines the final assessment result.

4.1 Integration tables of the results of subcriteria evaluations of the HELCOM MPAs

The outcomes of the subcriteria evaluations were compared against their specific targets and aggregated by using integration tables (adapted from Table 7 in Wolters et al. 2015).

Step-wise approach:

1. The subcriteria evaluated in chapters 3.1 - 3.4 aimed to assess how far the current state of the MPA network is from the target. To give this distance a value, the ratio of the subcriteria evaluation result and its target was attained (result divided by target) as a first step (Tables 9-12). A ceiling value of 2 was applied for this ratio in order to restrict disproportionally high values of subcriteria which highly exceed their targets.
2. In the second step, the uncertainties of the data, target and method of the analysis were estimated to be low, moderate or high, and translated into numerical values (1, 0.75 and 0.5, respectively). They were averaged to calculate a value for the mean uncertainty for each subcriteria. In the third step, the ratio and mean of uncertainties were used to calculate a weighted average for each subcriteria (subcriteria ratio × mean uncertainty). The rationale for the choice of uncertainty values are presented in Annex 6.

3. Finally, the score for the main criteria was attained by calculating a simple average of all subcriteria weighted averages. Based on these scores, the likelihood (very unlikely, unlikely, likely and very likely) that ecological coherence is reached was given for each main criterion (Table 8).

Table 8. Likelihood that target is currently achieved.

Likelihood target being achieved	Score of main criteria
VERY UNLIKELY	<0.5
UNLIKELY	0.5 - <1
LIKELY	1 - 1.5
VERY LIKELY	>1.5

Table 9. Integration table of the subcriteria of the representativity criterion in the ecological coherence assessment. For an explanation on the table calculations, please see chapter 4.1. The background colours indicate the different subcriteria of representativity (light blue = geographical representation in Baltic Sea sub-basins, lilac = geographical representation in Baltic Sea zones, light brown = representation of benthic marine landscapes within the MPAs).

Representativity of MPAs in	Subcriteria result	Subcriteria target	Subcriteria ratio (=result/target)	Uncertainty in data ⁶	Uncertainty in target ⁶	Uncertainty in method ⁶	Average uncertainty	Weighted average of subcriteria (=subcriteria ratio x mean uncertainty)
Kattegat	23.4	10.0	2.0	1.0	1.0	1.0	1.0	2.00
Great Belt	38.8	10.0	2.0	1.0	1.0	1.0	1.0	2.00
The Sound	14.3	10.0	1.4	1.0	1.0	1.0	1.0	1.43
Kiel Bay	38.9	10.0	2.0	1.0	1.0	1.0	1.0	2.00
Bay of Mecklenburg	17.4	10.0	1.7	1.0	1.0	1.0	1.0	1.74
Arkona Basin	14.7	10.0	1.5	1.0	1.0	1.0	1.0	1.47
Bornholm Basin	17.4	10.0	1.7	1.0	1.0	1.0	1.0	1.74
Gdansk Basin	16.3	10.0	1.6	1.0	1.0	1.0	1.0	1.63
Eastern Gotland Basin	7.0	10.0	0.7	1.0	1.0	1.0	1.0	0.70
Western Gotland Basin	3.6	10.0	0.4	1.0	1.0	1.0	1.0	0.36
Gulf of Riga	41.0	10.0	2.0	1.0	1.0	1.0	1.0	2.00
Northern Baltic Proper	3.2	10.0	0.3	1.0	1.0	1.0	1.0	0.32
Gulf of Finland	13.4	10.0	1.3	1.0	1.0	1.0	1.0	1.34
Åland Sea	5.6	10.0	0.6	1.0	1.0	1.0	1.0	0.56
Bothnian Sea	4.3	10.0	0.4	1.0	1.0	1.0	1.0	0.43
Quark	16.8	10.0	1.7	1.0	1.0	1.0	1.0	1.68
Bothnian Bay	3.8	10.0	0.4	1.0	1.0	1.0	1.0	0.38
Coastal sea	23.8	10.0	2.0	1.0	1.0	1.0	1.0	2.00
Outer coastal sea	11.0	10.0	1.1	1.0	1.0	1.0	1.0	1.10
Open sea	3.9	10.0	0.4	1.0	1.0	1.0	1.0	0.39
Representativity of land-scapes within MPAs								
Photic mud <5 psu	19.1	20.0	1.0	0.75	0.75	1.0	0.8	0.80
Photic mud 5-18 psu	34.8	20.0	1.7	0.75	0.75	1.0	0.8	1.45
Photic mud >18 psu	18.4	20.0	0.9	0.75	0.75	1.0	0.8	0.77
Aphotic mud <5 psu	9.1	20.0	0.5	0.75	0.75	1.0	0.8	0.38
Aphotic mud 5-18 psu	2.5	20.0	0.1	0.75	0.75	1.0	0.8	0.10
Aphotic mud >18 psu	11.2	20.0	0.6	0.75	0.75	1.0	0.8	0.47
Photic sand <5 psu	31.8	20.0	1.6	0.75	0.75	1.0	0.8	1.32
Photic sand 5-18 psu	52.7	20.0	2.0	0.75	0.75	1.0	0.8	1.67
Photic sand >18 psu	43.2	20.0	2.0	0.75	0.75	1.0	0.8	1.67
Aphotic sand <5 psu	4.4	20.0	0.2	0.75	0.75	1.0	0.8	0.18
Aphotic sand 5-18 psu	17.3	20.0	0.9	0.75	0.75	1.0	0.8	0.72
Aphotic sand >18 psu	20.9	20.0	1.0	0.75	0.75	1.0	0.8	0.87
Photic mixed sediment <5 psu	17.7	20.0	0.9	0.75	0.75	1.0	0.8	0.74
Photic mixed sediment 5-18 psu	22.1	20.0	1.1	0.75	0.75	1.0	0.8	0.92
Photic mixed sediment >18 psu	30.5	20.0	1.5	0.75	0.75	1.0	0.8	1.27
Aphotic mixed sediment <5 psu	6.2	20.0	0.3	0.75	0.75	1.0	0.8	0.26
Aphotic mixed sediment 5-18 psu	4.4	20.0	0.2	0.75	0.75	1.0	0.8	0.18
Aphotic mixed sediment >18 psu	24.0	20.0	1.2	0.75	0.75	1.0	0.8	1.00

⁵ See annex 6.

Representativity of land- scapes within MPAs	Subcriteria result	Subcriteria target	Subcriteria ratio (=result/ target)	Uncer- tainty in data ⁴	Uncer- tainty in target ⁶	Uncer- tainty in method ⁶	Average uncertainty	Weighted average of subcriteria (=subcriteria ratio x mean uncertainty)
Photic coarse sediment <5 psu	40.1	20.0	2.0	0.75	0.75	1.0	0.8	1.67
Photic coarse sediment 5-18 psu	55.9	20.0	2.0	0.75	0.75	1.0	0.8	1.67
Photic coarse sediment >18 psu	47.1	20.0	2.0	0.75	0.75	1.0	0.8	1.67
Aphotic coarse sediment <5 psu	21.0	20.0	1.0	0.75	0.75	1.0	0.8	0.87
Aphotic coarse sediment 5-18 psu	13.7	20.0	0.7	0.75	0.75	1.0	0.8	0.57
Aphotic coarse sediment >18 psu	35.1	20.0	1.8	0.75	0.75	1.0	0.8	1.46
Photic rock and other hard substrata <5 psu	14.0	20.0	0.7	0.75	0.75	1.0	0.8	0.58
Photic rock and other hard substrata 5-18 psu	19.9	20.0	1.0	0.75	0.75	1.0	0.8	0.83
Photic rock and other hard substrata >18 psu	36.8	20.0	1.8	0.75	0.75	1.0	0.8	1.53
Aphotic rock and other hard substrata <5 psu	20.1	20.0	1.0	0.75	0.75	1.0	0.8	0.84
Aphotic rock and other hard substrata 5-18 psu	11.1	20.0	0.6	0.75	0.75	1.0	0.8	0.46
Aphotic rock and other hard substrata >18 psu	36.0	20.0	1.8	0.75	0.75	1.0	0.8	1.50
Main criterion score of representativity								1.1
Likelihood								LIKELY

⁵ See annex 6.

Table 10. Integration table of the subcriteria of the replication criterion in the ecological coherence assessment. For an explanation on the table calculations, please see chapter 4.1. The background colours indicate the different subcriteria of replication (yellow = replication of marker species, green = replication of marker biotope complexes, light brown = replication of benthic marine landscapes).

Replication of species, biotope complexes and landscapes within MPAs	Subcriteria result	Subcriteria target	Subcriteria ratio (=result/target)	Uncertainty in data ⁵	Uncertainty in target ⁶	Uncertainty in method ⁶	Average uncertainty	Weighted average of subcriteria (=subcriteria ratio x mean uncertainty)
<i>Chara</i> spp.	21	4	2.0	1	0.5	0.75	0.8	1.50
<i>Anguilla anguilla</i>	22	4	2.0	1	0.5	0.75	0.8	1.50
<i>Lampetra fluviatilis</i>	34	4	2.0	1	0.5	0.75	0.8	1.50
<i>Salmo salar</i>	37	4	2.0	1	0.5	0.75	0.8	1.50
<i>Gavia arctica</i> & <i>Gavia stellata</i>	57	4	2.0	1	0.5	0.75	0.8	1.50
<i>Mergus serrator</i> & <i>Mergus merganser</i>	65	4	2.0	1	0.5	0.75	0.8	1.50
<i>Sternula albifrons</i>	32	4	2.0	1	0.5	0.75	0.8	1.50
<i>Tadorna tadorna</i>	39	4	2.0	1	0.5	0.75	0.8	1.50
<i>Somateria mollissima</i>	74	4	2.0	1	0.5	0.75	0.8	1.50
<i>Halichoerus grypus</i>	69	4	2.0	1	0.5	0.75	0.8	1.50
<i>Phoca vitulina</i> & <i>Phoca hispida</i> botnica	50	4	2.0	1	0.5	0.75	0.8	1.50
<i>Phocoena phocoena</i>	32	4	2.0	1	0.5	0.75	0.8	1.50
Biotope complex 1110	94	4	2.0	1	0.5	0.75	0.8	1.50
Biotope complex 1130	19	4	2.0	1	0.5	0.75	0.8	1.50
Biotope complex 1140	40	4	2.0	1	0.5	0.75	0.8	1.50
Biotope complex 1150	41	4	2.0	1	0.5	0.75	0.8	1.50
Biotope complex 1160	60	4	2.0	1	0.5	0.75	0.8	1.50
Biotope complex 1170	120	4	2.0	1	0.5	0.75	0.8	1.50
Biotope complex 1180	7	4	1.8	1	0.5	0.75	0.8	1.31
Biotope complex 1610	6	4	1.5	1	0.5	0.75	0.8	1.13
Biotope complex 1620	37	4	2.0	1	0.5	0.75	0.8	1.50
Biotope complex 1650	2	4	0.5	1	0.5	0.75	0.8	0.38
Biotope complex 8330	0	4	0.0	1	0.5	0.75	0.8	0.00
Photic mud <5 psu	66	4	2.0	0.75	0.5	0.5	0.6	1.17
Photic mud 5-18 psu	77	4	2.0	0.75	0.5	0.5	0.6	1.17
Photic mud >18 psu	111	4	2.0	0.75	0.5	0.5	0.6	1.17
Aphotic mud <5 psu	34	4	2.0	0.75	0.5	0.5	0.6	1.17
Aphotic mud 5-18 psu	80	4	2.0	0.75	0.5	0.5	0.6	1.17
Aphotic mud >18 psu	84	4	2.0	0.75	0.5	0.5	0.6	1.17
Photic sand <5 psu	106	4	2.0	0.75	0.5	0.5	0.6	1.17
Photic sand 5-18 psu	122	4	2.0	0.75	0.5	0.5	0.6	1.17
Photic sand >18 psu	166	4	2.0	0.75	0.5	0.5	0.6	1.17
Aphotic sand <5 psu	29	4	2.0	0.75	0.5	0.5	0.6	1.17
Aphotic sand 5-18 psu	77	4	2.0	0.75	0.5	0.5	0.6	1.17
Aphotic sand >18 psu	55	4	2.0	0.75	0.5	0.5	0.6	1.17

⁵ See annex 6.

Replication of species, biotope complexes and landscapes within MPAs	Subcriteria result	Subcriteria target	Subcriteria ratio (=result/target)	Uncertainty in data ⁵	Uncertainty in target ⁶	Uncertainty in method ⁶	Average uncertainty	Weighted average of subcriteria (=subcriteria ratio x mean uncertainty)
Photic mixed sediment <5 psu	115	4	2.0	0.75	0.5	0.5	0.6	1.17
Photic mixed sediment 5-18 psu	182	4	2.0	0.75	0.5	0.5	0.6	1.17
Photic mixed sediment >18 psu	170	4	2.0	0.75	0.5	0.5	0.6	1.17
Aphotic mixed sediment <5 psu	42	4	2.0	0.75	0.5	0.5	0.6	1.17
Aphotic mixed sediment 5-18 psu	134	4	2.0	0.75	0.5	0.5	0.6	1.17
Aphotic mixed sediment >18 psu	76	4	2.0	0.75	0.5	0.5	0.6	1.17
Photic coarse sediment <5 psu	26	4	2.0	0.75	0.5	0.5	0.6	1.17
Photic coarse sediment 5-18 psu	56	4	2.0	0.75	0.5	0.5	0.6	1.17
Photic coarse sediment >18 psu	8	4	2.0	0.75	0.5	0.5	0.6	1.17
Aphotic coarse sediment <5 psu	17	4	2.0	0.75	0.5	0.5	0.6	1.17
Aphotic coarse sediment 5-18 psu	40	4	2.0	0.75	0.5	0.5	0.6	1.17
Aphotic coarse sediment >18 psu	4	4	1.0	0.75	0.5	0.5	0.6	0.58
Photic rock and other hard substrata <5 psu	20	4	2.0	0.75	0.5	0.5	0.6	1.17
Photic rock and other hard substrata 5-18 psu	53	4	2.0	0.75	0.5	0.5	0.6	1.17
Photic rock and other hard substrata >18 psu	43	4	2.0	0.75	0.5	0.5	0.6	1.17
Aphotic rock and other hard substrata <5 psu	6	4	1.5	0.75	0.5	0.5	0.6	0.88
Aphotic rock and other hard substrata 5-18 psu	49	4	2.0	0.75	0.5	0.5	0.6	1.17
Aphotic rock and other hard substrata >18 psu	20	4	2.0	0.75	0.5	0.5	0.6	1.17
Main criterion score of replication								1.2
Likelihood								LIKELY

⁵ See annex 6.

Table 11. Integration table of the subcriterion of the adequacy criterion in the ecological coherence assessment. For an explanation on the table calculations, please see chapter 4.1. The background colour indicates the subcriteria MPA size (peach).

Adequacy of MPA size	Subcriteria result	Subcriteria target	Subcriteria ratio (=result/target)	Uncertainty in data ⁶	Uncertainty in target ⁶	Uncertainty in method ⁶	Average uncertainty	Weighted average of subcriteria (=subcriteria ratio x mean uncertainty)
MPAs with marine size of minimum 30km ²	68	80	0.9	1	0.75	1	0.92	0.78
MPAs with terrestrial size of minimum 10km ²	40	80	0.5	1	0.75	0.5	0.75	0.38
Main criterion score of adequacy								0.6
Likelihood								UNLIKELY

Table 12. Integration table of the subcriteria of the connectivity criterion in the ecological coherence assessment. For an explanation on the table calculations, please see the beginning of chapter 4.1. The background colours indicate the different subcriteria of connectivity (light brown = theoretical connectivity based on benthic marine landscapes, brown = species-specific connectivity, based on benthic marine landscapes representing suitable habitats for the chosen species).

Connectivity within/for	Subcriteria result	Subcriteria target	Subcriteria ratio (=result/target)	Uncertainty in data ⁶	Uncertainty in target ⁶	Uncertainty in method ⁶	Average uncertainty	Weighted average of subcriteria (=subcriteria ratio x mean uncertainty)
25 km for photic mud <5 psu	0	50	0.0	0.75	0.5	0.5	0.58	0.00
25 km for photic sand 5-18 psu	0.5	50	0.0	0.75	0.5	0.5	0.58	0.01
25 km for aphotic rock and other hard substrata <5 psu	0	50	0.0	0.75	0.5	0.5	0.58	0.00
25 km for aphotic sand 5-18 psu	0	50	0.0	0.75	0.5	0.5	0.58	0.00
25 km for aphotic mixed >18 psu	0	50	0.0	0.75	0.5	0.5	0.58	0.00
50 km for photic mud <5 psu	0.0	50	0.0	0.75	0.5	0.5	0.58	0.00
50 km for photic sand 5-18 psu	4.5	50	0.1	0.75	0.5	0.5	0.58	0.05
50 km for aphotic rock and other hard substrata <5 psu	0.0	50	0.0	0.75	0.5	0.5	0.58	0.00
50 km for aphotic sand 5-18 psu	0.0	50	0.0	0.75	0.5	0.5	0.58	0.00
50 km for aphotic mixed >18 psu	7.5	50	0.1	0.75	0.5	0.5	0.58	0.09
<i>Macoma baltica</i>	95.9	50	1.9	0.75	0.75	0.5	0.67	1.28
<i>Psetta maxima</i>	64.6	50	1.3	0.75	0.75	0.5	0.67	0.86
<i>Furcellaria lumbricalis</i>	62.1	50	1.2	0.75	0.75	0.5	0.67	0.83
<i>Idotea baltica</i>	62.1	50	1.2	0.75	0.75	0.5	0.67	0.83
<i>Fucus vesiculosus</i>	7.2	50	0.1	0.75	0.75	0.5	0.67	0.10
Main criterion score of connectivity								0.3
Likelihood								VERY UNLIKELY

⁵ See annex 6.

4.2 Final outcome of the ecological coherence assessment for the HELCOM MPA network, using the integration tables for aggregating the subcriteria results

The final step of assessing the likelihood that the HELCOM MPA network has reached ecological coherence is done by the one-out-all-out principle between the four main criteria, i.e. the criterion with the lowest likelihood determines the final assessment outcome. Based on the aggregated results of this assessment, it is very unlikely that the HELCOM MPA network is ecologically coherent (Table 13). While representativity and replication were evaluated to be likely to have reached ecological coherence, adequacy and connectivity were unlikely respective very unlikely to be sufficient to fulfil the conservation goals of the HELCOM MPA network (Table 13).

Using integration tables for calculating the main outcome of the ecological coherence assessment is a straightforward and transparent method. In addition, it accounts for uncertainties in the data, target and method, and can therefore be used despite differing accuracy of the available background data. The integration tables are also easy to reproduce in order to create a long term view of the development of the ecological coherence of the MPA network. The current integration tables are a first attempt at applying this method for aggregating the final outcome of the ecological coherence assessment of HELCOM MPAs, and this method is proposed to be applied in future ecological coherence assessments. However, some details of the subcriteria evaluations will inevitably be lost in the aggregation, and the integration tables will also in the future be accompanied by a descriptive outcome of the subcriteria evaluations, as in chapter 3 in this report. Future assessments should focus on developing the targets for the subcriteria analyses, as these have the largest influence when interpreting the results, both for the descriptive and the aggregated outcome.

Table 13. Scores of the main criteria and final aggregated outcome of the ecological coherence assessment.

Main criterion	Score	Likelihood	Ecological coherence of the HELCOM MPA network
REPRESENTATIVITY	1.1	LIKELY	It is very unlikely that ecological coherence is reached.
REPLICATION	1.2	LIKELY	
ADEQUACY	0.6	UNLIKELY	
CONNECTIVITY	0.3	VERY UNLIKELY	

5 Recommendation 35/1 on coastal and marine Baltic Sea protected areas (HELCOM MPAs⁶)

HELCOM Recommendation 35/1 ‘System of coastal and marine Baltic Sea protected areas (HELCOM MPAs⁶)’ was adopted on 1 April 2014, superseding HELCOM Recommendation 15/5. It recommends that the Governments of the Contracting Parties to the Helsinki Convention take all appropriate measures to step up efforts to establish an ecologically coherent and effectively managed network of coastal and marine Baltic Sea protected areas (HELCOM MPAs) and to improve the protection effectiveness of existing HELCOM MPAs.

The Recommendation also lists more detailed objectives which the Contracting Parties are recommended to achieve or carry out, labelled as paragraphs a-r. The Recommendation is followed up through the paragraphs that have specific targets, or time limits, for achievement by use of specific evaluation criteria. This chapter presents the current level of accomplishment of the commitments expressed in these paragraphs. In addition, the progress of objectives in a set of paragraphs without defined targets is also included in this chapter. The evaluation criteria, as well as additional guiding paragraphs from the Recommendation that are not included in the assessment of accomplishment, are included in Annex 7.

5.1 Current status of implementation

5.1.1 Accomplishment of Recommendation 35/1

Reach the target set by the HELCOM 2010 Moscow Ministerial Declaration that at least 10% of the marine area in all sub-basins of the Baltic Sea, including the EEZ areas beyond territorial waters, is covered by MPAs, where scientifically justified (paragraph a).

This objective overlaps with a subcriterion used to evaluate representativity as part of the assessment of ecological coherence of the MPA network. As shown in chapter 3.1.1, the 10% target is met in some sub-basins, as well as on the Baltic Sea level as a whole (Table 14 and Figure 7). However, since the target is not met in all sub-basins, the objective expressed in this paragraph is assessed as being “partly accomplished” (Table 16). This conclusion is furthermore supported, if the coverage of MPAs in the EEZ is considered: currently, MPAs are covering 5% of the EEZ (Table 14).

It should however be noted that paragraph a) includes the clause that the target of 10% areal coverage should be reached “where scientifically justified”. Thus, further consideration is needed to evaluate whether the 10% target is scientifically justified in the Eastern and Western Gotland Basin, as well as the Northern Baltic Proper, Åland Sea, Bothnian Sea and Bothnian Bay.

⁶ Former BSPAs.

Table 14. Percentage of HELCOM MPAs in HELCOM sub-basins – the basis for assessing the level of accomplishment of paragraph a) of Recommendation 35/1. The green cells show the sub-basins which have already reached the target of 10% areal protection.

Sub-basin	HELCOM MPA network coverage per sub-basin (%)	HELCOM MPA network coverage in the EEZ (% of tot EEZ per sub-basin)
Kattegat	23	20
Great Belt	39	5
The Sound	14	0.3
Kiel Bay	39	32
Bay of Mecklenburg	17	14
Arkona Basin	15	17
Bornholm Basin	17	13
Gdansk Basin	16	0
Eastern Gotland Basin	7	2
Western Gotland Basin	4	5
Gulf of Riga	41	100
Northern Baltic Proper	3	0
Gulf of Finland	13	2
Åland Sea	6	0
Bothnian Sea	4	0.7
The Quark	17	0
Bothnian Bay	4	0.1
Total area	12	4.7

Designate new sites as HELCOM MPAs where ecologically meaningful, especially in offshore areas beyond territorial waters (paragraph b).

Since the adoption of Recommendation 35/1 in 2014, only Finland has designated new sites. The 11 new Finnish sites increase the total area of HELCOM MPAs in the Baltic Sea by 725 km². Moreover, these sites are located in, or partly in, the EEZ, increasing the EEZ area covered by HELCOM MPAs by 82 km². However, the offshore area, and the landscapes typically encountered within it, are still in need of improved protection, as the assessment of the representativity criterion shows (Figures 8 and 9). Therefore, this objective should be seen as an ongoing process.

Ensure that HELCOM MPAs inter alia provide specific protection to those species, habitats, biotopes and biotope complexes included in the HELCOM Red Lists, as agreed upon in the HELCOM 2013 Copenhagen Ministerial Declaration, by considering these in the site selection procedure (paragraph c).

This objective is assessed as accomplished when the network of HELCOM MPAs provides protection to all species, biotopes and biotope complexes defined as threatened in the HELCOM Red Lists published in 2013 (HELCOM 2013b, c). The assessment of the accomplishment of this objective is based on information reported by the Contacting Parties to the HELCOM MPA database regarding the presence of threatened species, biotopes and biotope complexes in at least one MPA, and more specifically, information on whether the threatened species, biotope or biotope complex is protected in at least one MPA (i.e. justifies the site's establishment as an MPA). According to data reported in the HELCOM MPA database, this objective is currently assessed as being partly accomplished (Figure 29). It is accomplished for the threatened biotope complexes, as each one of them is reported to be protected in at least one HELCOM MPA. This objective has however not been reached for the threatened species and biotopes, as only 36%

of all threatened species, and 12% of threatened biotopes are protected within at least one MPA. The biotopes follow the HELCOM HUB classification and are in general very poorly reported in the HELCOM MPA database, which partly explains this low result.

According to the information, some threatened species are protected in many MPAs, for example the common eider, black-throated diver and velvet scoter, as well as the harbour porpoise. Threatened species which are protected in only one or a few MPAs include birds, macrophytes, fish and lamprey species and invertebrates (Table 15). In general, threatened species are the reason for establishing many HELCOM MPAs.

Figure 29. The percentage (%) of threatened species, biotopes and biotope complexes present (blue), or present and protected (green), in at least one MPA, as reported in the HELCOM MPA database. Present and protected means that the HELCOM MPA is established to protect this feature in particular (i.e. the feature justifies the site's establishment as an MPA). The threatened categories include regionally extinct (RE), critically endangered (CR), endangered (EN) and vulnerable (VU). The HELCOM Red Lists (HELCOM 2013b, c) include a total of 69 threatened species, 17 threatened biotopes and 8 threatened biotope complexes.




Table 15. Threatened species according to HELCOM Red List 2013 (HELCOM 2013 c) protected in HELCOM MPAs.

Species group	Scientific name	English name	Protected in no. MPAs	Countries
Fish and lamprey	<i>Anguilla anguilla</i>	European eel	1	Sweden
Birds	<i>Cepphus grylle arcticus</i> (wintering)	Black guillemot	1	Germany
Macrophytes	<i>Hippuris tetraphylla</i>		1	Sweden
Birds	<i>Rissa tridactyla</i> (breeding)	Black-legged kittiwake	1	Sweden
Fish and lamprey	<i>Salmo salar</i>	Salmon	1	Russia
Invertebrates	<i>Scrobicularia plana</i>		1	Denmark
Macrophytes	<i>Alisma wahlenbergii</i>		2	Sweden, Russia
Invertebrates	<i>Macoma calcarea</i>		2	Denmark
Birds	<i>Larus fuscus fuscus</i> (breeding)		3	Sweden, Germany
Macrophytes	<i>Zostera noltii</i>		3	Denmark
Birds	<i>Arenaria interpres</i> (breeding)	Ruddy turnstone	4	Sweden
Birds	<i>Hydroprogne caspia</i> (breeding)	Caspian tern	4	Sweden, Denmark, Poland
Fish and lamprey	<i>Salmo trutta</i>	Trout	4	Russia, Sweden
Birds	<i>Podiceps grisegena</i> (wintering)	Red-necked grebe	5	Poland, Russia, Germany
Birds	<i>Polysticta stelleri</i> (wintering)	Steller's eider	5	Finland
Fish and lamprey	<i>Petromyzon marinus</i>	Sea lamprey	6	Poland, Denmark, Russia
Mammals	<i>Phoca hispida botnica</i>	Baltic ringed seal	7	Sweden, Finland, Russia
Birds	<i>Anser fabalis fabalis</i> (wintering)	Taiga Bean goose	10	Poland, Russia, Denmark
Birds	<i>Clangula hyemalis</i> (wintering)	Long-tailed duck	11	Finland, Latvia, Denmark, Poland, Germany, Sweden
Birds	<i>Aythya marila</i> (breeding)	Greater scaup	12	Poland, Denmark, Russia, Germany
Birds	<i>Calidris alpina schinzii</i> (breeding)	Southern dunlin	14	Finland, Denmark, Sweden, Poland
Birds	<i>Melanitta nigra</i> (wintering)	Common scoter	14	Latvia, Denmark, Poland, Sweden, Germany
Birds	<i>Mergus serrator</i> (wintering)	Red-breasted merganser	14	Germany, Denmark, Latvia, Sweden, Poland
Birds	<i>Philomachus pugnax</i> (breeding)	Ruff	14	Sweden, Finland
Birds	<i>Melanitta fusca</i> (breeding)	Velvet scoter	18	Finland, Latvia, Denmark, Sweden, Germany, Poland
Birds	<i>Gavia stellata</i> (wintering)	Red-throated diver	20	Finland, Russia, Poland, Latvia, Sweden
Mammals	<i>Phocoena phocoena</i>	Harbour porpoise Western Baltic subpopulation	20	Denmark, Sweden, Germany
Birds	<i>Gavia arctica</i> (wintering)	Black-throated diver	22	Sweden, Finland, Latvia, Germany, Poland, Russia
Birds	<i>Somateria mollissima</i> (breeding)	Common eider	25	Finland, Germany, Denmark, Sweden

Ensure when selecting new areas that the network of HELCOM MPAs is ecologically coherent and takes into account connectivity between sites including, for example, migration routes, species mobility and areas of special ecological significance, such as spawning areas (paragraph d).

According to the assessment presented in chapter 3 of this report, the HELCOM MPA network in the Baltic Sea is not yet ecologically coherent. However, two of the four main criteria (representativity and replication; see Table 13) are likely to have reached ecological coherence, and the objective is thereby assessed as being partly accomplished.

Develop and apply management plans or measures for all existing HELCOM MPAs by 2015 (paragraph h1).

Paragraph h1) of the Recommendation is related to the development of management plans for the MPAs. The objective is that all existing MPAs should have implemented management plans or measures by 2015. Information provided to the HELCOM MPA database shows that this objective has not been met. On a regional level, the percentage of MPAs with management plans is the same as in 2014, i.e. 67%, and thus, the target has only been partly accomplished.

Assess the effectiveness of the management plans or measures of HELCOM MPAs by conducting monitoring, and where feasible, scientific research programmes, which are directly connected to the conservation interests of HELCOM MPAs, including the placement of monitoring stations inside the MPAs (paragraph k).

Currently, HELCOM has no joint approach on how to assess the effectiveness of management plans or measures and the objective is thus not accomplished. However, monitoring, which is required to assess the effectiveness of MPAs, takes place in 64% of MPAs, as reported in the MPA database.

Modernize the HELCOM MPA database, as agreed upon in the HELCOM 2013 Copenhagen Ministerial Declaration, taking into account and harmonizing with other similar databases (paragraph p).

This objective is accomplished, as the new HELCOM MPA database (mpas.helcom.fi) was published in October 2015. It contains a map interface, new reporting features on pressures and regulated human activities within the MPAs, and is harmonized with the OSPAR MPA database and Natura 2000 standard data forms, where feasible. Data, information and shapefiles reported to the database form the basis for the assessment of ecological coherence and accomplishment of Recommendation 35/1.

Update the management plans when necessary and in accordance with other legal requirements with a maximum of 12 year intervals (paragraph i).

The total number of management plans reported in the HELCOM MPA database by January 2016 is 216, and 140 of them are implemented. Seven of the implemented management plans have not been updated in the last 12 years (i.e. after 2004). Therefore, this target has not been accomplished. Management plans still need to be developed or finalized for several HELCOM MPAs, and those seven management plans that are older than 12 years need to be updated.

Establish management plan or measures for every new MPA within five years after its designation (paragraph h2).

Recommendation 35/1 was adopted in 2014 and, thus this target will be assessed in 2019.

5.1.2 Summary on Recommendation 35/1 follow-up

In conclusion, out of the eight paragraphs with agreed targets and time limits of the Recommendation 35/1, one has been accomplished, five have partly been accomplished, and two objectives have not been accomplished (Table 16). In addition, the second half of paragraph h of Recommendation 35/1 cannot yet be assessed, since its target year is set in the future.

Table 16. Evaluation of the follow-up of HELCOM Recommendation 35/1. The coloured cells describe the level of accomplishment; green = accomplished, yellow = partly accomplished, and red = not accomplished.

Paragraphs from Recommendation 35/1	Evaluation	Evaluation criteria supporting assessment	Additional info to justify accomplishment
a) reach the target set by the HELCOM 2010 Moscow Ministerial Declaration that at least 10% of the marine area in all sub-basins of the Baltic Sea, including the EEZ areas beyond territorial waters, is covered by MPAs, where scientifically justified;	Partly accomplished	10% of the marine area is covered by MPAs in some sub-basins, including the EEZ areas beyond territorial waters (presented per sub-basin).	In 11 of 17 sub-basins, MPAs cover at least 10% of the surface. The EEZ is sufficiently covered in some sub-basins (Table 14).
b) designate new sites as HELCOM MPAs, where ecologically meaningful, especially in offshore area beyond territorial waters;	Partly accomplished	The number and/or percentage of MPAs is increasing, and number and/or percentage of MPAs in the EEZ is increasing.	Only Finland has designated new sites. The 11 new Finnish sites increase the total area of HELCOM MPAs in the Baltic by 725km ² . Finland has designated three new sites in or partly in the EEZ, which increases the EEZ area covered by HELCOM MPAs by 82km ² .
c) ensure that HELCOM MPAs inter alia provide specific protection to those species, habitats, biotopes and biotope complexes included in the HELCOM Red Lists, as agreed upon in the HELCOM 2013 Copenhagen Ministerial Declaration, by considering these in the site selection procedure;	Partly accomplished	1)The HELCOM network of MPAs provides protection to some HELCOM threatened species 2) The HELCOM network of MPAs provides protection to some HELCOM threatened biotopes/habitats (both to be met)	All threatened biotope complexes are protected in at least one MPA, whereas only 36% of all threatened species, and 12% of threatened biotopes are protected in an MPA.
d) ensure when selecting new areas, that the network of HELCOM MPAs is ecologically coherent and takes into account connectivity between sites, including for example, migration routes, species mobility and areas of special ecological significance, such as spawning areas;	Partly accomplished	Some criteria considered in the evaluation are reflecting ecological coherence.	According to the assessment presented in chapter 3, the MPA network in the Baltic Sea is not ecologically coherent.
h1) develop and apply management plans or measures for all existing HELCOM MPAs by 2015,	Partly accomplished	At least 67% of HELCOM MPAs designated by 2014 have management plans or measures.	
h2) and establish management plan or measures for every new MPA within five years after its designation;	Not assessed		Not assessed - will be assessed in 2019 (5 years after the adoption of Rec35-1).
i) update the management plans when necessary and in accordance with other legal requirements with a maximum of 12 years intervals;	Not accomplished		Management plans still need to be developed or finalized for several HELCOM MPAs. Seven management plans are older than 12 years and need to be updated.
k) assess the effectiveness of the management plans or measures of HELCOM MPAs by conducting monitoring, and where feasible scientific research programmes, which are directly connected to the conservation interests of HELCOM MPAs, including the placement of monitoring stations inside the MPAs;	Not accomplished	Assessment management effectiveness not started.	Not accomplished - currently there is no joint approach on how to assess the effectiveness of management plans/measures. However, 64% of MPAs with management plans also have monitoring in place as reported to the MPA database.
p) modernize the HELCOM MPA database, as agreed upon in the HELCOM 2013 Copenhagen Ministerial Declaration, taking into account and harmonizing with other similar databases;	Accomplished	New database published online in October 2015	http://mpas.helcom.fi/apex/

5.1.3 Progress of additional commitments

In addition to the paragraphs with defined targets and time limits, Recommendation 35/1 includes a set of paragraphs, which only address some Contracting Parties, or which are expressed as encouragement, rather than commitments.

Encourage Contracting Parties which are also EU Member States to designate, when feasible, all appropriate Natura 2000 sites as HELCOM MPAs, and to consider all Natura 2000 sites, as well as other marine protected areas, when evaluating the network of marine protected areas (paragraph f).

Today 63% of the marine Natura 2000 area in the Baltic Sea is also designated into the HELCOM MPA network, and this has not changed since 2013.

In addition, there are six HELCOM MPAs within the EU borders which have not been designated as Natura 2000 sites (one in Finland, Denmark, Lithuania and Sweden, and two in Germany).

The second objective of the paragraph, “to consider all Natura 2000 sites, as well as other marine protected areas when evaluating the network of marine protected areas”, has been achieved with the current assessment of ecological coherence (chapter 3), which also considers the Natura 2000 sites.

Encourage Contracting Parties, which are also OSPAR Contracting Parties, to designate, when appropriate, OSPAR MPAs as HELCOM MPAs, in order to harmonize the networks where the conventions’ geographical scope overlap (paragraph g).

The geographical scope of the OSPAR and HELCOM regional sea conventions overlap in Kattegat, and there are 22 OSPAR MPAs in the Kattegat. All OSPAR MPAs in Kattegat are also HELCOM MPAs and have been so since 2005 or 2009. In addition, there are seven HELCOM MPAs in the Kattegat which are not designated as OSPAR MPAs.

Apply the newest IUCN categorization system when describing the HELCOM MPAs, in order to allow for global comparisons of regional networks (paragraph n).

According to data reported in the HELCOM MPA database, an IUCN category (Table 6) is provided for 118 of all 174 (68%) HELCOM MPAs. The most common IUCN category for the HELCOM MPAs is category IV Habitat/Species Management Area (Figure 23 and Table 6 in chapter 3). In many cases, one MPA has been assigned several IUCN categories, as the IUCN categories can be assigned to smaller areas inside the MPA.

6 Next steps for improving the network and assessments of HELCOM MPAs

From the perspective of a marine region, this report shows an impressive system of marine protected areas in the Baltic Sea region, by far exceeding the areal coverage of MPAs at the European and global level⁷. However, the ambition set by HELCOM for the MPA network is high and the commitments made by the Contracting Parties have been only partly met to date. In addition, the lack of information and consolidated methodology hampers the assessment of the ecological coherence of the network, as well as the effectiveness of its management. Some proposed steps of improvement require implementation by countries, while others are best achieved through joint efforts.

6.1 Measures by countries

Designate new sites as HELCOM MPAs, where scientifically meaningful, especially in offshore areas beyond territorial waters

The increase in areal coverage of HELCOM MPAs has slowed down in recent years. Although the target of 10% areal coverage has been met for the Baltic Sea as a whole, the current report shows that the target has not been met in all sub-basins, and also not in waters beyond territorial waters (chapter 3.1 and 5.1). While the designation of new MPAs is a national matter, such decisions can be supported by joint analyses to agree upon the need to improve protection in sub-basins and zones which do not yet reach the 10% target. As recognized in Recommendation 35/1, the relevance of spatial protection differs between areas, depending on the characteristics of the respective sub-basin. For example the offshore areas of the Baltic Sea are subject to oxygen depletion, and the designation of MPAs may not be meaningful in these areas from the point of protecting important or rare features.

Improve the protection of threatened species, biotopes and biotope complexes within the MPAs

Surprisingly few threatened species and biotopes have been reported as being protected in the MPAs, although the role of MPAs in protecting such species and biotopes is highlighted in Recommendation 35/1 (chapter 5.1). This evaluation is based on reporting by countries to the HELCOM MPA database. Thus, if countries have not reported consistently in this regard, the protection of threatened species may be underestimated. The results indicate the need to scrutinize the purpose of the MPAs with regard to protection of threatened species, but also the reporting to the HELCOM MPA database.

Develop management plans for MPAs still lacking them

Development of management plans for existing MPAs was laid down in the Baltic Sea Action Plan with an initial deadline of 2010, being extended for existing MPAs to 2015 in HELCOM Recommendation 35/1. Currently this is met for 67% of the MPAs (chapter 5.1). In addition, the agreement to update the management plans with a maximum of 12 year intervals has been exceeded for seven MPAs.

⁷ The total area of MPAs in the North-east Atlantic Ocean, Mediterranean Sea and Black Sea was estimated at 4.2, 9.5 and 5.9% respective in 2015 (EEA 2015, Technical Report No 17/2015), and 2.1% of the global oceans in 2016 (www.mpatlas.org).

Increase monitoring and information on monitoring in MPAs

Currently, monitoring is taking place in 64% of HELCOM MPAs as reported to the HELCOM MPA database. Monitoring in MPAs is a prerequisite to evaluate the effectiveness of MPAs and their management (see also below). The frequency of monitoring has however not been reported to the database and it is therefore difficult to estimate whether the monitoring is sufficient for analysing temporal changes in those MPAs where monitoring exists.

Ensure complete reporting to the HELCOM MPA database

This assessment is, to a large extent, based on the reporting by countries to the HELCOM MPA database. If the reporting is not complete, the conclusions drawn from the information are incorrect. This refers to reporting on protected species, biotopes and biotope complexes (see above), as well as other features such as pressures within the MPA, management plan status and regulated activities within the MPA.

Encourage Contracting Parties, which are also EU Member States to designate, when feasible, all appropriate Natura 2000 sites as HELCOM MPAs

There is still 37% of marine Natura 2000 sites which are not designated to the HELCOM MPA network. In order to improve the HELCOM MPA network, the Contracting Parties are encouraged to designate these areas as HELCOM MPAs, when feasible.

6.2 Joint measures

Maximize the possibility of enhancing the ecological coherence when designating new MPAs

One of the key outcomes of the assessment is the lack of ecological coherence of the MPA network, and, in particular, the aspect of connectivity that reflects the possibilities of species to migrate and disperse within and between MPAs. To achieve better connectivity, countries need to cooperate in the designation of the MPAs, to ensure that the network as a whole will meet the desired features, such as connectivity. The use of site selection tools to identify areas suitable for new MPAs have been used previously in HELCOM (HELCOM 2010), and are proposed to be explored again.

Further develop the assessment methodology for the next ecological coherence assessment, including the targets for the subcriteria

As detailed in chapter 3.5.1, several improvements can be made for the next ecological coherence assessment. These include improving data availability, setting science-based targets for all subcriteria and scrutinizing the methods for some of the analyses. The integration table provides a calculative approach for aggregating the results of the subcriteria analyses, and is encouraged to be used along with the descriptive approach.

Include all Baltic Sea MPAs in the next assessment

In the next assessment, all MPAs established in the Baltic Sea, such as national protected areas, Ramsar sites and Important Bird Areas should preferentially be included in addition to HELCOM MPAs and the marine Natura 2000 sites. The prerequisite for such an assessment is that the appropriate data for the analysis is available for all protected area networks in the Baltic Sea.

Develop an approach and carry out an assessment of management effectiveness for HELCOM MPAs

While management plans are a required component in the process of designating MPAs, and are being regularly followed-up in HELCOM, an assessment of the effectiveness of such plans or other measures applied in the MPAs has not yet been carried out. A first step is to develop a common HELCOM approach to assess the management effectiveness.

Update HELCOM guidelines

HELCOM Recommendation 35-1 recommends Contracting Parties to 'update, when necessary, HELCOM MPA related guidelines and guiding documents, in order to keep them in line with new knowledge and compatible with other international criteria, such as MSFD requirements, in particular those concerning spatial protection measures'. The latest HELCOM MPA guidelines are from 2006 (BSEP 105, Planning and management of Baltic Sea Protected Areas: guidelines and tools). It is thus timely to review and tentatively revise the existing guidelines.

Improve spatial data on the distribution of species and biotopes

A lack of data has hampered the assessment of ecological coherence. Better data on the distribution of species and biotope complexes would have improved the analysis of, for example, representativity.

References

- Al-Hamdani Z., Reker J.(2007). Towards marine landscapes in the Baltic Sea. BALANCE Interim Report No. 10. 116 pp.
- Andersson Å., Korponen S., Liman A.S., Nilsson P.R., Piekäinen H.(2007). Ecological coherence and principles for MPA assessment, selection and design. BALANCE Technical Summary Report No. 3/4.
- Anonymous 1979, (2009). Council Directive 09/147/EEC on the conservation of wild birds (the codified version of Directive 79/409/EEC as amended).
- Anonymous(1992). Council Directive 92/43/EEC on the Conservation of natural habitats and of wild fauna and flora.
- BirdLife International (2015). Detailed regional assessment and species account from the European Red List.
- Catchpole R. (2012). Ecological Coherence Definitions in Policy and Practice. Contract Report to Scottish Natural Heritage, No. 41102, 28pp.
- CBD(2008). Ninth meeting of the Conference of the Parties to the Convention on Biological Diversity, Bonn, Germany, 19-30 May 2008. Available at: <http://www.cbd.int/decision/cop/?id=11663>.
- CBD (2010). Strategic Plan for Biodiversity 2011-2020, including Aichi Biodiversity targets. Available at: <https://www.cbd.int/sp/>.
- EC (2007). Guidelines for the establishment of the Natura 2000 network in the marine environment. Application of the Habitats and Birds Directives.
- EEA (2015). Technical Report No 17/2015.
- EUSEaMap (2015). Interim draft Baltic Sea Broad-Scale Predictive Habitat Map.
- HELCOM (1994). HELCOM Recommendation 15/5 on the system of coastal and marine Baltic Sea Protected Areas. Available online: <http://helcom.fi/Recommendations/Rec%2015-5.pdf>.
- HELCOM (2003). Guidelines for Designating Marine and Coastal Baltic Sea Protected Areas (BSPA) and Proposed Protection Categories. Updated and approved by HELCOM HOD 11/2003.
- HELCOM (2006a). Initial analysis on status of ecological coherence of the BSAP network. HELCOM HABITAT 8/2006, document 5.2/5.
- HELCOM (2007a). HELCOM Baltic Sea Action Plan. 102 pp.
- HELCOM (2007b). Assessment of the ecological coherence of the network of Baltic Sea Protected Areas, HELCOM 28/2007, document 3/13.
- HELCOM (2009). Biodiversity in the Baltic Sea – an integrated thematic assessment on biodiversity and nature conservation in the Baltic Sea. Baltic Sea Environmental Proceedings No. 116B.
- HELCOM (2010). Towards an ecologically coherent network of well-managed Marine Protected Areas – Implementation report on the status and ecological coherence of the HELCOM BSPA network. Baltic Sea Environmental Proceedings No. 124B.
- HELCOM (2012). Checklist of Baltic Sea Macro-species. Baltic Sea Environment Proceedings No. 130
- HELCOM (2013a). HELCOM PROTECT-Overview of the status of the network of Baltic Sea marine protected areas. 31pp.
- HELCOM (2013b). Red List of Baltic Sea underwater biotopes, habitats and biotope complexes. Baltic Sea Environmental Proceedings No. 138.
- HELCOM (2013c). HELCOM Red List of Baltic Sea species in danger of becoming extinct. Baltic Sea Environmental Proceedings No. 140.
- HELCOM (2013d). Monitoring and Assessment Strategy (2013).

- HELCOM (2013e). HELCOM Red List of Baltic Sea species in danger of becoming extinct. Species Information Sheets. Available at: [http://helcom.fi/baltic-sea-trends/biodiversity/red-list-of-species/\(under species groups\)](http://helcom.fi/baltic-sea-trends/biodiversity/red-list-of-species/(under%20species%20groups)).
- HELCOM (2013f). HELCOM Red List of Baltic Sea underwater biotopes, habitats and biotope complexes. Biotope Information Sheets: Available at: <http://helcom.fi/baltic-sea-trends/biodiversity/red-list-of-biotopes-habitats-and-biotope-complexes/biotope-information-sheets/>.
- HELCOM (2014). HELCOM Recommendation 35/1 on the system of coastal and marine Baltic Sea Protected Areas. Available online: <http://www.helcom.fi/Recommendations/Rec%2035-1.pdf>.
- HELCOM S&C 3-2015 (2015). Outcome of the Third Meeting of HELCOM State and Conservation Working Group (STATE & CONSERVATION 3-2015). Available at: <https://portal.helcom.fi/meetings/STATE-CONSERVATION%203-2015-276/MeetingDocuments/Outcome%20of%20STATE-CONSERVATION%203-2015.pdf>
- HELCOM/OSPAR (2003). Joint HELCOM/OSPAR Work Programme on Marine Protected Areas.
- ICES (2015). Fishing abrasion pressure maps for mobile bottom-contacting gears in the HELCOM area relating to both seafloor integrity and management of HELCOM MPAs. HELCOM requested advice from ICES (based on VMS/logbook data).
- IUCN World Commission on Protected Areas (IUCN-WCPA) (2008). Establishing Marine Protected Area Networks—Making It Happen. Washington, D.C.: IUCN-WCPA, National Oceanic and Atmospheric Administration and the Nature Conservancy. 118 pp.
- Piekäinen H., Korpinen S. (2007). Towards an Assessment of Ecological Coherence of the Marine Protected Areas Network in the Baltic Sea Region. BALANCE Interim Report 25. 141 pp.
- Wolters H.A., Galparsoro I., Castro R., Korpinen S., Nurmi M., Tsangaris C., Reizopoulou S., van der Meulen M., Schipper, C.A., Roeleveld G., Uriarte A., Uyarra M.C. (2015). Proposal for an assessment method of the ecological coherence of networks of marine protected areas in Europe. Deltares, Report 1208917-000-ZKS-0018 to the European Commission, 123 pp.
- UNEP-WCMC (2008). National and Regional Networks of Marine Protected Areas: A Review of Progress. UNEP-WCMC, Cambridge.

Annex 1.


Coverage of the HELCOM MPA network in 2013.

Total number, total area and marine fraction of HELCOM MPAs, as well as total marine area and protected marine area per each country. The data is based on HELCOM MPA shapefiles from March 2016, recalculated to correspond to the status of the HELCOM MPA network in 2013.

Country and number of HELCOM MPAs	Total area of HELCOM MPAs km ²	Marine fraction of HELCOM MPAs		Total marine area per country			Size of protected marine area (km ²) and fraction (%) of the national territorial waters (TW), exclusive economic zone (EEZ) and total area					
		km ²	(%)	TW km ²	EEZ km ²	Total km ²	TW km ²	(%)	EEZ km ²	(%)	Total km ²	(%)
Denmark 66	11 181	10 411	(93%)	32 917	13 074	45 991	7 599	(23%)	1 091	(8%)	10 411	(23%)
Estonia 7	7 192	6 050	(84%)	25 139	11 830	36 970	5 954	(24%)	43	(0%)	6 050	(17%)
Finland 22	5 798	5 553	(96%)	53 176	28 660	81 836	2 423	(5%)	0	(0%)	5 553	(7%)
Germany 12	5 840	5 526	(95%)	10 852	4 505	15 357	2 202	(20%)	2 217	(49%)	5 526	(36%)
Latvia 7	4 364	4 364	(100%)	12 692	16 125	28 816	4 150	(33%)	214	(1%)	4 364	(15%)
Lithuania 6	1 393	1 005	(72%)	2 274	4 259	6 534	1 393	(61%)	0	(0%)	1 005	(15%)
Poland 9	8 052	7 361	(91%)	10 172	19 491	29 663	5 426	(53%)	888	(5%)	7 361	(25%)
Russia 6	1 435	977	(68%)	16 315	7 373	23 902	894	(6%)	0	(0%)	977	(4%)
Sweden 28	8 387	7 375	(88%)	83 013	65 272	148 284	3 599	(4%)	2 750	(4%)	7 375	(5%)
Total 163	53 642	48 621	(91%)	246 550	170 589	417 352	33 640	(13.6%)	7 203	(4.2%)	48 621	(11.6%)

Annex 2.

Overview of the assessment design of the HELCOM ecological coherence assessment in 2016.



* excluded due to lack of comprehensive distribution data of species and biotope complexes in the entire Baltic Sea.
 ** the 60% target originates from discussions under the Habitats Directive, and was originally intended only for rare habitats.

Annex 3.

Overview of the ecological coherence assessment carried out for (1) the HELCOM MPA network and (2) the combined network of the HELCOM MPAs and the marine Natura 2000 sites in the Baltic Sea.

Criteria and subcriteria	(1) HELCOM MPA network	(2) combined network of the HELCOM MPAs and marine Natura 2000 sites in the Baltic Sea
I. Representativity	x	x
Geographical representation	x	x
Benthic marine landscapes	x	x
II. Replication	x	x
Marker species and biotope complexes	x	-
Benthic marine landscapes	x	x
III. Adequacy	x	-
Marine size of MPAs	x	-
Terrestrial size of MPAs	x	-
IV. Connectivity	x	x
Theoretical connectivity of benthic marine landscapes	x	x
Species-specific connectivity	x	x

Annex 4.

Overview table of the data used for the HELCOM ecological coherence assessment in 2016.

Data layer	Source and year of last update	Representativity of landscapes	Representativity, geographical distribution	Replication of marker species	Replication of marker biotope complexes	Replication of landscapes	Adequacy of MPA size	Connectivity, theoretical	Connectivity, species-specific
Shapefile of HELCOM MPAs	HELCOM Data & Map service (2016).	x	x	x	x	x	x	x	x
Shapefile of HELCOM MPAs and marine Natura 2000 sites	HELCOM Data & Map service (2016) and the EEA data bank (2015).	x	x			x		x	x
Shapefile of marine benthic landscape maps	EuSeaMap2 Interim draft Baltic Sea Broad-Scale Predictive Habitat Map (2015). ⁹	x				x		x	x
Shapefile of Baltic Sea sub-basins	HELCOM Data & Map service (2013). "HELCOM Sub-basins" Based on HELCOM Monitoring and Assessment Strategy (2013).		x						
Shapefile of coastal zones	Created for this assessment based on "HELCOM Sub-basins with coastal WFD divisions (2013)".		x						
Shapefiles of species distribution (per sub-basin)	HELCOM Species Information Sheets (2013).			x					
Shapefiles of biotope complexes distribution (100x100 km grid)	HELCOM Biotope complexes Information Sheets (2013).				x				
Species data reported per MPA	HELCOM MPA database (2015).			x					
Biotope complexes data reported per MPA	HELCOM MPA database (2015).				x				
Shapefile of spatial distribution of fisheries within the Baltic Sea during 2013 (VMS)	HELCOM Data & Map service (2015).						x		
Shapefile of spatial distribution of shipping traffic within the Baltic Sea during 2014 (AIS)	Unpublished, will be made available in HELCOM Data & Map service in 2016.						x		

⁹ To cover the whole Baltic Sea, the shapefile was amended with EuSeaMap North & Celtic Seas habitat map (including Kattegat). This dataset was supplemented with a salinity dataset from EuSeaMap project and reclassified to match the classes used in the Baltic habitat map.

Annex 5.

Fishing effort within HELCOM MPAs.

The fishing effort is given in total hours in 2013 for longlines, midwater trawl, and mobile bottom-contacting gears, based on VMS data (ICES 2015). HELCOM MPAs where no fishing occurred have been omitted from the table.

MPA	Longlines	Midwater trawl	Mobile bottom-contact gear
Adler Grund og Rønne Banke	8		9
Æbelø og havet syd for og Nærå			0
Anholt og havet nord for		7	235
Bakkebrædt og Bakkegrund			2
Bogskar		14	
Davids Banke	49	1	7
Centrale Storebælt og Vresen		14	1 028
Eckernförder Bucht mit Flachgründen, Südküste der Eckernförder Bucht und vorgelagerte Flachgründe		13	128
Fehmarnbelt		20	755
Femern Bælt			5
Finngrundet-Östra Banken		36	
Fladen			42
Flensborg Fjord, Bredgrund og farvandet omkring Als		60	2 587
Gilleleje Flak og Tragten		61	37
The open sea area southeast from Hanko		28	
Havet mellem Romsø og Hindsholm samt Romsø			
Havet og kysten mellem Hundested og Rørvig			2
Havet omkring Nordre Rønner			27
Herthas Flak			8
Hesselø med omliggende stenrev		3	
Hirsholmene, havet vest herfor og Ellinge Å's udløb		1	8
Hvideodde Rev		1	11
Irbes saurums		22	
Jasmund National Park		23	0
Kadetrinne		8	1
Kims Top og den Kinesiske Mur			1 083
Kopparstenarna/Gotska Sandön/Salvoren Area		10	
Kristiinankaupunki Archipelago		202	
Küstenbereiche Flensburger Förde von Flensburg bis Geltinger Birk, Flengurger Förde		4	131
Küstenlandschaft Bottsand - Marzkamp u. vorgelagerte Flachgründe, Östlichen Kieler Bucht		1	83
Lahemaa		36	
Lawica Slupska	125	18	8
Lilla Middelgrund		6	133
Lillebælt			78
Maden på Helnæs og havet vest for		3	
Outer Bothnian Threshold Archipelago (The Quark)		27	
Morups Bank			6
Nakskov Fjord og Inderfjord			1
Nida-Perkone			
Northern Midsjöbanken		3	
Ostoja Slowinska	2		
Pakri		616	
Pommersche Bucht-Rönnebank		261	49
Przybrzezne Wody Baltyku	270	6	46

MPA	Longlines	Midwater trawl	Mobile bottom-contact gear
Rīgas līča rietumu piekraste		249	
Røsnæs, Røsnæs Rev og Kalundborg Fjord			3
Archipelago Sea		7	
Schlei incl. Schleimünde und vorgelagerter Flachgründe			22
Schultz og Hastens Grund samt Briseis Flak		1	10
Selga uz rietumiem no Tujas		820	
Signilskär - Märket		33	
Skælskør Fjord og havet og kysten mellem Agersø og Glænø			4
Staberhuk, Großenbrode Meeresbereiche, Wagrien, Sagas-Bank			113
Stenrev sydøst for Langeland			9
Stevns Rev		1	2
Stora Middelgrund och Röde Bank			4
Strandenge på Læsø og havet syd herfor			1
Sydfynske Øhav			38
Tammisaari and Hanko Archipelago-and Pojo Bay marine protection area		33	
Torhamns Archipelago		1	
Tulliniemi bird protection area		33	
Uusikaupunki Archipelago		254	
Walkyriengrund			12
Vilsandi		6	
West-Pommeranian Lagoon National Park			1
Väinameri		83	
Zatoka Pomorska	1	6	1 346
Zatoka Pucka	0	20	37
Ålborg Bugt, østlige del		10	1

Annex 6.

Rationale for uncertainty ranking and values in the integration tables used for aggregating the subcriteria results (chapter 4.1).

Uncertainty rankings and their corresponding values are low (1), moderate (0.75) and high (0.5).

Main criterion	Subcriterion	Uncertainty value of data	Justification	Uncertainty value of target	Justification	Uncertainty value of method	Justification
Representativity	Geographical representation - subbasins	LOW (1)	The data is shapefiles of MPAs provided by countries.	LOW (1)	The target is based on recommendations from the CBD.	LOW (1)	The method is a standard GIS analysis of the shapefiles.
	Geographical representation - zonation	LOW (1)	The data is shapefiles of MPAs provided by countries.	LOW (1)	The target is based on recommendations from the CBD.	LOW (1)	The method is a standard GIS analysis of the shapefiles.
	Benthic marine landscapes	MODERATE (0.75)	The data is shapefiles derived from modelled maps produced by the EUSEaMap project. Modelled data always includes some uncertainty.	MODERATE (0.75)	The lower of two targets (20%) was chosen for the assessment. Choosing the higher target (60%) would have resulted in a lower outcome of the assessment.	LOW (1)	The method is a standard GIS analysis of the shapefiles.
Replication	Marker species	LOW (1)	Data on species comes from the MPA database, reported by the countries.	HIGH (0.5)	Theoretical target of 3 replications is based on the previous assessment (HELCOM 2010). It is rather low and a species-specific target would be better.	MODERATE (0.75)	The analysis method assumes that a species is present throughout the sub-basin where it is reported to occur, which is a rather coarse scale for such an assessment.
	Marker biotope complexes	LOW (1)	Data on biotope complexes comes from the MPA database, reported by the countries.	HIGH (0.5)	Theoretical target of 3 replications is based on the previous assessment (HELCOM 2010). It is rather low and a biotope complex-specific target would be better.	MODERATE (0.75)	The analysis method assumes that a biotope complex is present throughout the 100x100 square where it is reported to occur, which is a rather coarse scale for such an assessment.
	Benthic marine landscapes	MODERATE (0.75)	The data is shapefiles derived from modelled maps produced by the EUSEaMap project. Modelled data always includes some uncertainty.	HIGH (0.5)	Theoretical target of 3 replications is based on the previous assessment (HELCOM 2010). It is rather low and a landscape-specific target would be better.	HIGH (0.5)	The method is easily biased by the scale of the data. The fine scale of the current data makes the outcome overly positive.

Main criterion	Subcriterion	Uncertainty value of data	Justification	Uncertainty value of target	Justification	Uncertainty value of method	Justification
Adequacy	MPA marine size	LOW (1)	The data is shapefiles of MPAs provided by countries.	MODERATE (0.75)	The target was set by S&C 3-2015 meeting, and could perhaps be too high (depending on how important size is seen for efficient protection).	LOW (1)	The method is a standard GIS analysis of the shapefiles.
	MPA terrestrial size	LOW (1)	The data is shapefiles of MPAs provided by countries.	MODERATE (0.75)	The target was set by S&C 3-2015 meeting, and could perhaps be too high (depending on how important size is seen for efficient protection).	HIGH (0.5)	The method might possibly be skewed by small islands inside larger marine MPAs which are interpreted as their own sites.
Connectivity	Theoretical connectivity	MODERATE (0.75)	The data is shapefiles derived from modelled maps produced by the EUSEaMap project. Modelled data always includes some uncertainty.	HIGH (0.5)	Based on recommendations in the Deltares report (2015), but no further reasoning was found for the amount of connections.	HIGH (0.5)	The GIS analysis is complex and easily biased by the scale of the data.
	Species-specific connectivity	MODERATE (0.75)	The data is shapefiles derived from modelled maps produced by the EUSEaMap project. Modelled data always includes some uncertainty.	MODERATE (0.75)	Based on recommendations in the Deltares report (2015), but no further reasoning was found for the amount of connections. However, species-specific dispersal distances are reliable.	HIGH (0.5)	The GIS analysis is complex and easily biased by the scale of the data.

Annex 7.

Follow up of HELCOM Recommendation 35/1 on the system of coastal and marine Baltic Sea protected areas (HELCOM MPAs¹⁰).

Outline of the follow-up of actions in recommendation 35/1. Please note, that the rules for assessing the level of accomplishment (green, yellow, red) refer to the regional level. NA = Not applicable.

¹⁰ Former BSPAs.

Action from Recommendation 35/1	Level of implementation	Indicator/required information for national reporting	Indicator for aggregation at regional level	Accomplished (Regional)	Partly accomplished (Regional)	Not accomplished (Regional)
a) reach the target set by the HELCOM 2010 Moscow Ministerial Declaration that at least 10% of the marine area in all sub-basins of the Baltic Sea including the EEZ areas beyond territorial waters is covered by MPAs, where scientifically justified;	National	National reporting of numerical/geographical data on MPAs, as required to the HELCOM MPA database.	% area covered by MPAs in the respective sub-basin, in particular in EEZ areas beyond territorial waters.	At least 10% of the marine area in all sub-basins of the Baltic Sea, including the EEZ areas beyond territorial waters, is covered by MPAs (presented per sub-basin).	10% of the marine area is covered by MPAs in some sub-basins, including the EEZ areas beyond territorial waters (presented per sub-basin).	Less than 10% of the marine area is covered by MPAs in all sub-basins, including the EEZ areas beyond territorial waters (presented per sub-basin).
b) designate new sites as HELCOM MPAs, where ecologically meaningful, especially in offshore areas beyond territorial waters;	National/ Regional	National reporting of numerical/geographical data on MPAs, as required to the HELCOM MPA database.	Increase in the nr and % of MPAs in offshore areas beyond territorial waters.	The nr and/or % of MPAs in offshore areas is increasing in all sub-basins where the 10% aim is not met <i>(will level off and become redundant when the 10% aim is reached)</i>	The nr and/or % of MPAs in offshore areas is increasing in some sub-basins where the 10% aim is not met <i>(will level off and become redundant when the 10% aim is reached)</i>	The nr and/or % of MPAs in offshore areas is not increasing.
c) ensure that HELCOM MPAs inter alia provide specific protection to those species, habitats, biotopes and biotope complexes included in the HELCOM Red Lists, as agreed upon in the HELCOM 2013 Copenhagen Ministerial Declaration, by considering these in the site selection procedure;	National	National reporting of numerical/geographical data and information on MPAs, as required to the HELCOM MPA database.	Nr of threatened species present in MPAs. Nr and area of threatened biotopes/habitats present in MPAs.	1) The HELCOM network of MPAs provides protection to all HELCOM threatened species. 2) The HELCOM network of MPAs provides protection to all HELCOM threatened biotopes/habitats. (Both statements to be met for the recommendation to be accomplished).	1) The HELCOM network of MPAs provides protection to some HELCOM threatened species. 2) The HELCOM network of MPAs provides protection to some HELCOM threatened biotopes/habitats.	The HELCOM network of MPAs does not provide any known protection to HELCOM threatened species, biotopes, biotope complexes and habitats.
d) ensure when selecting new areas, that the network of HELCOM MPAs is ecologically coherent and takes into account connectivity between sites, including, for example, migration routes, species mobility and areas of special ecological significance, such as spawning areas;	National	National reporting of numerical/geographical data and information on MPAs, as required to the HELCOM MPA database.	Regular regional evaluation of the ecological coherence of HELCOM MPAs, using agreed criteria for assessment.	The network of HELCOM MPAs is assessed as being ecologically coherent.	Some criteria considered in the evaluation are reflecting ecological coherence.	None of the criteria used in the assessment of the network of HELCOM MPAs are reflecting ecological coherence.

Action from Recommendation 35/1	Level of implementation	Indicator/required information for national reporting	Indicator for aggregation at regional level	Accomplished (Regional)	Partly accomplished (Regional)	Not accomplished (Regional)
h) 1) develop and apply by 2015 management plans or measures for all existing HELCOM MPAs;	National	National reporting on the status of management plans as required to the HELCOM MPA database.	Nr or % of MPAs with management plans or measures.	All HELCOM MPAs designated by 2014* have management plans or measures. * year of adoption of recommendation	> 67%* of HELCOM MPAs designated by 2014 have management plans or measures. *% MPAs with management plans in 2013 HELCOM assessment prepared for Ministerial Meeting (no new areas designated by 2014).	67%* of HELCOM MPAs designated by 2014 have management plans or measures. *% MPAs with management plans in 2013 HELCOM assessment prepared for Ministerial Meeting.
h) 2) and establish management plan or measures for every new MPA within five years after its designation;	National	National reporting on the status of management plans as required to the HELCOM MPA database.	Nr or % of new MPAs with management plans within 5 years of designation.	All HELCOM MPAs designated after 2014 have a management plan within 5 years of designation.	Some HELCOM MPAs designated after 2014 have a management plan within 5 years of designation.	No HELCOM MPAs designated after 2014 have a management plan within 5 years of designation.
i) update the management plans when necessary and in accordance with other legal requirements with a maximum of 12 years intervals;	National	National reporting on the implementation date of current management plans as required to the HELCOM MPA database.	All management plans are updated with a maximum of 12 years interval.		Some management plans are updated with 12 years interval.	No management plans are updated after 12 years.
k) assess the effectiveness of the management plans or measures of HELCOM MPAs by conducting monitoring, and where feasible scientific research programmes, which are directly connected to the conservation interests of HELCOM MPAs, including the placement of monitoring stations inside the MPAs;	National/ Regional	National reporting on monitoring within MPAs as required to the HELCOM MPA database.	Nr of MPAs with monitoring stations within the designated area.	All (or certain %) HELCOM MPAs has monitoring stations with the designated area. Assessment of the management effectiveness has been carried out.	Some HELCOM MPAs has monitoring stations with the designated area. Assessment of management effectiveness ongoing.	No HELCOM MPA has monitoring station with the designated area. Assessment management effectiveness not started.

In addition to the paragraphs assessed in this report, the following paragraphs are included in the Recommendation 35/1.

- e) make use of computer-based site selection tools, such as MARXAN for a HELCOM-wide approach which maximizes the chance of creating a coherent network of HELCOM MPAs and, at the same time, minimizing the impact of pressures and conflicts with other interests;
- j) harmonise the designation of neighbouring HELCOM MPAs in transboundary marine areas, and, where appropriate, to join forces between neighbouring states when setting up management plans or measures for such HELCOM MPAs
- l) include HELCOM MPAs as areas of particular ecological significance in coastal and maritime spatial planning processes and incorporate their management provisions in spatial plans and Integrated Marine and Coastal Management Strategies, respectively
- m) update, when necessary, HELCOM MPA related guidelines and guiding documents, in order to keep them in line with new knowledge and compatible with other
- o) perform identification, designation and legal protection of HELCOM MPAs, according to HELCOMs criteria and guidelines and base all management plans or measures on relevant HELCOM publications, such as "Planning and management of Baltic Sea Protected Areas: guidelines and tools" (BSEP 105). For EU Member States, the respective EU requirements and guidelines are regarded as being adequate for designating and managing HELCOM MPAs;
- q) continuously report the most recent numerical and descriptive data on HELCOM MPAs to HELCOMs data systems (HELCOM MPA database, GIS based map and data service)
- r) regularly assess the status and development of HELCOMs MPAs, according to the time tables set by HELCOM and to ensure that the assessments are applicable for corresponding EU and global reporting



www.helcom.fi