HELCOM Monitoring Programme topic

Concentrations of contaminants

Programme:

Contaminants in biota

Contents

a.	Metadata on monitoring strategies and monitoring programmes	2
	a.1 Responsible HELCOM subsidiary body	2
	a.2 Regional Cooperation (RegionalCooperation)	3
b.	Monitoring strategies	3
	b.1 Descriptor	3
	b.2 BSAP segments	4
	b.3 Monitoring strategy description	4
	b.4 BSAP Ecological objectives	4
	b.5 Gaps in monitoring	5
c.	Monitoring programmes	5
	c.1 Purpose of monitoring	5
	c.2 Other legislation	9
	c.3 Implementation of Regional Cooperation (RegionalCooperation_implementation)	10
	c.4 Monitoring concepts	11
	c.5 Monitoring and assessment requirements	27
	c.6 Data providers and access	28
	c.7 MSFD Criteria (GES Criteria)	31
٦	Poforoneos	20

a. Metadata on monitoring strategies and monitoring programmes

a.1 Responsible HELCOM subsidiary body

Please indicate the relevant expert group/network if available, otherwise the responsible HELCOM Working Group. or the responsible working group understanding is that they are the EN or EG if available, otherwise the WGs

	Permament Groups	
	Gear – Group on the Implementation of the Ecosystem Approach	
	Maritime – Maritime Working Group	
	Pressure – Working Group on Reduction of Pressures from the Baltic Sea Catchment Area	
	Response – Response Working Group	
	State and Conservation — Working Group on the State of the Environmental and Nature Conservation	
	Time-limited Groups	
	Agri – Group on Sustainable Agricultural Practices	
	Fish – Group on Ecosystem-based Sustainable Fisheries	
	HELCOM-VASAB MSP WG - Joint HELCOM-VASAB Maritime Spatial Planning Working Group	
	0.044	
	Expert Groups	
	Expert Groups	
_	Expert Groups AIS EWG — Expert Working Group for Mutual Exchange and Deliveries of AIS data	
_	Expert Groups AIS EWG – Expert Working Group for Mutual Exchange and Deliveries of AIS data EN Hazardous Substances – Expert Network on hazardous substances	
_	Expert Groups AIS EWG – Expert Working Group for Mutual Exchange and Deliveries of AIS data EN Hazardous Substances – Expert Network on hazardous substances EN Marine Litter – Expert Network on Marine Litter	
_	Expert Groups AIS EWG – Expert Working Group for Mutual Exchange and Deliveries of AIS data EN Hazardous Substances – Expert Network on hazardous substances EN Marine Litter – Expert Network on Marine Litter EN Noise – Expert Network on Underwater Noise	
_	Expert Groups AIS EWG – Expert Working Group for Mutual Exchange and Deliveries of AIS data EN Hazardous Substances – Expert Network on hazardous substances EN Marine Litter – Expert Network on Marine Litter EN Noise – Expert Network on Underwater Noise ESA – Expert Network on Economic and Social Analyses	
_	Expert Groups AIS EWG – Expert Working Group for Mutual Exchange and Deliveries of AIS data EN Hazardous Substances – Expert Network on hazardous substances EN Marine Litter – Expert Network on Marine Litter EN Noise – Expert Network on Underwater Noise ESA – Expert Network on Economic and Social Analyses EWG OWR – Expert Working Group on Oiled Wildlife Response	

	Management Convention Exemptions
	IN Benthic habitat – Intersessional Network on habitat monitoring
	IWGAS – Informal Working Group on Aerial Surveillance
	JWG Bird – HELCOM-OSPAR-ICES Joint Working Group on Seabirds
\boxtimes	MORS EG – Expert group on monitoring of radioactive substances in the Baltic Sea
	PRF Cooperation Platform – Cooperation Platform on Port Reception Facilities in the Baltic Sea
	SAFE NAV – Group of Experts on Safety of Navigation
	SUBMERGED – Expert Group on Environmental Risks of Hazardous Submerged Objects
⊠ Fully co	ng of this programme is: pordinated coordinated. Indicate missing component(s):
•	
	nated monitoring is under development. Indicate by which group/project and by when a ndation on coordinated monitoring can be expected.
	mon monitoring guidelines: HELCOM COMBINE manual, MORS Guidelines and guidelines (see References).
DIN meth manu	mon quality assurance programme: HELCOM COMBINE manual, QUASIMEME, EN ISO/IEC 17025. Radioacve substances: MORS Guidelines defines odologies for sample treatment, analysis and intercomparison. Reported data is rally quality assured by HELCOM secretariat and results reported and verified in al MORS EG meeng.
- Comr	mon databases: ICES DOME and MORS.
b. Moni	itoring strategies
b.1 Desc The programm boxes.	riptor ne supports the following obligatory MSFD Monitoring Strategies. Tick one or more relevant
□ D1	Biodiversity
□ n 2	Non-indigenous Species

□ D3	Commercial fish and shellfish
□ D4	Food webs
□ D 5	Eutrophication
□ D 6	Seafloor integrity
□ D7	Hydrographical conditions
⊠ D8	Contaminants
⊠ D9	Contaminants in seafood
□ D10	Marine litter
□ D11	Energy including underwater noise
b.2 BSAP s The sub-program	segments nme serves the following BSAP segments. Tick one or more relevant boxes.
□Eutrophicatio	า
⊠ Hazardous sub	ostances
\square Biodiversity	
☐ Maritime activ	vities
b.3 Monito	oring strategy description
	tegy: Monitoring is to be carried out to fulfill assessment requirements of HELCOM cives that are specified through HELCOM core indicators. The requirements on include number of stations, the sampling frequency and replication.
	Ecological objectives most relevant option(s). Tick one or more boxes below.
Eutrophication	☐ Concentrations of nutrients close to natural levels
	☐ Clear water
	\square Natural level of algal blooms
	$\hfill\square$ Natural distribution and occurrence of plants and animals
	☐ Natural oxygen levels
Hazardous substances	□ Concentrations of hazardous substances close to natural levels
SUDSTAILES	

	☐ Healthy wildlife
	☐ Radioactivity at pre-Chernobyl levels
Biodiversity	☐ Natural landscapes and seascapes
	\square Thriving and balanced communities of plants and animals
	☐ Viable populations of species
Maritime activities	☐ No illegal pollution
activities	\square Safe maritime traffic without accidental pollution
	☐ Efficient response capability
	\square No introductions of alien species from ships
	☐ Minimum air pollution from ships
	\square Zero discharges from offshore platforms
Sufficient covera ☐ Adequate mo ☐ Monitoring is	e GES criteria addressed, indicate when sufficient monitoring was in place or by when ge will be in place (Coverage_GEScriteria) nitoring was in place in 2014 nitoring was in place by 2018 nitoring is in place by July 2020 nitoring will be in place by 2024 not being put in place for this descriptor due to a low risk r this descriptor is not relevant
this descriptor m	te implementation gaps and plans to complete the establishment and implementation of nonitoring strategy (Gaps_Plans): carried out in the indicator assessment regarding areas lacking data and areas o scared to be statistically robust.

c. Monitoring programmes

c.1 Purpose of monitoring

c.1a Assessment purpose in general

The programme supports the assessment of:

Tick the relevant box

TICK THE FEIEVAIL BOX.		
Temporal trends	Spatial distribution	State classification
NZ		∇
		<u> </u>

The **programme** supports the assessment of: (MonitoringPurpose).

Note that the answer to this question will be decisive for whether to answer upcoming questions e.g. upcoming questions on pressures should only be answered if the monitoring is defined as supporting the assessment of pressures.

Tick the relevant boxes.

Environmental state and impacts	Pressures in the marine environment	Pressures at source (land-based, riverine, sea-based ¹ and atmospheric sources)	Human activities causing the pressures	Effectiveness of measures
If this is selected fill in the following questions: c.1b	If this is selected fill in the following questions: c.1c, d	If this is selected fill in the following questions: c.1c, d		If this is selected fill in the following questions: c.1c, d

Give any other monitoring purpose e.g. if the programmes include supporting parameters for other monitoring programmes

Supporting also health assessments of biota and pelagic habitats as well as food safety.

For questions 1b-1d, select when applicable for the sub-programme, the link from the Reporting on the 2020 update of Article 11 for the Marine Strategy Framework Directive (MSFD Guidance Document 17, 2020) (Features) to:

- Ecosystem components (relevant for monitoring and assessment for Article 8(1a) for D1C2-C5, D3, D4, D6C3-C5, D7C2)
- Pressures and impacts in the marine environment (relevant for monitoring and assessment for Article 8(1b) for D1C1, D2, D5, D6C1-C2, D7C1, D8, D9, D10, D11)
- Pressure inputs to the marine environment (relevant for monitoring and assessment for Article 10)
- Uses and human activities (relevant for monitoring and assessment for Article 8(1c) and 13)

c.1b • Ecosystem components (Features)

Choose only the most relevant option(s). Tick one or more boxes below.

Theme	Sub-theme	Label feature
Species	☐ Birds	☐ Grazing birds
		☐ Wading birds
		☐ Surface-feeding birds
		☐ Pelagic-feeding birds
		☐ Benthic-feeding birds

¹ Sea-based 'Pressures at source' refers to monitoring pressures from sea-based activities where the monitoring is directly at the activity rather than at a distance from or time period after it is generated by the activity (e.g. D1 incidental by-catch when fishing, D2 ballast water discharges, D6 use of bottom fishing gear, D8 contaminant discharges and pollution events from a vessel or pipeline, D11 impulsive sound events from a vessel or platform).

	☐ Mammals	☐ Small toothed cetaceans		
		\square Deep-diving toothed cetaceans		
		☐ Baleen whales		
		□ Seals		
	☐ Reptiles	☐ Turtles		
	⊠ Fish	☐ Coastal fish		
		☐ Pelagic shelf fish		
		\square Demersal shelf fish		
		☐ Deep-sea fish		
		\square Commercially exploited fish and shellfish		
	☐ Cephalopods	\square Coastal/shelf cephalopods		
		☐ Deep-sea cephalopods		
Habitats	☑ Benthic habitats	\square Benthic broad habitats		
		\square Other benthic habitats		
	☑ Pelagic habitats	☐ Pelagic broad habitats		
		☐ Other pelagic habitats		
Ecosystems		characteristics		
	□ Chemical characteristics			
	☐ Ecosystems, including food webs	☐ Coastal ecosystems		
		☐ Shelf ecosystems		
		☐ Oceanic/deep-sea ecosystems		
	• Pressures and impacts in the most relevant option(s). Ti	the marine environment (Features) ck one or more boxes below.		
Theme	Label: Feature			
Biological	☐ Newly introduced non-	-indigenous species		
	☐ Established non-indigenous species			
	\square Species affected by inc	idental by-catch		
Physical and	d 🗆 Hydrographical changes			
hydrological	☐ Physical disturbance to seabed			
	☐ Physical loss of the seabed			
Substances,	☐ Eutrophication			
litter and		☐ Contaminants - non UPBT substances		
energy	□ Contaminants - UPBT substances			

	⊠ Contaminants – in seafood		
	☑ Adverse effects on species or habitats		
	☐ Acute pollution events		
	☐ Litter in the environment		
	☐ Impulsive sound in water		
	☐ Continuous low frequency sound		
c.1d • P	ressure inputs to the marine environment (Features)		
Theme	Label: Feature		
Biological	☐ Input or spread of non-indigenous species		
	☐ Input of microbial pathogens		
	☐ Input of genetically modified species and translocation of native species		
	$\hfill\Box$ Loss of, or change to, natural biological communities due to cultivation of animal or plant species		
	$\hfill\Box$ Disturbance of species (e.g. where they breed, rest and feed) due to human presence		
	☐ Extraction of, or mortality/injury to, wild species (by commercial and recreational fishing and other activities)		
Substances,	☐ Input of nutrients — diffuse sources, point sources, atmospheric deposition		
litter and energy	☐ Input of organic matter — diffuse sources and point sources		
chergy	☑ Input of other substances (e.g. synthetic substances, non-synthetic substances, radionuclides) — diffuse sources, point sources, atmospheric deposition, acute events		
	☐ Input of litter (solid waste matter, including micro-sized litter)		
	☐ Input of anthropogenic sound (impulsive, continuous)		
	☐ Input of other forms of energy (including electromagnetic fields, light and heat)		
	☐ Input of water — point sources (e.g. brine)		
c.1e • U	ses and human activities (Features)		
Choose only the	most relevant option(s). Tick one or more boxes below.		
Theme Label: Feature			
Physical	☐ Land claim		
restructuring of rivers, coastline	☐ Canalisation and other watercourse modifications		
or seabed (wate	T Coastal defence and flood protection		

management)	☐ Offshore structures (other than for oil/gas/renewables)
	☐ Restructuring of seabed morphology, including dredging and depositing of materials
Extraction of	☐ Extraction of minerals (rock, metal ores, gravel, sand, shell)
non-living resources	☐ Extraction of oil and gas, including infrastructure
resources	☐ Extraction of salt
	☐ Extraction of water
Production of energy	\square Renewable energy generation (wind, wave and tidal power), including infrastructure
	☐ Non-renewable energy generation
	☐ Transmission of electricity and communications (cables)
Extraction of	☐ Fish and shellfish harvesting (professional, recreational)
living resources	☐ Fish and shellfish processing
	☐ Marine plant harvesting
	☐ Hunting and collecting for other purposes
Cultivation of	☐ Aquaculture — marine, including infrastructure
living resources	☐ Aquaculture — freshwater
	□Agriculture
	☐ Forestry
Transport	☐ Transport infrastructure
	☐Transport — shipping
	□Transport — air
	☐ Transport — land
Urban and	□Urban uses
industrial uses	□Industrial uses
	☐ Waste treatment and disposal
Tourism and	☐ Tourism and leisure infrastructure
leisure	☐ Tourism and leisure activities
Security/defence	☐Military operations (subject to Article 2(2))
Education and research	☐ Research, survey and educational activities

c.2 Other legislation

The sub-programme links with the following other international legislation (OtherPoliciesConventions). Tick one or more relevant boxes.

☐ Bathing Water Directive
☐ Common Fisheries Policy and Data Collection Framework
☐ Habitats Directive
☐ Birds Directive
□ Nitrates Directive
☐ Urban Waste Water Treatment Directive
⊠Water Framework Directive
⊠OSPAR Convention
☐Trilateral Wadden Sea Convention
☑Other, Specify: Maritime Spatial Planning Directive; Environmental Quality Standards Directive (2008/105/EC, amended by 2013/39/EU); Regulation (EC) No 1881/2006 on certain contaminants in foodstuffs
c.3 Implementation of Regional Cooperation (RegionalCooperation_implementation) Indicate the level of implementation by selecting one of the following:
☐ Agreed data collection methods
☐Common monitoring strategy (spatial and temporal design of programme)
⊠Coordinated data collection (delivered separately by each country)
\Box Joint data collection (multinational delivery using same platform and/or algorithms)

c.4 Monitoring concepts

Monitoring concepts table²:

Current means of coordination	Features or Elements	Parameter	Method	QA/QC	Frequenc y ³	Spatial resolution (density) of sampling	Link to HELCOM core indicators ⁴	Spatial scope	Monitorin g started (year)	CPs monitoring ⁵
	Elements (Features) (Features_enu m)	Parameters (Parameter) (ParametersOther)	MonitoringMethod (Monitoring Method) MonitoringMethodOthe r)	(Free text)	MonitoringF requency	(ProgrammeDescripti on)	(RelatedIndicator) (RelatedIndicator_n ame	(SpatialSco pe)	(TemporalSc ope)	(CountryCode_E num)
Regional (Combine)	PCBs, dioxins and furans	Concentration of chemical/nutrient/pollutant in biota	HELCOM COMBINE manual, PartD and different approaches e.g. CEMP manual, ICES guidelines, ISO/CEN standards (see References)	HELCOM COMBINE manual Part B, Annex B12, Appendix 1 and 3, QUASIMEME and DIN EN ISO/IEC 17025	Other	See map for details	Polychlorinated biphenyls and dioxins and furans	Coastal waters/ transitio nal waters/ EEZ	DE: 1995 DK: 1998 EE: 1996 FI: 1998 PL: 1998 SE: 1979	DE, DK, LT, EE, FI, PL, SE
Regional (Combine)	РАН	Concentration of chemical/nutrient/ pollutant in biota	HELCOM COMBINE manual, PartD and different approaches e.g. CEMP manual, ICES guidelines, ISO/CEN standards (see References)	HELCOM COMBINE manual Part B, Annex B12, Appendix 1 and 2, QUASIMEME and DIN EN ISO/IEC 17025	Other	See map for details	Polyaromac hydrocarbons and their metabolites	Coastal waters/ transitio nal waters/ EEZ	EE:2002 PL: 2012 SE: 1984	DE, DK, LT, EE, PL, SE

² Needed codelists can be found on 2020 update of Article 11 for the Marine Strategy Framework Directive (MSFD Guidance Document 17, 2020).

³ The option "Different for each country - see MORE overview" refers to the <u>overview</u> carried out in 2013

⁴ Give the name of HELCOM core indicators that are based on the monitoring parameter.

⁵ Provide information on the Contracting Partie(s) that are monitoring the parameter.

Current means of coordination	Features or Elements	Parameter	Method	QA/QC	Frequenc y ³	Spatial resolution (density) of sampling	Link to HELCOM core indicators ⁴	Spatial scope	Monitorin g started (year)	CPs monitoring ⁵
National	PBDE	Concentration of chemical/nutrient/ pollutant in biota	Different approaches e.g. CEMP manual, ICES guidelines, ISO/CEN standards (see References)	QUASIMEME and DIN EN ISO/IEC 17025	Other	See map for details	PBDE	Coastal waters/ transitio nal waters/ EEZ	DE: 2007 DK: 2004 FI: 2012 EE: 2002 PL: 2012 SE: 1969 (guillemot egg) 1980 (fish)	DE, DK. FI, EE, PL, SE
National	PFOA, PFOS and PFOSA	Concentration of chemical/nutrient/pollutant in biota	Different approaches e.g. CEMP manual, ICES guidelines, ISO/CEN standards (see References)	QUASIMEME and DIN EN ISO/IEC 17025	Other	See map for details	PFOS	Coastal waters/ transitio nal waters/ EEZ	DE: 2011 DK: 2011 FI: 2012 EE: 2002 PL: 2014 SE: 1973 (guillemot egg), 1980 (fish)	DE, DK, FI, EE, PL, SE
National	HBCDD	Concentration of chemical/nutrient/pollutant in biota	Different approaches e.g. CEMP manual, ICES guidelines, ISO/CEN standards (see References)	QUASIMEME and DIN EN ISO/IEC 17025	Other	See map for details	HBCD	Coastal waters/ transitio nal waters/ EEZ	DE: 2011 DK: 2011 FI: 2012 EE: 2002 PL: 2012 SE: 1969 (guillemot egg) 1980 (fish)	DE, DK, FI, EE, PL, SE

Current means of coordination	Features or Elements	Parameter	Method	QA/QC	Frequenc y ³	Spatial resolution (density) of sampling	Link to HELCOM core indicators ⁴	Spatial scope	Monitorin g started (year)	CPs monitoring ⁵
Regional (Combine)	Tributyltin (TBT)	Concentration of chemical/nutrient/pollutant in biota	HELCOM COMBINE manual, PartD and different approaches e.g. CEMP manual, ICES guidelines, ISO/CEN standards (see References)	HELCOM COMBINE manual, QUASIMEME and DIN EN ISO/IEC 17025	Other	See map for details	TributyIn (TBT) and imposex	Coastal waters/ transitio nal waters/ EEZ	DE: 2011 DK: 1998 EE: 2002 PL: 2012 SE: 2008, sporadic years reported before 2008	DE, DK, EE, PL, SE
Regional (Combine)	Metals	Concentration of chemical/nutrient/pollutant in biota	HELCOM COMBINE manual, PartD and different approaches e.g. CEMP manual, ICES guidelines, ISO/CEN standards (see References)	HELCOM COMBINE manual Part B, Annex B12, Appendix 1, 4 and 5, QUASIMEME and DIN EN ISO/IEC 17025	Other	See map for details	Metals (lead, cadmium and mercury)	Coastal waters/ transitio nal waters/ EEZ	DE: 1995, data also reported for 1992 DK: 1998 EE: 1996, some data also from 1986 FI: 1998 LT: 2007 PL: (fish, mussel:Hg, Pb, Cd, Zn, Cu-1998; As -2020;) SE: 1975 (guillemot egg) 1980 (gerring)	DE, DK, EE, FI, LT, PL, SE

Current means of coordination	Features or Elements	Parameter	Method	QA/QC	Frequenc y ³	Spatial resolution (density) of sampling	Link to HELCOM core indicators ⁴	Spatial scope	Monitorin g started (year)	CPs monitoring ⁵
National	BDE	Concentration of chemical/nutrient/pollutant in biota	Different approaches e.g. CEMP manual, ICES guidelines, ISO/CEN standards (see References)	Other	Other			EEZ	DK: 2006, 2008	DK
National	1. Organochlori ne pesticides (DDT and its metabolites, hexachloroc yclohxsane, hexachlorob enzene, endosulfan) 2. Aquatic plants - Metals:cadm ium (Cd), lead (Pb), mercury (Hg), nickel (Ni), cooper (Cu) i zinc (Zn)	Concentration of chemical/nutrient/pollutant in biota	Different approaches e.g. CEMP manual, ICES guidelines, ISO/CEN standards , national	Other	Other			Coastal waters/ transitio nal waters/ / EEZ	PL: 1.1998 - Yearly at all stations (from 2020: 7 stations - fish; 1 station - mussels); 2. 2012 - yearly /4 stations/1- 7 samples aquatic plants	PL
Regional (MORS)	Radionucli des: gammaemi ers	Concentration of chemical/nutrient/pollutant in biota	MORS Guidelines	MORS Guidelines	Yearly	Yearly, See map for details	Radioactive substances: Cs137 in fish and surface waters	EEZ	1984 EE: 1997	DE, DK, EE, FI, LT, PL, SE

Element/Parameter pair

PCB, Dioxins, furans / Concentration of chemical/pollutant in biota

METHOD (Monitoring Details)

Element/parameter

Measured in: In tissue of Platichthys flesus, Zoarces viviparous (in Denmakr, Zoarces vivparous is often not available anymore due to competetion from Neogobius melanostomus, which is used in fjords where Zoarces are not caught anymore), Mytilus edulis, Perca fluvialis, Clupea harengus, Limanda limanda (by DE in Kiel Bay). In the eggs of Uria aalge, Sterna hirundo, Haematopus ostralegus.

Sampling and analytical methods are reported per sample and per parameter respecvely in the data. HELCOM COMBINE manual.

EE: measured in Perca fluviatilis in WFD coastal water bodies, Clupea harengus in offshore areas (other fish species are also included for food safety (D9) monitoring.

QA/QC

Element/Parameter pair

Quality assurance is a laboratory's whole sampling and analytical process from start to finish. See HELCOM COMBINE manual.Quality assurance is a laboratory's whole sampling and analytical process from start to finish. See HELCOM COMBINE manual.

FREQUENCY

Frequency

Element/Parameter pair

DE - UBA: kelb , mussels: Baltic Sea: 2x/year(06/12), annually LUNG: mussels, every 2nd or 3rd year LLUR: 2x/yr (Mar/Oct)

FI: 1-6 years intervals, depending on staon; me series staons yearly

EE: in coastal waters 1 or 2 times per WFD 6-years cycle; in EEZ trend stations yearly

SE: Yearly at all stations and 2 times per year at two stations

DK: Yearly in fish and every second year in mussels

PL: Yearly at all stations (from 2020: 7 stations - fish; 1 station - mussels)

SPATIAL SCOPE

Spatial Scope

Element/Parameter pair

Coastal waters/ EEZ / Whole Baltic Sea for assessment

SPATIAL RESOLUTION (DENSITY) OF SAMPLING

Spatial resolution

Element/Parameter pair

Polychlorinated biphenyls (PCB) and dioxins and furans are measured in the following subbasins: Great Belt, Little Belt, Kattegat, Southern Baltic Proper, Kiel Bay, Bay of Mecklenburg, Arkona Basin, Bothnian Bay, Bothnian Sea, Northern Baltic Proper, Gulf of Gdansk, Gulf of Riga, Gulf of Finland, Quark, Archipelago Sea.

See map for details.

Element/Parameter pair

PAH / Concentration of chemical/pollutant in biota

METHOD (Monitoring Details)

Element/parameter

FI, DK (Mytilus edulis) and SE: Monitored in bivalves

EE: Monitored in bivalves - Mytilus trossulus.

PL: PAHs monitored in bivalves (Mytilus edulis) and their metabolites in bile on fish (perch, herring and flounder) See HELCOM COMBINE manual.

QA/QC

Element/Parameter pair

See HELCOM COMBINE manual

FREQUENCY

Frequency

Element/Parameter pair

FI: Survey type

SE: Yearly

EE: in coastal waters 1 or 2 times per WFD 6-years cycle

DK: Yearly

PL: Yearly at all stations (from 2020: 7 stations - fish; 1 station - mussels)

SPATIAL SCOPE

Spatial Scope

Element/Parameter pair

Coastal waters/ transitional waters/ EEZ

FI: Coastal sites

EE: Coastal sites

SE: Coastal sites

PL: Coastal waters/ transitional waters/ EEZ

SPATIAL RESOLUTION (DENSITY) OF SAMPLING

Spatial resolution

Element/Parameter pair

See map for details

Coastal Waters / EEZ

PARAMETER

Element/Parameter pair

PBDE / Concentration of chemical/pollutant in biota

METHOD (Monitoring Details)

Element/parameter

Measured in: Plachthys flesus, Zoarces viviparous, Mytilus edulis, Clupea harengus, Gadus morhua, Uria aalge, Sterna hirundo, Haematopus ostralegus, Limanda limanda (by DE in Kiel Bay), Perca fluvialis. Sampling and analytical methods are reported per sample and per parameter respectively in the data.

QA/QC

Element/Parameter pair

Quality assurance is a laboratory's whole sampling and analycal process from start to finish.

FREQUENCY

Frequency

Element/Parameter pair

DE:

• UBA: kelb , mussels: Baltic Sea: 2x/year(06/12), annually

• LUNG: mussels, every 2nd or 3rd year

LLUR: 2x/yr (Mar/Oct)

FI: 1-6 years intervals, depending on station; time series stations yearly

EE: in coastal waters 1 or 2 times per WFD 6-years cycle; in EEZ trend stations yearly

SE: Annually at all stations and 2 times per year at two stations

DK: Annually in fish

PL: Yearly at all stations (from 2020: 7 stations - fish; 1 station - mussels)

SPATIAL SCOPE

Spatial Scope

Element/Parameter pair

Coastal Waters / transitional waters/ EEZ / Whole Baltic Sea for assessment.

SPATIAL RESOLUTION (DENSITY) OF SAMPLING

Spatial resolution

Element/Parameter pair

See map for details

PBDE is measured in the following areas: Great Belt, Little Belt, Kattegat, Kiel Bay, Arkona Basin, Southern Baltic Proper, Gulf of Gdansk, Gulf of Riga, Bothnian Bay, the Quarck, Bothnian Sea, Archipelago Sea, Northern Baltic Proper, Gulf of Finland.

Element/Parameter pair

PFOA, PFOS, PFOSA / Concentration of chemical/pollutant in biota

METHOD (Monitoring Details)

Element/parameter

Measured in the following species: Platichthys flesus, Zoarces viviparous, Clupea harengus, Uria aalge, Sterna hirundo, Haematopus ostralegus, Perca fluvialis. Sampling and analytical methods are reported per sample and per parameter respectively in the data.

QA/QC

Element/Parameter pair

Quality assurance is a laboratory's whole sampling and analytical process from start to finish.

FREQUENCY

Frequency

Element/Parameter pair

Yearly DE:

• UBA: kelb , mussels: Baltic Sea: 2x/year(06/12), annually

• LUNG: mussels, every 2nd or 3rd year

LLUR: 2x/yr (Mar/Oct)

EE: in coastal waters 1 or 2 times per WFD 6-years cycle; in EEZ trend stations yearly

SE: Annually at all stations

DK: Annually in fish

PL: Yearly at all stations (from 2020: 7 stations - fish)

SPATIAL SCOPE

Spatial Scope

Element/Parameter pair

Coastal Waters / transitional waters/ EEZ / Whole Baltic Sea for assessment.

SPATIAL RESOLUTION (DENSITY) OF SAMPLING

Spatial resolution

Element/Parameter pair

PFOA, PFOS and PFOSA are measured in the following areas: Great Belt, Kattegat, Little Belt, Southern Baltic Proper, Bothnian Bay, Northern Baltic Proper, Bothnian Sea, Gulf of Riga, Gulf of Finland, Gulf of Gdansk. See map for details

Element/Parameter pair

HBCDD / Concentration of chemical/pollutant in biota

METHOD (Monitoring Details)

Element/parameter

Measured in the following species: Mytius edulis, Platichthys flesus, Clupea harengus, Zoarces viviparous, Gadus morhua, Uria aalge, Sterna hirundo, Haematopus ostralegus, Perca fluvialis.

Sampling and analytical methods are reported per sample and per parameter respectively in the data.

QA/QC

Element/Parameter pair

Quality assurance is a laboratory's whole sampling and analycal process from start to finish.

FREQUENCY

Frequency

Element/Parameter pair

DE:

- UBA: kelb , mussels: Baltic Sea: 2x/year(06/12), annually
- LUNG: mussels, every 2nd or 3rd year
- LLUR: 2x/yr (Mar/Oct) FI: 1-6 years intervals, depending on station; time series stations yearly

EE: in coastal waters 1 or 2 times per WFD 6-years cycle; in EEZ trend stations yearly

SE: Annually at all stations and 2 times per year at two stations

DK: Annually in fish

PL: Yearly at all stations (from 2020: 7 stations - fish; 1 station - mussels)

SPATIAL SCOPE

Spatial Scope

Element/Parameter pair

Coastal Waters /EEZ / Whole Baltic Sea for assessment.

SPATIAL RESOLUTION (DENSITY) OF SAMPLING

Spatial resolution

Element/Parameter pair

HBCDD: are measured in the following areas: Gulf of Gdansk, Southern Baltic Proper, Great Belt, Little Belt, Kattegat, Northern Baltic Proper, Bothnian Bay, Bothnian Sea, Gulf of Riga, Gulf of Finland.

See map for details

PARAMETER

Element/Parameter pair

TBT / Concentration of chemical/pollutant in biota

METHOD (Monitoring Details)

Element/parameter

Method Measured in the following species: Mylus edulis, Zoarces viviparous, Mya arenaria, Plachthys flesus, Clupea harengus, Perca fluvialis and Tritia nitida.

Sampling and analycal methods are reported per sample and per parameter respecvely in the data. See HELCOM COMBINE manual.

QA/QC

Element/Parameter pair

Quality assurance is a laboratory's whole sampling and analycal process from start to finish. See HELCOM COMBINE manual.

FREQUENCY

Frequency

Element/Parameter pair

DE:

UBA: kelb and mussels:Balc Sea: 2x/year (06/12)

• LUNG: mussels: every 2nd or 3rd year

• LLUR: 2x/yr (Mar/Oct)

EE: Rotation three times per 6 years.

SE: Annually at all stations in perch and snail

DK: Annually in fish and mussels

PL: Yearly at all stations (from 2020: 7 stations - fish; 1 station - mussels)

SPATIAL SCOPE

Spatial Scope

Element/Parameter pair

Coastal Waters / transitional waters/ EEZ / Whole Baltic Sea for assessment.

SPATIAL RESOLUTION (DENSITY) OF SAMPLING

Spatial resolution

Element/Parameter pair

TBT is measured in the following areas: The Sound, Kattegat, Great Belt, Southern Baltic Proper, Northern Baltic Proper, Bothnian Sea, Little Belt, Bay of Mecklenburg, Gulf of Gdansk, Gulf of Riga, Gulf of Finland. See map for details

PARAMETER

Element/Parameter pair

Metals / Concentration of chemical/pollutant in biota

METHOD (Monitoring Details)

Element/parameter

Measured in the following species: Platichthys flesus, Mytilus edulis, Zoarces viviparous, Mya arenaria, Perca fluviatilis, Macoma balthica, Clupea harengus, Gadus morhua, Uria aalge, Sterna hirundo, Haematopus ostralegus, Limanda limanda (by DE in Kiel Bay).

Sampling and analytical methods are reported per sample and per parameter respectively in the data. See HELCOM COMBINE manual.

QA/QC

Element/Parameter pair

Quality assurance is a laboratory's whole sampling and analycal process from start to finish. See HELCOM COMBINE manual.

FREQUENCY

Frequency

Element/Parameter pair

DE:

UBA: kelb and mussels:Balc Sea: 2x/year (06/12)

• LUNG: mussels: every 2nd or 3rd year, annually

SE: Annually at all stations and 2 times per year at two stations

DK: Annually in fish (Hg only from2019) and mussels

EE: Annually fish at open sea; 3 times per six years in coastal waters.

PL: Yearly at all stations (from 2020: 7 stations - fish; 1 station - mussels for Hg, Cd, Pb, Cu, Zn, 4 stations - fish for As

SPATIAL SCOPE

Spatial Scope

Element/Parameter pair

Coastal Waters / transitional waters/ EEZ / Whole Baltic Sea for assessment.

SPATIAL RESOLUTION (DENSITY) OF SAMPLING

Spatial resolution

Element/Parameter pair

Metals are measured in the following subbasins: Great Belt, The Sound, Southern Baltic Proper, Kattegat, Little Belt, Kiel Bay, Bay of Mecklenburg, Arkona Basin, Bothnian Bay, Bothnian Sea, Northern Baltic Proper, Gulf of Gdansk.

See map for details

Element/Parameter pair

Radionuclides: Gamma-emitters / Concentration of chemical/pollutant in biota

METHOD (MonitoringDetails)

Element/parameter

Measured in the following species:

- Fish (measured species depends on the location): Clupea harengus, Gadus morhua, Platichthys flesus, Pleuronectes plates, Perca fluviatilis, Esox luciu, Limanda limanda (by DE in Kiel Bay).
- Benthic invertebrates: Mya arenaria, Macoma balthica, Saduria entomon
- Aquatic plants: Fucus vesiculosus

Measured radionuclides:

Obligatory:

• Gamma-emitters: K-40, Cs-137 and other γ -emitters identifies in the γ -spectrum .

Voluntary:

- Fish: Sr-90; natural radionuclides (e.g. Po-210)
- Benthic invertebrates: Sr-90; Tc-99; natural radionuclides (e.g. Po-210); Pu-239, 240; Am-241
- Aquatic plants: Sr-90; Tc-99; Pu-239, 240; Am-241; natural radionuclides

Sampling and analytical methods are reported per sample and per parameter respectively in the MORS database. See MORS Guidelines for detailed descripon.

QA/QC

Element/Parameter pair

Quality assurance is a laboratory's whole sampling and analytical process from start to finish. MORS Guidelines defines methodologies for sample treatment and analysis and intercomparison. Reported data is manually quality assured by HELCOM secretariat and results reported and verified

in annual MORS EG meeting.	
FREQUENCY	
Frequency	
Element/Parameter pair	
Yearly	
SPATIAL SCOPE	
Spatial Scope	
Element/Parameter pair	
EEZ / Whole Baltic Sea for assessment.	
SPATIAL RESOLUTION (DENSITY) OF SAMPLING	
Spatial resolution	
Element/Parameter pair	
See map for details	
Provide considerations for the scale of aggregation of data for an indication of the scale of aggregation of data for an indication of the scale of aggregation of data for an indication of the scale of aggregation of data for an indication of the scale of aggregation of data for an indication of the scale of aggregation of data for an indication of the scale of aggregation of data for an indication of the scale of aggregation of data for an indication of the scale of aggregation of data for an indication of the scale of aggregation of data for an indication of the scale of aggregation of data for an indication of the scale of aggregation of data for an indication of the scale of aggregation of data for an indication of the scale of aggregation of data for an indication of the scale of th	ator-based assessmen
⊠HELCOM assessment unit Level 4: Subbasins with coastal WFD division	
\square HELCOM assessment unit Level 3: Subbasins with coastal and offshore division	
⊠HELCOM assessment unit Level 2: Subbasin	
⊠HELCOM assessment unit Level 1: Baltic Sea	
☐MSFD Region	
□EU	
□Other (specify)	
□Unknown	
c.5 Monitoring and assessment requirements	
Monitoring requirements:	
There is a common agreement concerning statistical requirements for data tindicator assessments.	to be included in the
Adequacy for assessment of GES:	

•	•	information to enable the periodic assessmen as towards GES as required by MSFD under Article 9
	Yes	No
Adequate data?	\boxtimes	
Established methods for assessment?	\boxtimes	
Adequate understanding of GES?		
Adequate capacity to perform assessments?	\boxtimes	
Assessment of natural variability		
Quantitave. The programme ger further used in power analysis .	nerates informati	ion both of within and between - year variation
From which database the data can HELCOM HELCOM PLC COMBINE		
☑Other: ICES DOME and	d MORS, EEA EIO	NET
If the previous answer is "Other" p the HELCOM Secretariat will do it)	lease fill in the nex	xt questions (In case the answer is a HELCOM datab
Data type Tick the relevant box	es below:	
☐Unprocessed/raw Data		
⊠Processed Data sets		
☐ Data Products		
☐Modelled data		
Data management: General descri	ption of data man	nagement (DataManagement, Free text)
No information.		
What method/mechanism will be provide location (DataAccess):	used to make the	data available? Tick the relevant boxes below and
☐ Providing URL to view data: Clic	ck here to enter tex	st.

\square Providing URL to down	load data: Click here to enter text.				
☐ Provide location of data in national data centre: Click here to enter text.					
☐ Provide location of date text.	\square Provide location of data in international data centre (e.g. RSC, ICES, EEA, EMODnet): Click here to enter text.				
When will the data first b	pecome available? (DataPublicationDate)				
Enter the date of reportin	g, or even a past date if desired (MM/YYYY):				
EE: March next year after	er the monitoring.				
How frequently are the d	ata expected to be updated thereafter? Tick the relevant box below:				
□Every 6 years	□Weekly				
☐ Every 3 years	□Daily				
□Every 2 years	□Hourly				
⊠Yearly	□ Continually				
\Box 6-monthly	⊠One-off				
\square 3-monthly	☐ As needed				
\square Monthly	□Other (specify) Click here to enter text.				
\square 2-weekly	□Unknown				
List providing contact noi	nts in the Contracting Portion				
	nts in the Contracting Parties				
EE: Estonian Environme	nt Agency				
EE: Estonian Environme Has the data been used o					
EE: Estonian Environme	nt Agency				
EE: Estonian Environme Has the data been used o	nt Agency				
EE: Estonian Environme Has the data been used o Yes No Select if data is used in the	nt Agency or is it planned to be used in HELCOM assessments? Tick the relevant box below:				
EE: Estonian Environme Has the data been used of Yes	nt Agency or is it planned to be used in HELCOM assessments? Tick the relevant box below:				
EE: Estonian Environme Has the data been used of Yes	nt Agency or is it planned to be used in HELCOM assessments? Tick the relevant box below: ne following Baltic Sea Environment Fact Sheets (BSEF) Tick the relevant boxes ution of marenzelleria species				
EE: Estonian Environme Has the data been used of the last of the	nt Agency or is it planned to be used in HELCOM assessments? Tick the relevant box below: ne following Baltic Sea Environment Fact Sheets (BSEF) Tick the relevant boxes ution of marenzelleria species ution of Round goby				
EE: Estonian Environme Has the data been used of the last of the	nt Agency or is it planned to be used in HELCOM assessments? Tick the relevant box below: ne following Baltic Sea Environment Fact Sheets (BSEF) Tick the relevant boxes ution of marenzelleria species ution of Round goby ution of the Zebra mussel				
EE: Estonian Environme Has the data been used of the last of the	nt Agency or is it planned to be used in HELCOM assessments? Tick the relevant box below: ne following Baltic Sea Environment Fact Sheets (BSEF) Tick the relevant boxes ution of marenzelleria species ution of Round goby ution of the Zebra mussel				
EE: Estonian Environme Has the data been used of the last of the	nt Agency or is it planned to be used in HELCOM assessments? Tick the relevant box below: the following Baltic Sea Environment Fact Sheets (BSEF) Tick the relevant boxes ution of marenzelleria species ution of Round goby ution of the Zebra mussel substant of the Zebra mussel substant of the Baltic Sea				
EE: Estonian Environme Has the data been used of the last of the	nt Agency or is it planned to be used in HELCOM assessments? Tick the relevant box below: the following Baltic Sea Environment Fact Sheets (BSEF) Tick the relevant boxes ution of marenzelleria species ution of Round goby ution of the Zebra mussel the sea and cryptogenic species in the Baltic Sea ant of Great Cormorant				
EE: Estonian Environmee Has the data been used of the last with the last is used in the below: Biodiversity Abundance and distribut the last with last wi	Int Agency In is it planned to be used in HELCOM assessments? Tick the relevant box below: Interpolation of Baltic Sea Environment Fact Sheets (BSEF) Tick the relevant boxes Interpolation of Round goby Interpolation of the Zebra mussel Interpolation of Great Cormorant Interpolation of Sandwich Tern				

☐ Temporal development of Baltic coastal fish communities and key species
Eutrophication
☐ Bacterioplankton growth
□Chlorophyll-a concentrations, temporal variations and regional differences from satellite remote sensing
☐ Cyanobacteria biomass
☐ Cyanobacterial blooms in the Baltic Sea
☐Cyanobacteria bloom index
\square Impacts of invasive phytoplankton species on the Baltic Sea ecosystem in 1980-2008
\square Nitrogen atmospheric deposition to the Baltic Sea
\square Nitrogen emissions to the air in the Baltic Sea area
☐ Phytoplankton biomass and species succession
\square Shifts in the Baltic Sea summer phytoplankton communities in 1992-2006
\square Spatial distribution of the winter nutrient pool
☐Unusual phytoplankton event
Hazardous substances
☐ Atmospheric deposition of heavy metals on the Baltic Sea
☐ Atmospheric deposition of PCDD/Fs on the Baltic Sea
☐ Atmospheric emissions of heavy metals in the Baltic Sea region
☐ Atmospheric emissions of PCDD/Fs in the Baltic Sea region
☐ Cesium-137 in Baltic Sea sediments
☑Temporal trends in contaminants in Herring in the Baltic Sea in the period 1980-2010
☐ Emissions from Baltic Sea shipping
□ Illegal discharges of oil in the Baltic Sea
☐ Liquid discharges of Cs-137, Sr-90 and Co-60 into the Baltic Sea
☐ Trace metal concentrations and trends in Baltic surface and deep waters
Hydrography
☐ Development of Sea Surface Temperature in the Baltic Sea
☐ Hydrography and Oxygen in the Deep Basins
□lce season
□Total and regional runoff to the Baltic Sea
\square Water Exchange between the Baltic Sea and the North Sea, and conditions in the Deep Basins

Γ	71/21/0	climate	in tha	Daltic	500
- 1	IVVAVE	cumare	in the	ванис	763

c.7 MSFD Criteria (GES Criteria)Choose only the most relevant option(s). Tick one or more boxes below.

Descriptor 1	□ D1C1 – Primary:
	The mortality rate per species from incidental by-catch is below levels which threaten the species, such that its long- term viability is ensured.
	Member States shall establish the threshold values for the mortality rate from incidental by-catch per species, through regional or subregional cooperation.
	□ D1C2 – Primary:
	The population abundance of the species is not adversely affected due to anthropogenic pressures, such that its long-term viability is ensured.
	Member States shall establish threshold values for each species through regional or subregional cooperation, taking account of natural variation in population size and the mortality rates derived from D1C1, D8C4 and D10C4 and other relevant pressures. For species covered by Directive 92/43/EEC, these values shall be consistent with the Favourable Reference Population values established by the relevant Member States under Directive 92/43/EEC.
	$\ \square$ D1C3 – Primary for commercially- exploited fish and cephalopods and secondary for other species:
	The population demographic characteristics (e.g. body size or age class structure, sex ratio, fecundity, and survival rates) of the species are indicative of a healthy population which is not adversely affected due to anthropogenic pressures.
	Member States shall establish threshold values for specified characteristics of each species through regional or subregional cooperation, taking account of adverse effects on their health derived from D8C2, D8C4 and other relevant pressures.
	\square D1C4 – Primary for species covered by Annexes II, IV or V to Directive 92/43/EEC and secondary for other species:
	The species distributional range and, where relevant, pattern is in line with prevailing physiographic, geographic and climatic conditions.
	Member States shall establish threshold values for each species through regional or subregional cooperation. For species covered by Directive 92/43/EEC, these shall be consistent with the Favourable Reference Range values established by the relevant Member States under Directive 92/43/EEC.
	\square D1C5 – Primary for species covered by Annexes II, IV and V to Directive 92/43/EEC and secondary for other species:
	The habitat for the species has the necessary extent and condition to support the different stages in the life history of the species.
	□ D1C6 – Primary
	The condition of the habitat type, including its biotic and abiotic structure and its functions (e.g. its typical species composition and their relative abundance, absence of particularly sensitive or fragile species or species providing a key function, size

	structure of species), is not adversely affected due to anthropogenic pressures.
	Member States shall establish threshold values for the condition of each habitat type, ensuring compatibility with related values set under Descriptors 2, 5 and 8, through regional or subregional cooperation.
Descriptor 2	□ D2C1 – Primary:
	The number of non-indigenous species which are newly introduced via human activity into the wild, per assessment period (6 years), measured from the reference year as reported for the initial assessment under Article 8(1) of Directive 2008/56/EC, is minimised and where possible reduced to zero.
	Member States shall establish the threshold value for the number of new introductions of non-indigenous species, through regional or subregional cooperation.
	□ D2C2 — Secondary:
	Abundance and spatial distribution of established non-indigenous species, particularly of invasive species, contributing significantly to adverse effects on particular species groups or broad habitat types.
	□ D2C3 — Secondary:
	Proportion of the species group or spatial extent of the broad habitat type which is adversely altered due to non-indigenous species, particularly invasive non-indigenous species.
	Member States shall establish the threshold values for the adverse alteration to species groups and broad habitat types due to non-indigenous species, through regional or subregional cooperation.
Descriptor 3	□ D3C1 — Primary:
	The Fishing mortality rate of populations of commercially-exploited species is at or below levels which can produce the maximum sustainable yield (MSY). Appropriate scientific bodies shall be consulted in accordance with Article 26 of Regulation (EU) No 1380/2013.
	□ D3C2 — Primary:
	The Spawning Stock Biomass of populations of commercially-exploited species are above biomass levels capable of producing maximum sustainable yield. Appropriate scientific bodies shall be consulted in accordance with Article 26 of Regulation (EU) No 1380/2013.
	□ D3C3 — Primary:
	The age and size distribution of individuals in the populations of commercially-exploited species is indicative of a healthy population. This shall include a high proportion of old/large individuals and limited adverse effects of exploitation on genetic diversity.
	Member States shall establish threshold values through regional or subregional cooperation for each population of species in accordance with scientific advice obtained pursuant to Article 26 of Regulation (EU) No 1380/2013.
Descriptor 4	□ D4C1 — Primary:

	The diversity (species composition and their relative abundance) of the trophic guild is not adversely affected due to anthropogenic pressures.
	Member States shall establish threshold values through regional or subregional cooperation.
	□ D4C2 — Primary:
	The balance of total abundance between the trophic guilds is not adversely affected due to anthropogenic pressures.
	Member States shall establish threshold values through regional or subregional cooperation.
	☐ D4C3 — Secondary:
	The size distribution of individuals across the trophic guild is not adversely affected due to anthropogenic pressures.
	Member States shall establish threshold values through regional or subregional cooperation.
	\square D4C3 — Secondary (to be used in support of criterion D4C2, where necessary):
	Productivity of the trophic guild is not adversely affected due to anthropogenic pressures.
	Member States shall establish threshold values through regional or subregional cooperation.
Descriptor 5	□ D5C1 — Primary:
	Nutrient concentrations are not at levels that indicate adverse eutrophication effects.
	The threshold values are as follows:
	(a) in coastal waters, the values set in accordance with Directive 2000/60/EC;
	(b) beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation
	□ D5C2 — Primary:
	Chlorophyll a concentrations are not at levels that indicate adverse effects of nutrient enrichment.
	The threshold values are as follows:
	(c) in coastal waters, the values set in accordance with Directive 2000/60/EC;
	(d) beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation.
	□ D5C3 — Secondary:
	The number, spatial extent and duration of harmful algal bloom events are not at levels that indicate adverse effects of nutrient enrichment.
	□ D5C4 — Secondary:
	The photic limit (transparency) of the water column is not reduced, due to increases in

suspended algae, to a level that indicates adverse effects of nutrient enrichment.
The threshold values are as follows:
(e) in coastal waters, the values set in accordance with Directive 2000/60/EC;
(f) beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation.
\square D5C5 — Primary (may be substituted by D5C8):
The concentration of dissolved oxygen is not reduced, due to nutrient enrichment, to levels that indicate adverse effects on benthic habitats (including on associated biota and mobile species) or other eutrophication effects.
The threshold values are as follows:
(g) in coastal waters, the values set in accordance with Directive 2000/60/EC;
(h) beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation.
☐ D5C6 — Secondary:
The abundance of opportunistic macroalgae is not at levels that indicate adverse effects of nutrient enrichment.
The threshold values are as follows:
(a) in coastal waters, the values set in accordance with Directive 2000/60/EC;
(b) should this criterion be relevant for waters beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation.
☐ D5C7 — Secondary:
The species composition and relative abundance or depth distribution of macrophyte communities achieve values that indicate there is no adverse effect due to nutrient enrichment including via a decrease in water transparency, as follows:
(a) in coastal waters, the values set in accordance with Directive 2000/60/EC;
(b) should this criterion be relevant for waters beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation.
\square D5C8 — Secondary: (except when used as a substitute for D5C5):
The species composition and relative abundance of macrofaunal communities, achieve values that indicate that there is no adverse effect due to nutrient and organic enrichment, as follows:
(a) in coastal waters, the values for benthic biological quality elements set in accordance with Directive 2000/60/EC;
(b) beyond coastal waters, values consistent with those for coastal waters under

	Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation.
Descriptor 6	□ D6C1 – Primary:
	Spatial extent and distribution of physical loss (permanent change) of the natural seabed.
	□ D6C2 – Primary:
	Spatial extent and distribution of physical disturbance pressures on the seabed.
	□ D6C3 – Primary:
	Spatial extent of each habitat type which is adversely affected, through change in its biotic and abiotic structure and its functions (e.g. through changes in species composition and their relative abundance, absence of particularly sensitive or fragile species or species providing a key function, size structure of species), by physical disturbance.
	Member States shall establish threshold values for the adverse effects of physical disturbance, through regional or subregional cooperation.
	☐ D6C4 – Primary:
	The extent of loss of the habitat type, resulting from anthropogenic pressures, does not exceed a specified proportion of the natural extent of the habitat type in the assessment area.
	Member States shall establish the maximum allowable extent of habitat loss as a proportion of the total natural extent of the habitat type, through cooperation at Union level, taking into account regional or subregional specificities.
	□ D6C5 – Primary:
	The extent of adverse effects from anthropogenic pressures on the condition of the habitat type, including alteration to its biotic and abiotic structure and its functions (e.g. its typical species composition and their relative abundance, absence of particularly sensitive or fragile species or species providing a key function, size structure of species), does not exceed a specified proportion of the natural extent of the habitat type in the assessment area.
	Member States shall establish threshold values for adverse effects on the condition of each habitat type, ensuring compatibility with related values set under Descriptors 2, 5, 6, 7 and 8, through cooperation at Union level, taking into account regional or subregional specificities. Member States shall establish the maximum allowable extent of those adverse effects as a proportion of the total natural extent of the habitat type, through cooperation at Union level, taking into account regional or subregional specificities.
Descriptor 7	□ D7C1 – Secondary:
	Spatial extent and distribution of permanent alteration of hydrographical conditions (e.g. changes in wave action, currents, salinity, temperature) to the seabed and water column, associated in particular with physical loss(1) of the natural seabed.
	□ D7C2 – Secondary: Snatial extent of each benthic babitat type adversely affected (physical and
	Spatial extent of each benthic habitat type adversely affected (physical and

	hydrographical characteristics and associated biological communities) due to permanent alteration of hydrographical conditions.
Descriptor 8	☑ D8C1 – Primary:
	Within coastal and territorial waters, the concentrations of contaminants do not exceed the following threshold values:
	(a) for contaminants set out under point 1(a) of criteria elements, the values set in accordance with Directive 2000/60/EC;
	(b) when contaminants under point (a) are measured in a matrix for which no value is set under Directive 2000/60/EC, the concentration of those contaminants in that matrix established by Member States through regional or subregional cooperation;
	(c) for additional contaminants selected under point 1(b) of criteria elements, the concentrations for a specified matrix (water, sediment or biota) which may give rise to pollution effects. Member States shall establish these concentrations through regional or subregional cooperation, considering their application within and beyond coastal and territorial waters.
	Beyond territorial waters, the concentrations of contaminants do not exceed the following threshold values:
	(a) for contaminants selected under point 2(a) of criteria elements, the values as applicable within coastal and territorial waters;
	(b) for contaminants selected under point 2(b) of criteria elements, the concentrations for a specified matrix (water, sediment or biota) which may give rise to pollution effects. Member States shall establish these concentrations through regional or subregional cooperation.
	□ D8C2 – Secondary:
	The health of species and the condition of habitats (such as their species composition and relative abundance at locations of chronic pollution) are not adversely affected due to contaminants including cumulative and synergetic effects.
	Member States shall establish those adverse effects and their threshold values through regional or subregional cooperation.
	□ D8C3 – Primary:
	The spatial extent and duration of significant acute pollution events are minimised.
	\square D8C4 – Secondary (to be used when a significant acute pollution event has occurred):
	The adverse effects of significant acute pollution events on the health of species and on the condition of habitats (such as their species composition and relative abundance) are minimised and, where possible, eliminated.
Descriptor 9	☑ D9C1 – Primary:
	The level of contaminants in edible tissues (muscle, liver, roe, flesh or other soft parts, as appropriate) of seafood (including fish, crustaceans, molluscs, echinoderms, seaweed and other marine plants) caught or harvested in the wild (excluding fin-fish from mariculture) does not exceed:

	(a) for contaminants listed in Regulation (EC) No 1881/2006, the maximum levels laid down in that Regulation, which are the threshold values for the purposes of this Decision;(b) for additional contaminants, not listed in Regulation (EC) No 1881/2006,
	threshold values, which Member States shall establish through regional or subregional cooperation.
Descriptor 10	□ D10C1 – Primary:
	The composition, amount and spatial distribution of litter on the coastline, in the surface layer of the water column, and on the seabed, are at levels that do not cause harm to the coastal and marine environment.
	Member States shall establish threshold values for these levels through cooperation at Union level, taking into account regional or subregional specificities.
	□ D10C2 — Primary:
	The composition, amount and spatial distribution of micro-litter on the coastline, in the surface layer of the water column, and in seabed sediment, are at levels that do not cause harm to the coastal and marine environment.
	Member States shall establish threshold values for these levels through cooperation at Union level, taking into account regional or subregional specificities.
	□ D10C3 — Secondary:
	The amount of litter and micro-litter ingested by marine animals is at a level that does not adversely affect the health of the species concerned. Member States shall establish threshold values for these levels through regional or subregional cooperation.
	□ D10C4 — Secondary:
	The number of individuals of each species which are adversely affected due to litter, such as by entanglement, other types of injury or mortality, or health effects. Member States shall establish threshold values for the adverse effects of litter, through regional or subregional cooperation.
Descriptor 11	□ D11C1 – Primary:
	The spatial distribution, temporal extent, and levels of anthropogenic impulsive sound sources do not exceed levels that adversely affect populations of marine animals.
	Member States shall establish threshold values for these levels through cooperation at Union level, taking into account regional or subregional specificities.
	□ D11C2 – Primary:
	The spatial distribution, temporal extent and levels of anthropogenic continuous low-frequency sound do not exceed levels that adversely affect populations of marine animals.
	Member States shall establish threshold values for these levels through cooperation at Union level, taking into account regional or subregional specificities.

d. References

Make a list of cited references and literature for further supportive information.

HELCOM, 2019. Monitoring guideline on biological material sampling and sample handling for the analysis of persistent organic pollutants (PAHs, PCBs and OCPs) and metallic trace elements, https://helcom.fi/wp-content/uploads/2020/04/HELCOM-Guidelines-on-biological-material-sampling-1.pdf. HELCOM, 2019. Guidelines for monitoring of radioactive substances to be followed when implementing HELCOM Recommendation 26/3 (https://helcom.fi/media/publications/Guidelines-for-Monitoring-of-Radioactive-Substances.pdf)ological-material-sampling-1.pdf.