HELCOM Monitoring Programme topic

Concentrations of contaminants

Programme:

Contaminants in sediment

Contents

a.	Metadata on monitoring strategies and monitoring programmes	2
	a.1 Responsible HELCOM subsidiary body	2
	a.2 Regional Cooperation (RegionalCooperation)	3
b.	Monitoring strategies	4
	b.1 Descriptor	4
	b.2 BSAP segments	4
	b.3 Monitoring strategy description	4
	b.4 BSAP Ecological objectives	4
	b.5 Gaps in monitoring	5
c.	Monitoring programmes	6
	c.1 Purpose of monitoring	6
	c.2 Other legislation	10
	c.3 Implementation of Regional Cooperation (RegionalCooperation_implementation)	10
	c.4 Monitoring concepts	11
	c.5 Monitoring and assessment requirements	24
	c.6 Data providers and access	25
	c.7 MSFD Criteria (GES criteria)	27
٦	Poforoncos	21

a. Metadata on monitoring strategies and monitoring programmes

a.1 Responsible HELCOM subsidiary body

Please indicate the relevant expert group/network if available, otherwise the responsible HELCOM Working Group. or the responsible working group understanding is that they are the EN or EG if available, otherwise the WGs

	Permament Groups
	Gear – Group on the Implementation of the Ecosystem Approach
	Maritime – Maritime Working Group
	Pressure – Working Group on Reduction of Pressures from the Baltic Sea Catchment Area
	Response – Response Working Group
\boxtimes	State and Conservation – Working Group on the State of the Environmental and Nature Conservation
	Time-limited Groups
	Agri – Group on Sustainable Agricultural Practices
	Fish – Group on Ecosystem-based Sustainable Fisheries
	HELCOM-VASAB MSP WG - Joint HELCOM-VASAB Maritime Spatial Planning Working Group
	Expert Groups
	AIS EWG – Expert Working Group for Mutual Exchange and Deliveries of AIS data
\boxtimes	EN Hazardous Substances – Expert Network on hazardous substances
	EN Marine Litter – Expert Network on Marine Litter
	EN Noise – Expert Network on Underwater Noise
	ESA – Expert Network on Economic and Social Analyses
	EWG OWR – Expert Working Group on Oiled Wildlife Response
	EWG SHORE – Expert Working Group on Response on the Shore
	Green Technology and Alternative Fuels Platform for Shipping
	HELCOM/OSPAR TG BALLAST – Joint HELCOM/OSPAR Task Group on Ballast

	Management Convention Exemptions
	IN Benthic habitat – Intersessional Network on habitat monitoring
	IWGAS – Informal Working Group on Aerial Surveillance
	JWG Bird – HELCOM-OSPAR-ICES Joint Working Group on Seabirds
\boxtimes	MORS EG – Expert group on monitoring of radioactive substances in the Baltic Sea
	PRF Cooperation Platform – Cooperation Platform on Port Reception Facilities in the Baltic Sea
	SAFE NAV – Group of Experts on Safety of Navigation
	SUBMERGED – Expert Group on Environmental Risks of Hazardous Submerged Objects

a.2 Regional Cooperation (RegionalCooperation)

THE HIGHHOLDS OF THIS DIOSIALITIE IS.	The	monitoring	of this	programme	is:
---------------------------------------	-----	------------	---------	-----------	-----

\square Fully coordinated	
□ Partly coordinated.	Indicate missing component(s

 \Box Coordinated monitoring is under development. Indicate by which group/project and by when a recommendation on coordinated monitoring can be expected.

- Common monitoring guidelines in COMBINE manual, ISO/IEC 17025, Guidelines for determination of chlorinated hydrocarbons in sediment, Guidelines for determination of PAH in sediment and other guidelines (see References).
- Common quality assurance programme: HELCOM COMBINE manual, ISO/CEN standards and QUASIMEME. Radioactive substances: MORS Guidelines defines methodologies for sample treatment, analysis and intercomparison. Reported data is manually quality assured by the HELCOM Secretariat and results are reported and verified in annual MORS EG meetings.
- Common databases: COMBINE (ICES DOME), MORS.

There is no current plan for coordinated monitoring of contaminants in sediments, other than radionuclides. Under the Water Framework Directive, no EQS values have so far been set in sediments. However, countries are encouraged to perform long term trend analysis of concentrations of persistent substances that tend to accumulate in sediment, in order to ensure that such concentrations do not significantly increase (such as Hg, PAHs, HCHs, HCB, PBDE, TBT, PCB, PFOS, PCDD/F, HBCDD).

b. Monitoring strategies

b.1 Descriptor

supports the following obligatory MSFD Monitoring Strategies. Tick one or more relevant
Biodiversity
Non-indigenous Species
Commercial fish and shellfish
Food webs
Eutrophication
Seafloor integrity
Hydrographical conditions
Contaminants
Contaminants in seafood
Marine litter
Energy including underwater noise
segments nme serves the following BSAP segments. Tick one or more relevant boxes.
า
ostances
rities
oring strategy description
egy: Monitoring is to be carried out to fulfill assessment requirements of HELCOM cives that are specified through HELCOM core indicators. The requirements on include number of stations, the sampling frequency and replication.
Ecological objectives most relevant option(s). Tick one or more boxes below.
☐ Concentrations of nutrients close to natural levels

	☐ Clear water
	☐ Natural level of algal blooms
	$\hfill\square$ Natural distribution and occurrence of plants and animals
	☐ Natural oxygen levels
Hazardous	☑ Concentrations of hazardous substances close to natural levels
substances	☐ All fish safe to eat
	□ Healthy wildlife
	☑ Radioactivity at pre-Chernobyl levels
Biodiversity	☐ Natural landscapes and seascapes
	\square Thriving and balanced communities of plants and animals
	☐ Viable populations of species
Maritime	☐ No illegal pollution
activities	\square Safe maritime traffic without accidental pollution
	☐ Efficient response capability
	\square No introductions of alien species from ships
	☐ Minimum air pollution from ships
	☐ Zero discharges from offshore platforms
In relation to the	n monitoring GES criteria addressed, indicate when sufficient monitoring was in place or by when ge will be in place (Coverage_GEScriteria)
☐ Adequate mo	nitoring was in place in 2014
☐ Adequate mo	nitoring was in place by 2018
☐ Adequate mo	nitoring is in place by July 2020
☐ Adequate mo	nitoring will be in place by 2024
☐ Monitoring is	not being put in place for this descriptor due to a low risk
\square Monitoring fo	r this descriptor is not relevant
•	e implementation gaps and plans to complete the establishment and implementation of conitoring strategy (Gaps_Plans):
Assessment of	gaps has not been carried out.
	·

c. Monitoring programmes

c.1 Purpose of monitoring

c.1a Assessment purpose in general

The programme supports the assessment of:

_					
T	irl	tha	ro	levant	hov
- 11	-	uie	10	ıevaiii	DUX.

Temporal trends	Spatial distribution	State classification
	\boxtimes	

The **programme** supports the assessment of: (MonitoringPurpose).

Note that the answer to this question will be decisive for whether to answer upcoming questions e.g. upcoming questions on pressures should only be answered if the monitoring is defined as supporting the assessment of pressures.

Tick the relevant boxes.

Environmental state and impacts	Pressures in the marine environment	Pressures at source (land-based, riverine, sea-based ¹ and atmospheric sources)	Human activities causing the pressures	Effectiveness of measures
If this is selected fill in the following questions: c.1b	If this is selected fill in the following questions: c.1c, d	If this is selected fill in the following questions: c.1c, d	If this is selected fill in the following questions: c.1c, d	If this is selected fill in the following questions: c.1c, d

Give any other monitoring purpose e.g. if the programmes include supporting parameters for other monitoring programmes

Gives input into marine ecosystems health assessments (bottom species), also for secondary pollution risk assessments. Gives input into marine ecosystems health assessments (bottom species), also for secondary pollution risk assessments.

For questions 1b-1d, select when applicable for the sub-programme, the link from the Reporting on the 2020 update of Article 11 for the Marine Strategy Framework Directive (MSFD Guidance Document 17, 2020) (Features) to:

- Ecosystem components (relevant for monitoring and assessment for Article 8(1a) for D1C2-C5, D3, D4, D6C3-C5, D7C2)
- Pressures and impacts in the marine environment (relevant for monitoring and assessment for Article 8(1b) for D1C1, D2, D5, D6C1-C2, D7C1, D8, D9, D10, D11)
- Pressure inputs to the marine environment (relevant for monitoring and assessment for Article 10)
- Uses and human activities (relevant for monitoring and assessment for Article 8(1c) and 13)

¹ Sea-based 'Pressures at source' refers to monitoring pressures from sea-based activities where the monitoring is directly at the activity rather than at a distance from or time period after it is generated by the activity (e.g. D1 incidental by-catch when fishing, D2 ballast water discharges, D6 use of bottom fishing gear, D8 contaminant discharges and pollution events from a vessel or pipeline, D11 impulsive sound events from a vessel or platform).

c.1b • Ecosystem components (Features)Choose only the most relevant option(s). Tick one or more boxes below.

Theme	Sub-theme	Label feature
Species	☐ Birds	☐ Grazing birds
		☐ Wading birds
		☐ Surface-feeding birds
		☐ Pelagic-feeding birds
		☐ Benthic-feeding birds
	☐ Mammals	\square Small toothed cetaceans
		\square Deep-diving toothed cetaceans
		☐ Baleen whales
		☐ Seals
	☐ Reptiles	☐ Turtles
	☐ Fish	☐ Coastal fish
		☐ Pelagic shelf fish
		☐ Demersal shelf fish
		☐ Deep-sea fish
		☐ Commercially exploited fish and shellfish
	☐ Cephalopods	☐ Coastal/shelf cephalopods
		☐ Deep-sea cephalopods
Habitats	\square Benthic habitats	⊠ Benthic broad habitats
		○ Other benthic habitats
	☐ Pelagic habitats	☐ Pelagic broad habitats
		☐ Other pelagic habitats
Ecosystems	☐ Physical and hydrological c	characteristics
	□ Chemical characteristics	
	☐ Ecosystems, including	☐ Coastal ecosystems
	food webs	☐ Shelf ecosystems
		☐ Oceanic/deep-sea ecosystems
	• Pressures and impacts in the most relevant option(s). Tick	he marine environment (Features) k one or more boxes below.
Theme	Label: Feature	

Biological	☐ Newly introduced non-indigenous species
	☐ Established non-indigenous species
	☐ Species affected by incidental by-catch
Physical and	☐ Hydrographical changes
hydrological	☐ Physical disturbance to seabed
	☐ Physical loss of the seabed
Substances,	☐ Eutrophication
litter and energy	□ Contaminants - non UPBT substances
energy	□ Contaminants - UPBT substances
	☐ Contaminants – in seafood
	☐ Acute pollution events
	☐ Litter in the environment
	☐ Impulsive sound in water
	☐ Continuous low frequency sound
c.1d •	Pressure inputs to the marine environment (Features)
c.1d •	Pressure inputs to the marine environment (Features)
c.1d •	Pressure inputs to the marine environment (Features) Label: Feature
Theme	Label: Feature
Theme	Label: Feature ☐ Input or spread of non-indigenous species
Theme	Label: Feature ☐ Input or spread of non-indigenous species ☐ Input of microbial pathogens
Theme	Label: Feature ☐ Input or spread of non-indigenous species ☐ Input of microbial pathogens ☐ Input of genetically modified species and translocation of native species ☐ Loss of, or change to, natural biological communities due to cultivation of
Theme	Label: Feature Input or spread of non-indigenous species Input of microbial pathogens Input of genetically modified species and translocation of native species Loss of, or change to, natural biological communities due to cultivation of animal or plant species Disturbance of species (e.g. where they breed, rest and feed) due to human
Theme	Label: Feature ☐ Input or spread of non-indigenous species ☐ Input of microbial pathogens ☐ Input of genetically modified species and translocation of native species ☐ Loss of, or change to, natural biological communities due to cultivation of animal or plant species ☐ Disturbance of species (e.g. where they breed, rest and feed) due to human presence ☐ Extraction of, or mortality/injury to, wild species (by commercial and
Theme Biological Substances, litter and	Label: Feature ☐ Input or spread of non-indigenous species ☐ Input of microbial pathogens ☐ Input of genetically modified species and translocation of native species ☐ Loss of, or change to, natural biological communities due to cultivation of animal or plant species ☐ Disturbance of species (e.g. where they breed, rest and feed) due to human presence ☐ Extraction of, or mortality/injury to, wild species (by commercial and recreational fishing and other activities)
Theme Biological Substances,	Label: Feature ☐ Input or spread of non-indigenous species ☐ Input of microbial pathogens ☐ Input of genetically modified species and translocation of native species ☐ Loss of, or change to, natural biological communities due to cultivation of animal or plant species ☐ Disturbance of species (e.g. where they breed, rest and feed) due to human presence ☐ Extraction of, or mortality/injury to, wild species (by commercial and recreational fishing and other activities) ☐ Input of nutrients — diffuse sources, point sources, atmospheric deposition
Theme Biological Substances, litter and	Label: Feature ☐ Input or spread of non-indigenous species ☐ Input of microbial pathogens ☐ Input of genetically modified species and translocation of native species ☐ Loss of, or change to, natural biological communities due to cultivation of animal or plant species ☐ Disturbance of species (e.g. where they breed, rest and feed) due to human presence ☐ Extraction of, or mortality/injury to, wild species (by commercial and recreational fishing and other activities) ☐ Input of nutrients — diffuse sources, point sources, atmospheric deposition ☐ Input of other substances (e.g. synthetic substances, non-synthetic substances, radionuclides) — diffuse sources, point sources, atmospheric deposition, acute
Theme Biological Substances, litter and	Label: Feature ☐ Input or spread of non-indigenous species ☐ Input of microbial pathogens ☐ Input of genetically modified species and translocation of native species ☐ Loss of, or change to, natural biological communities due to cultivation of animal or plant species ☐ Disturbance of species (e.g. where they breed, rest and feed) due to human presence ☐ Extraction of, or mortality/injury to, wild species (by commercial and recreational fishing and other activities) ☐ Input of nutrients — diffuse sources, point sources, atmospheric deposition ☐ Input of other substances (e.g. synthetic substances, non-synthetic substances, radionuclides) — diffuse sources, point sources, atmospheric deposition, acute events
Theme Biological Substances, litter and	Label: Feature ☐ Input or spread of non-indigenous species ☐ Input of microbial pathogens ☐ Input of genetically modified species and translocation of native species ☐ Loss of, or change to, natural biological communities due to cultivation of animal or plant species ☐ Disturbance of species (e.g. where they breed, rest and feed) due to human presence ☐ Extraction of, or mortality/injury to, wild species (by commercial and recreational fishing and other activities) ☐ Input of nutrients — diffuse sources, point sources, atmospheric deposition ☐ Input of organic matter — diffuse sources and point sources ☑ Input of other substances (e.g. synthetic substances, non-synthetic substances, radionuclides) — diffuse sources, point sources, atmospheric deposition, acute events ☐ Input of litter (solid waste matter, including micro-sized litter)

heat)
☐ Input of water — point sources (e.g. brine)

c.1e • Uses and human activities (Features)

Choose only the most relevant option(s). Tick one or more boxes below.

Theme	Label: Feature						
Physical	☐ Land claim						
restructuring of rivers, coastline	☐ Canalisation and other watercourse modifications						
or seabed (water	☐ Coastal defence and flood protection						
management)	☐ Offshore structures (other than for oil/gas/renewables)						
	☐ Restructuring of seabed morphology, including dredging and depositing of materials						
Extraction of	☐ Extraction of minerals (rock, metal ores, gravel, sand, shell)						
non-living resources	☐ Extraction of oil and gas, including infrastructure						
	☐ Extraction of salt						
	☐ Extraction of water						
Production of energy	$\hfill\square$ Renewable energy generation (wind, wave and tidal power), including infrastructure						
	☐ Non-renewable energy generation						
	☐ Transmission of electricity and communications (cables)						
Extraction of	☐ Fish and shellfish harvesting (professional, recreational)						
living resources	☐ Fish and shellfish processing						
	☐ Marine plant harvesting						
	☐ Hunting and collecting for other purposes						
Cultivation of	☐ Aquaculture — marine, including infrastructure						
living resources	☐ Aquaculture — freshwater						
	☐ Agriculture						
	□ Forestry						
Transport	☐ Transport infrastructure						
	☐ Transport — shipping						
	□ Transport — air						
	☐ Transport — land						
Urban and	☐ Urban uses						
industrial uses	□ Industrial uses						

	☐ Waste treatment and disposal
Tourism and	☐ Tourism and leisure infrastructure
leisure	☐ Tourism and leisure activities
Security/defence	☐ Military operations (subject to Article 2(2))
Education and research	☐ Research, survey and educational activities
c.2 Other le	ne links with the following other international legislation (OtherPoliciesConventions). Tick
☐Bathing Water [Directive
\square Common Fisher	ies Policy and Data Collection Framework
☐ Habitats Directi	ve
\square Birds Directive	
□ Nitrates Directiv	ve
□Urban Waste W	ater Treatment Directive
⊠Water Framewo	ork Directive
⊠OSPAR Convent	ion
☐Trilateral Wadd	en Sea Convention
⊠Other, Specify:	Maritime Spatial Planning Directive
(RegionalC	entation of Regional Cooperation ooperation_implementation) of implementation by selecting one of the following:
☐Agreed data col	lection methods
☐Common monit	oring strategy (spatial and temporal design of programme)
⊠Coordinated da	ta collection (delivered separately by each country)
☐Joint data collec	ction (multinational delivery using same platform and/or algorithms)

c.4 Monitoring concepts

Monitoring concepts table²:

Current means of coordination	Features or Elements	Parameter	Method	QA/QC	Freque ncy ³	Spatial resolution (density) of sampling	Link to HELCOM core indicators ⁴	Spatial scope	(year)	CPs monitoring ⁵
	Elements (Features) (Features_enu m)	Parameters (Parameter) (ParametersOther)	MonitoringMethod (Monitoring Method) MonitoringMethod Other)	(Free text)	Monitori ngFreque ncy	(ProgrammeDesc ription)	(RelatedIndicator) (RelatedIndicator_n ame	(SpatialScope)	(TemporalScope)	(CountryCode_E num)
Regional (Combine)	PCBs	Concentration of chemical/nutri ent/pollutant in/on seabed substrate	Guidelines for determination of chlorinated hydrocarbons in sediment	HELCOM COMBINE manual, ISO/CEN standards and QUASIMEME	Other	See map for details		EEZ/ Transitional waters	DE: 2000, 2008 DK: 1999 with some interupons in the mid 2000's EE: 2010 (as research, monitoring 2014) PL: 1998 SE: 2003 (now repeated every 6 years)	DE, DK, PL, SE, EE

-

² Needed codelists can be found on 2020 update of Article 11 for the Marine Strategy Framework Directive (MSFD Guidance Document 17, 2020).

³ The option "Different for each country - see MORE overview" refers to the <u>overview</u> carried out in 2013

 $^{^{\}rm 4}$ Give the name of HELCOM core indicators that are based on the monitoring parameter.

⁵ Provide information on the Contracting Partie(s) that are monitoring the parameter.

Current means of coordination	Features or Elements	Parameter	Method	QA/QC	Freque ncy ³	Spatial resolution (density) of sampling	Link to HELCOM core indicators ⁴	Spatial scope		CPs monitoring ⁵
Regional (Combine)	РАН	Concentration of chemical/nutri ent/pollutant in/on seabed substrate	Guidelines for determination of PAH in sediment	HELCOM COMBINE manual, Annex B13, Appendix 1 and 2. ISO/CEN standards and QUASIMEME	Other, LT: 1 time per year (summ er)	See map for details		EEZ, Teriorial waters, Transional waters/coast al waters	EE: 2014 (research 2010) LT: 2007 PL:2012 SE: 2003 (repeated every 6 years)	DE, DK until 2017, LT, PL, SE, EE
Regional (Combine)	Tributyltin (TBT)	Concentration of chemical/nutri ent/pollutant in/on seabed substrate	HELCOM COMBINE Manual, Part D. ISO/IEC 17025 and other guidelines (see References)	HELCOM COMBINE manual, ISO/CEN standards and QUASIMEME	Other	See map for details		EEZ/ Transitional waters/coast al waters	DE: 2000, 2004, 2008 DK: 1999 with a few gaps EE:2014 LT: 2011 LV: 2011 PL: 2018 SE: 2003 (repeated every 6 years)	DE, DK until 2017, FI, LT, LV, SE, PL, EE

Current means of coordination	Features or Elements	Parameter	Method	QA/QC	Freque ncy ³	Spatial resolution (density) of sampling	Link to HELCOM core indicators ⁴	Spatial scope	_	CPs monitoring ⁵
Regional (Combine)	Metals	Concentration of chemical/nutri ent/pollutant in/on seabed substrate	Guideline for the determination of heavy metals in sediment	Guideline for the determination of heavy metals in sediment	Other	See map for details		EEZ/ Transitional waters/ Coastal waters	DE: 2000 DK: 1999 EE: 2010 (research; monitoring 2014) LT: 2007, data also from 2004 LV: 2007, data also from 2004 PL: Cd, Pb,Cu, Zn - 1998; Hg – 2002; Al – 2012; As – 2020) SE: 2003 (repeated every 6 years)	DE, DK until 2017, LT, LV, SE, PL, EE
National	Furans (SE: and dioxins)	Concentration of chemical/nutri ent/pollutant in/on seabed substrate	Different approaches e.g. CEMP manual, ICES guidelines, ISO/CEN standards (see References)	Other	Other			EEZ	DK: 2007 SE: 2014	DK until 2017, SE

Current means of coordination	Features or Elements	Parameter	Method	QA/QC	Freque ncy ³	Spatial resolution (density) of sampling	Link to HELCOM core indicators ⁴	Spatial scope	Monitoring started (year)	CPs monitoring ⁵
Other	1.Organochl orine pesticides (DDT and its metabolites, hexachloroc yclohxsane, hexachlorob enzene, endosulfan) 2. Mustard gas with its derivatives	Concentration of chemical/nutri ent/pollutant in/on seabed substrate	ISO/IEC 17025	ISO/IEC 17025	Other			EEZ/ Transitional waters/ Coastal waters	1 – 1998 2 - 2020	PL
Regional (MORS)	Radionuclid es: Gammaemi ers	Concentration of chemical/nutri ent/pollutant in/on seabed substrate	HELCOM guidelines for Monitoring of Radioactive Substances	HELCOM guidelines for Monitoring of Radioactive Substances	Yearly	See map for details		EEZ	1984	All HELCOM Contracng Pares

Current means of coordination	Features or Elements	Parameter	Method	QA/QC	Freque ncy ³	Spatial resolution (density) of sampling	Link to HELCOM core indicators ⁴	Spatial scope		CPs monitoring ⁵
National	PBDEs, PFASs, organotins (in addition to TBT), nonylphenol s, octylphenol s, phthalates, HBCDD, pesticides (atrazine, chlorpyrifos, isoproturon e, diurone, simazine, trifluraline, chlorfenvinf os, alachlor, DCOIT, capsaicin, cybutryn (Irgarol), HCHs, DDTs, endosulfane s), other chlorinated	Concentration of chemical/nutri ent/pollutant in/on seabed substrate	https://www.n aturvardsverke t.se/upload/sto d-i- miljoarbetet/v agledning/milj oovervakning/ handledning/m etoder/unders okningstyper/l andskap/under sokningstyp- organiska- miljogifter- sedimnet- 2016-06-29.pdf https://www.n aturvardsverke t.se/upload/st od-i- miljoarbetet/v agledning/milj oovervakning/ handledning/M anualer/Metall er-sediment- 2017-12-20.pdf	See Method	Every six years (analyt es have change d betwee n years, some only analyse d one year)	13 stations (Bothnian Bay, Bothnian Sea, Northern Baltic Proper, Southern Baltic Proper		EEZ, territorial waters	2003 (repeated every 6 years)	SE

Current means of coordination	Features or Elements	Parameter	Method	QA/QC	Freque ncy ³	, ,,	Link to HELCOM core indicators ⁴	Spatial scope		CPs monitoring ⁵
National	Nonylpheno	Concentration			Yearly	Mainly		Coastal	2018	DK
	Is and	of				coastal		waters		
	phthalates	chemical/nutri				stations				
		ent/pollutant								
		in/on seabed								
		substrate								

PARAMETER

Element/Parameter pair

PCBs / Concentration of chemical/pollutant in/on seabed substrate

METHOD (MonitoringDetails)

Element/parameter

Sampling and analytical methods are reported per sample and per parameter respectively in the data. See HELCOM COMBINE manual.

QA/QC

Element/Parameter pair

Quality assurance is a laboratory's whole sampling and analytical process from start to finish. See HELCOM COMBINE manual.

In Sweden: https://www.naturvardsverket.se/upload/stod-i-miljoarbetet/vagledning/miljoovervakning/handledning/metoder/undersokningstyper/landskap/undersokningstyp-organiska-miljogifter-sedimnet-2016-06-29.pdf

FREQUENCY

Frequency

Element/Parameter pair

Varies from 1-2 to 24 samples/station/year – every six years, depending on country.

DE- BSH: May/Jun, annually IOW: Jun/Jul, annually LLUR: Jul/Aug, every 2nd year LUNG: Jul/Aug/Sep, every 3rd year

PL: 8 samples/layer every six years (4 stations); 8 samples/layer every three years in transitional waters (2 stations).

SE: Every six years.

EE: rotation, three times per 6 years, planning to reduce to one time per 6 years.

SPATIAL SCOPE

Spatial Scope

Element/Parameter pair

EEZ / Transitional waters / WFD Coastal water boides/ Whole Balc Sea for assessment

SPATIAL RESOLUTION (DENSITY) OF SAMPLING

Spatial resolution

Element/Parameter pair

Measured in the following HELCOM sub basins: Kiel Bay, Kattegat, The Sound, Great Belt, Bay of Mecklenburg and Lile Belt, Southern Baltic Proper, Northern Baltic Proper, Bothnian Sea and Bothnian Bay, Gulf of Gdansk. Gulf of Finland, Gulf of Riga.

See map for details.

PARAMETER

Element/Parameter pair

PAH / Concentration of chemical/pollutant in/on seabed substrate

METHOD (Monitoring Details)

Element/parameter

Sampling and analytical methods are reported per sample and per parameter respectively in the data. See HELCOM COMBINE manual.

QA/QC

Element/Parameter pair

Quality assurance is a laboratory's whole sampling and analytical process from start to finish. See HELCOM COMBINE manual.

FREQUENCY

Frequency

Element/Parameter pair

Varies from 1-2 to 24 samples/station/year – every six years, depending on country.

PL: 8 samples/layer every six years (4 stations); 8 samples/layer every three years in transitional waters (2 stations) and 1 sample every year (5 stations)

EE: rotation, three times per 6 years, planning to reduce to one time per 6 years.

SPATIAL SCOPE

Spatial Scope

Element/Parameter pair

EEZ / Transitional waters/ Coastal waters / Whole Baltic Sea for assessment.

DK: every third year to yearly from 2000-2017

EE: WFD coastal waters

SPATIAL RESOLUTION (DENSITY) OF SAMPLING

Spatial resolution

Element/Parameter pair

Measured in: Kiel Bay, Kattegat, Great Belt, Bay of Mecklenburg, Little Belt, The Sound, Southern Baltic Proper, Northern Baltic Proper, Bothnian Sea and Bothnian Bay, Gulf of Gdansk. Gulf of Finland, Gulf of Riga.

See map for details

PARAMETER

Element/Parameter pair

TBT / Concentration of chemical/pollutant in/on seabed substrate

METHOD (Monitoring Details)

Element/parameter

Sampling and analytical methods are reported per sample and per parameter respectively in the data. See HELCOM COMBINE manual.

QA/QC

Element/Parameter pair

Quality assurance is a laboratory's whole sampling and analytical process from start to finish. See HELCOM COMBINE manual

In Sweden: https://www.naturvardsverket.se/upload/stod-i-miljoarbetet/vagledning/miljoovervakning/handledning/metoder/undersokningstyper/land

FREQUENCY

Frequency

Element/Parameter pair

Varies from 1-2 to 24 samples/station/year – every six years, depending on country.

SE: Every six years

PL: 8 samples/layer every six years (4 stations); 8 samples/layer every three years in transitional waters (2 stations)

DK: every third year to yearly from 2000-2017

EE: rotation, three times per 6 years, planning to reduce to one time per 6 years.

SPATIAL SCOPE

Spatial Scope

Element/Parameter pair

EEZ / Transitional waters / Coastal waters / Whole Baltic Sea for assessment.

SPATIAL RESOLUTION (DENSITY) OF SAMPLING

Spatial resolution

Element/Parameter pair

Measured in: Kiel Bay, Kattegat, Great Belt, Bay of Mecklenburg, Little Belt, The Sound, Southern Baltic Proper, Northern Baltic Proper, Bothnian Sea and Bothnian Bay, Gulf of Gdansk, Bothnian Sea, Gulf of Finland, Gulf of Riga.

See map for details

PARAMETER

Element/Parameter pair

Metals / Concentration of chemical/pollutant in/on seabed substrate

METHOD (Monitoring Details)

Element/parameter

Sampling and analytical methods are reported per sample and per parameter respectively in the data. See HELCOM COMBINE manual.

QA/QC

Element/Parameter pair

Quality assurance is a laboratory's whole sampling and analytical process from start to finish. See HELCOM COMBINE manual – Part B, Annex B13, Appendix 3 and 4.

In Sweden: https://www.naturvardsverket.se/upload/stod-i-miljoarbetet/vagledning/miljoovervakning/handl

FREQUENCY

Frequency

Element/Parameter pair

Varies from 1-2 to 24 samples/station/year – every six years, depending on country.

DE - BSH: Mar, annually IOW: Jun/Jul, annually LLUR: Jul/Aug, every 2nd year LUNG: Jul/Aug/Sep, every 3rd year.

SE: Every six years.

PL: 8 samples/layer every six years (4 stations); 8 samples/layer every three years in transitional waters (2 stations) for Hg, Cd, Pb, Cu, Zn, Al and 1 sample every year (9 stations) for As.

DK: every third year to yearly from 2000-2017.

EE: rotation, three times per 6 years, planning to reduce to one time per 6 years.

SPATIAL SCOPE

Spatial Scope

Element/Parameter pair

EEZ / Transitional waters/ Coastal waters /Whole Baltic Sea for assessment.

SPATIAL RESOLUTION (DENSITY) OF SAMPLING

Spatial resolution

Element/Parameter pair

Measured in: Bay of Mecklenburg, Kiel Bay, Kaegat, The Sound, Great Belt, Lile Belt, Southern Baltic Proper, Northern Baltic Proper, Bothnian Sea and Bothnian Bay, Gulf of Gdansk. Gulf of Finland, Gulf of Riga.

See map for details.

PARAMETER

Element/Parameter pair

Other contaminants / Concentration of chemical/pollutant in/on seabed substrate

METHOD (Monitoring Details)

Element/parameter

Sampling and analytical methods are reported per sample and per parameter respectively in the data. See HELCOM COMBINE manual.

QA/QC

Element/Parameter pair

Quality assurance is a laboratory's whole sampling and analytical process from start to finish. See HELCOM COMBINE manual

FREQUENCY

Frequency

Element/Parameter pair

Varies from 1-2 to 24 samples/station/year, depending on country.

PL: 8 samples/layer every six years (4 stations); 8 samples/layer every three years in transitional waters (2 stations) for Organochlorine pesticides and 1 sample every year (6 stations) for Mustard gas with its derivatives.

SPATIAL SCOPE

Spatial Scope

Element/Parameter pair

EEZ / Transitional waters/ Coastal waters / Whole Baltic Sea for assessment.

SPATIAL RESOLUTION (DENSITY) OF SAMPLING

Spatial resolution

Element/Parameter pair

All other contaminants in sediment are measured in: Kiel Bay, Kattegat, Southern Baltic Proper, The Sound, Great Belt, Bay of Mecklenburg, Little Belt. For CEMP: furans are monitored in: Kattegat and Skagerrak, TBT only in Kattegat. National monitoring is only reported for Denmark and

Germany, of which furans are monitored in Great Belt, Kiel Bay, Little Belt. TBT is measured in Bay of Mecklenburg, Kiel Bay, Little Belt, Great Belt. Metals are reported from Great Belt Kiel Bay Bay of Mecklenburg Little Belt. All the remaining national contaminants data is reported from Great Belt, Kiel Bay, Bay of Mecklenburg and Little Belt.

PARAMETER

Element/Parameter pair

Radionuclides / Concentration of chemical/pollutant in/on seabed substrate

METHOD (Monitoring Details)

Element/parameter

Obligatory radionuclides: Gamma-emitters: K-40, Cs-137 and other γ -emitters identified in the γ -spectrum. Voluntary radionuclides: Sr-90; Pu-239, 240; Am-241; natural radionuclides (e.g. Po-210).

Sampling and analytical methods are reported per sample and per parameter respectively in the data. See MORS Guidelines.

QA/QC

Element/Parameter pair

Quality assurance is a laboratory's whole sampling and analycal process from start to finish. MORS Guidelines define methodologies for sample treatment, analysis and intercomparison. Reported data is manually quality assured by HELCOM Secretariat and results reported and verified in annual MORS EG meetings.

FREQUENCY

Frequency

Element/Parameter pair

Annual

SPATIAL SCOPE

Spatial Scope

Element/Parameter pair

EEZ / Whole Baltic Sea for assessment.

SPATIAL RESOLUTION (DENSITY) OF SAMPLING

Spatial resolution

 	
Element/Parameter pair	
See map for details	

Provide considerations for the scale of aggregation of data for an indicator-based assessment Tick one or more relevant boxes below:
⊠ HELCOM assessment unit Level 4: Subbasins with coastal WFD division
\square HELCOM assessment unit Level 3: Subbasins with coastal and offshore division
⊠ HELCOM assessment unit Level 2: Subbasin
⊠ HELCOM assessment unit Level 1: Baltic Sea
☐MSFD Region
□Other (specify)
□Unknown

c.5 Monitoring and assessment requirements

Monitoring requirements:

Core indicators rely primarily on monitoring data from biota, secondarily from sediments and lastly from water. Under the Water Framework Direcve, no EQS values have so far been set in sediments. However, countries are encouraged to perform long term trend analysis of concentraons of persistent substances that tend to accumulate in sediment, in order to ensure that such concentraons do not significantly increase (such as Hg, PAHs, HCHs, HCB, PBDE, TBT, PCB, PFOS, PCDD/F, HBCDD). To this end, sediments are suitable for revealing past recent history of contaminants.

The MSFD descriptors and associated criteria relevant to the monitoring programme are given in question on MSFD GES Criteria above.

Adequacy for assessment of GES:

Monitoring should provide adequate data and information to enable the periodic assessment of environmental status, and distance from and progress towards GES as required by MSFD under Article 9 and Article 11.

	Yes	No
Adequate data?	\boxtimes	
Established methods for		

assessment?			
Adequate unders	tanding of GES?	\boxtimes	
Adequate capacit assessments?	y to perform		
Assessment of na	tural variability		
Not relevant.			
	oviders and base the data can be		the relevant boxes below:
☐ HELCOM COMBINE	☐ HELCOM PLC	⊠HELCOM MORS	;
⊠Other:	ICES DOME		
the HELCOM Secr	etariat will do it)		stions (In case the answer is a HELCOM databas
	the relevant boxes	below:	
☐ Unprocessed/ra			
☐ Data Products	3013		
☐Modelled data			
Data managemer	nt: General descript	ion of data manageme	ent (DataManagement, Free text)
National.			
what method/method/method		sed to make the data a	available? Tick the relevant boxes below and
\square Providing URL	to view data: Click	here to enter text.	
\square Providing URL	to download data: (Click here to enter text.	i.
☐ Provide location	on of data in nationa	al data centre: Click he	ere to enter text.
☐ Provide location text.	on of data in interna	ational data centre (e. _i	.g. RSC, ICES, EEA, EMODnet): Click here to en
When will the da	ta first become ava	ilable? (DataPublicatio	onDate)
Enter the date of	reporting, or even a	a past date if desired (N	MM/YYYY):

EE: March next year after monitoring.		
How frequently are the	data expected to be updated thereafter? Tick the relevant box below:	
☐ Every 6 years	□Weekly	
☐ Every 3 years	□Daily	
☐ Every 2 years	□Hourly	
⊠Yearly	□ Continually	
\Box 6-monthly	⊠One-off	
\square 3-monthly	☐As needed	
\square Monthly	□Other (specify) Click here to enter text.	
\square 2-weekly	□Unknown	
	nts in the Contracting Parties	
EE: Estonian Environmer	nt Agency	
Has the data been used or	r is it planned to be used in HELCOM assessments? Tick the relevant box below:	
⊠Yes □No		
Select if data is used in the following Baltic Sea Environment Fact Sheets (BSEF) Tick the relevant boxes below:		
Biodiversity		
☐Abundance and distribu	tion of marenzelleria species	
☐Abundance and distribu	tion of Round goby	
☐Abundance and distribu	tion of the Zebra mussel	
☐Biopollution level index		
☐Observed non-indigenor	us and cryptogenic species in the Baltic Sea	
\square Population developmen	t of Great Cormorant	
☐ Population development of Sandwich Tern		
Population development of Southern Dunlin		
☐ Population Development of White-tailed Sea Eagle		
☐ Temporal development of Baltic coastal fish communities and key species		
Eutrophication		
• □Bacterioplankton growtl	h	
,		

☐ Chlorophyll-a concentrations, temporal variations and regional differences from satellite remote sensing	
☐ Cyanobacteria biomass	
☐ Cyanobacterial blooms in the Baltic Sea	
□Cyanobacteria bloom index	
☐ Impacts of invasive phytoplankton species on the Baltic Sea ecosystem in 1980-2008	
□ Nitrogen atmospheric deposition to the Baltic Sea	
□ Nitrogen emissions to the air in the Baltic Sea area	
☐ Phytoplankton biomass and species succession	
☐ Shifts in the Baltic Sea summer phytoplankton communities in 1992-2006	
☐ Spatial distribution of the winter nutrient pool	
☐Unusual phytoplankton event	
Hazardous substances	
☐ Atmospheric deposition of heavy metals on the Baltic Sea	
☐ Atmospheric deposition of PCDD/Fs on the Baltic Sea	
☐ Atmospheric emissions of heavy metals in the Baltic Sea region	
☐ Atmospheric emissions of PCDD/Fs in the Baltic Sea region	
⊠Cesium-137 in Baltic Sea sediments	
\Box Temporal trends in contaminants in Herring in the Baltic Sea in the period 1980-2010	
☐ Emissions from Baltic Sea shipping	
□Illegal discharges of oil in the Baltic Sea	
☐ Liquid discharges of Cs-137, Sr-90 and Co-60 into the Baltic Sea	
\Box Trace metal concentrations and trends in Baltic surface and deep waters	
Hydrography	
\square Development of Sea Surface Temperature in the Baltic Sea	
☐ Hydrography and Oxygen in the Deep Basins	
□Ice season	
☐ Total and regional runoff to the Baltic Sea	
\square Water Exchange between the Baltic Sea and the North Sea, and conditions in the Deep Basins	
☐ Wave climate in the Baltic Sea	
c.7 MSFD Criteria (GES criteria)	
Choose only the most relevant option(s). Tick one or more boxes below.	
Descriptor 1	

	The mortality rate per species from incidental by-catch is below levels which threaten the species, such that its long- term viability is ensured.
	Member States shall establish the threshold values for the mortality rate from incidental by-catch per species, through regional or subregional cooperation.
	□ D1C2 – Primary:
	The population abundance of the species is not adversely affected due to anthropogenic pressures, such that its long-term viability is ensured.
	Member States shall establish threshold values for each species through regional or subregional cooperation, taking account of natural variation in population size and the mortality rates derived from D1C1, D8C4 and D10C4 and other relevant pressures. For species covered by Directive 92/43/EEC, these values shall be consistent with the Favourable Reference Population values established by the relevant Member States under Directive 92/43/EEC.
	☐ D1C3 — Primary for commercially- exploited fish and cephalopods and secondary for other species:
	The population demographic characteristics (e.g. body size or age class structure, sex ratio, fecundity, and survival rates) of the species are indicative of a healthy population which is not adversely affected due to anthropogenic pressures.
	Member States shall establish threshold values for specified characteristics of each species through regional or subregional cooperation, taking account of adverse effects on their health derived from D8C2, D8C4 and other relevant pressures.
	☐ D1C4 — Primary for species covered by Annexes II, IV or V to Directive 92/43/EEC and secondary for other species:
	The species distributional range and, where relevant, pattern is in line with prevailing physiographic, geographic and climatic conditions.
	Member States shall establish threshold values for each species through regional or subregional cooperation. For species covered by Directive 92/43/EEC, these shall be consistent with the Favourable Reference Range values established by the relevant Member States under Directive 92/43/EEC.
	☐ D1C5 – Primary for species covered by Annexes II, IV and V to Directive 92/43/EEC and secondary for other species:
	The habitat for the species has the necessary extent and condition to support the different stages in the life history of the species.
	□ D1C6 – Primary
	The condition of the habitat type, including its biotic and abiotic structure and its functions (e.g. its typical species composition and their relative abundance, absence of particularly sensitive or fragile species or species providing a key function, size structure of species), is not adversely affected due to anthropogenic pressures.
	Member States shall establish threshold values for the condition of each habitat type, ensuring compatibility with related values set under Descriptors 2, 5 and 8, through regional or subregional cooperation.
Descriptor 2	□ D2C1 – Primary:

	The number of non-indigenous species which are newly introduced via human activity into the wild, per assessment period (6 years), measured from the reference year as reported for the initial assessment under Article 8(1) of Directive 2008/56/EC, is
	minimised and where possible reduced to zero.
	Member States shall establish the threshold value for the number of new introductions of non-indigenous species, through regional or subregional cooperation.
	☐ D2C2 — Secondary:
	Abundance and spatial distribution of established non-indigenous species, particularly of invasive species, contributing significantly to adverse effects on particular species groups or broad habitat types.
	□ D2C3 — Secondary:
	Proportion of the species group or spatial extent of the broad habitat type which is adversely altered due to non-indigenous species, particularly invasive non-indigenous species.
	Member States shall establish the threshold values for the adverse alteration to species groups and broad habitat types due to non-indigenous species, through regional or subregional cooperation.
Descriptor 3	□ D3C1 — Primary:
	The Fishing mortality rate of populations of commercially-exploited species is at or below levels which can produce the maximum sustainable yield (MSY). Appropriate scientific bodies shall be consulted in accordance with Article 26 of Regulation (EU) No 1380/2013.
	□ D3C2 — Primary:
	The Spawning Stock Biomass of populations of commercially-exploited species are above biomass levels capable of producing maximum sustainable yield. Appropriate scientific bodies shall be consulted in accordance with Article 26 of Regulation (EU) No 1380/2013.
	□ D3C3 — Primary:
	The age and size distribution of individuals in the populations of commercially-exploited species is indicative of a healthy population. This shall include a high proportion of old/large individuals and limited adverse effects of exploitation on genetic diversity.
	Member States shall establish threshold values through regional or subregional cooperation for each population of species in accordance with scientific advice obtained pursuant to Article 26 of Regulation (EU) No 1380/2013.
Descriptor 4	□ D4C1 — Primary:
	The diversity (species composition and their relative abundance) of the trophic guild is not adversely affected due to anthropogenic pressures.
	Member States shall establish threshold values through regional or subregional cooperation.
	□ D4C2 — Primary:

	The balance of total abundance between the trophic guilds is not adversely affected due to anthropogenic pressures.
	Member States shall establish threshold values through regional or subregional cooperation.
	☐ D4C3 — Secondary:
	The size distribution of individuals across the trophic guild is not adversely affected due to anthropogenic pressures.
	Member States shall establish threshold values through regional or subregional cooperation.
	\square D4C3 — Secondary (to be used in support of criterion D4C2, where necessary):
	Productivity of the trophic guild is not adversely affected due to anthropogenic pressures.
	Member States shall establish threshold values through regional or subregional cooperation.
Descriptor 5	□ D5C1 — Primary:
	Nutrient concentrations are not at levels that indicate adverse eutrophication effects.
	The threshold values are as follows:
	(a) in coastal waters, the values set in accordance with Directive 2000/60/EC;
	(b) beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation
	□ D5C2 — Primary:
	Chlorophyll a concentrations are not at levels that indicate adverse effects of nutrient enrichment.
	The threshold values are as follows:
	(c) in coastal waters, the values set in accordance with Directive 2000/60/EC;
	(d) beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation.
	□ D5C3 — Secondary:
	The number, spatial extent and duration of harmful algal bloom events are not at levels that indicate adverse effects of nutrient enrichment.
	□ D5C4 — Secondary:
	The photic limit (transparency) of the water column is not reduced, due to increases in suspended algae, to a level that indicates adverse effects of nutrient enrichment.
	The threshold values are as follows:
	(e) in coastal waters, the values set in accordance with Directive 2000/60/EC;
	(f) beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through

	regional or subregional cooperation.
	\square D5C5 — Primary (may be substituted by D5C8):
	The concentration of dissolved oxygen is not reduced, due to nutrient enrichment, to levels that indicate adverse effects on benthic habitats (including on associated biota and mobile species) or other eutrophication effects.
	The threshold values are as follows:
	(g) in coastal waters, the values set in accordance with Directive 2000/60/EC;
	(h) beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation.
	□ D5C6 — Secondary:
	The abundance of opportunistic macroalgae is not at levels that indicate adverse effects of nutrient enrichment.
	The threshold values are as follows:
	(a) in coastal waters, the values set in accordance with Directive 2000/60/EC;
	(b) should this criterion be relevant for waters beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation.
	□ D5C7 — Secondary:
	The species composition and relative abundance or depth distribution of macrophyte communities achieve values that indicate there is no adverse effect due to nutrient enrichment including via a decrease in water transparency, as follows:
	(a) in coastal waters, the values set in accordance with Directive 2000/60/EC;
	(b) should this criterion be relevant for waters beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation.
	\square D5C8 — Secondary: (except when used as a substitute for D5C5):
	The species composition and relative abundance of macrofaunal communities, achieve values that indicate that there is no adverse effect due to nutrient and organic enrichment, as follows:
	(a) in coastal waters, the values for benthic biological quality elements set in accordance with Directive 2000/60/EC;
	(b) beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation.
Descriptor 6	□ D6C1 – Primary:
	Spatial extent and distribution of physical loss (permanent change) of the natural seabed.

	□ D6C2 – Primary:
	Spatial extent and distribution of physical disturbance pressures on the seabed.
	☐ D6C3 — Primary:
	Spatial extent of each habitat type which is adversely affected, through change in its biotic and abiotic structure and its functions (e.g. through changes in species composition and their relative abundance, absence of particularly sensitive or fragile species or species providing a key function, size structure of species), by physical disturbance.
	Member States shall establish threshold values for the adverse effects of physical disturbance, through regional or subregional cooperation.
	□ D6C4 – Primary:
	The extent of loss of the habitat type, resulting from anthropogenic pressures, does not exceed a specified proportion of the natural extent of the habitat type in the assessment area.
	Member States shall establish the maximum allowable extent of habitat loss as a proportion of the total natural extent of the habitat type, through cooperation at Union level, taking into account regional or subregional specificities.
	□ D6C5 – Primary:
	The extent of adverse effects from anthropogenic pressures on the condition of the habitat type, including alteration to its biotic and abiotic structure and its functions (e.g. its typical species composition and their relative abundance, absence of particularly sensitive or fragile species or species providing a key function, size structure of species), does not exceed a specified proportion of the natural extent of the habitat type in the assessment area.
	Member States shall establish threshold values for adverse effects on the condition of each habitat type, ensuring compatibility with related values set under Descriptors 2, 5, 6, 7 and 8, through cooperation at Union level, taking into account regional or subregional specificities. Member States shall establish the maximum allowable extent of those adverse effects as a proportion of the total natural extent of the habitat type, through cooperation at Union level, taking into account regional or subregional specificities.
Descriptor 7	□ D7C1 – Secondary:
	Spatial extent and distribution of permanent alteration of hydrographical conditions (e.g. changes in wave action, currents, salinity, temperature) to the seabed and water column, associated in particular with physical loss(1) of the natural seabed.
	□ D7C2 – Secondary:
	Spatial extent of each benthic habitat type adversely affected (physical and hydrographical characteristics and associated biological communities) due to permanent alteration of hydrographical conditions.
Descriptor 8	⊠ D8C1 – Primary:
	Within coastal and territorial waters, the concentrations of contaminants do not exceed the following threshold values:

	(a) for contaminants set out under point 1(a) of criteria elements, the values set in accordance with Directive 2000/60/EC;
	(b) when contaminants under point (a) are measured in a matrix for which no value is set under Directive 2000/60/EC, the concentration of those contaminants in that matrix established by Member States through regional or subregional cooperation;
	(c) for additional contaminants selected under point 1(b) of criteria elements, the concentrations for a specified matrix (water, sediment or biota) which may give rise to pollution effects. Member States shall establish these concentrations through regional or subregional cooperation, considering their application within and beyond coastal and territorial waters.
	Beyond territorial waters, the concentrations of contaminants do not exceed the following threshold values:
	(a) for contaminants selected under point 2(a) of criteria elements, the values as applicable within coastal and territorial waters;
	(b) for contaminants selected under point 2(b) of criteria elements, the concentrations for a specified matrix (water, sediment or biota) which may give rise to pollution effects. Member States shall establish these concentrations through regional or subregional cooperation.
	□ D8C2 – Secondary:
	The health of species and the condition of habitats (such as their species composition and relative abundance at locations of chronic pollution) are not adversely affected due to contaminants including cumulative and synergetic effects.
	Member States shall establish those adverse effects and their threshold values through regional or subregional cooperation.
	□ D8C3 – Primary:
	The spatial extent and duration of significant acute pollution events are minimised.
	$\ \square$ D8C4 – Secondary (to be used when a significant acute pollution event has occurred):
	The adverse effects of significant acute pollution events on the health of species and on the condition of habitats (such as their species composition and relative abundance) are minimised and, where possible, eliminated.
Descriptor 9	☐ D9C1 – Primary:
	The level of contaminants in edible tissues (muscle, liver, roe, flesh or other soft parts, as appropriate) of seafood (including fish, crustaceans, molluscs, echinoderms, seaweed and other marine plants) caught or harvested in the wild (excluding fin-fish from mariculture) does not exceed:
	(a) for contaminants listed in Regulation (EC) No 1881/2006, the maximum levels laid down in that Regulation, which are the threshold values for the purposes of this Decision;
	(b) for additional contaminants, not listed in Regulation (EC) No 1881/2006, threshold values, which Member States shall establish through regional or

	subregional cooperation.
Descriptor 10	□ D10C1 – Primary:
	The composition, amount and spatial distribution of litter on the coastline, in the surface layer of the water column, and on the seabed, are at levels that do not cause harm to the coastal and marine environment.
	Member States shall establish threshold values for these levels through cooperation at Union level, taking into account regional or subregional specificities.
	□ D10C2 — Primary:
	The composition, amount and spatial distribution of micro-litter on the coastline, in the surface layer of the water column, and in seabed sediment, are at levels that do not cause harm to the coastal and marine environment.
	Member States shall establish threshold values for these levels through cooperation at Union level, taking into account regional or subregional specificities.
	□ D10C3 — Secondary:
	The amount of litter and micro-litter ingested by marine animals is at a level that does not adversely affect the health of the species concerned. Member States shall establish threshold values for these levels through regional or subregional cooperation.
	□ D10C4 — Secondary:
	The number of individuals of each species which are adversely affected due to litter, such as by entanglement, other types of injury or mortality, or health effects. Member States shall establish threshold values for the adverse effects of litter, through regional or subregional cooperation.
Descriptor 11	□ D11C1 – Primary:
	The spatial distribution, temporal extent, and levels of anthropogenic impulsive sound sources do not exceed levels that adversely affect populations of marine animals.
	Member States shall establish threshold values for these levels through cooperation at Union level, taking into account regional or subregional specificities.
	□ D11C2 – Primary:
	The spatial distribution, temporal extent and levels of anthropogenic continuous low-frequency sound do not exceed levels that adversely affect populations of marine animals.
	Member States shall establish threshold values for these levels through cooperation at Union level, taking into account regional or subregional specificities.

d. References

Make a list of cited references and literature for further supportive information.

HELCOM monitoring guidelines for sediments (https://helcom.fi/action-areas/monitoring-and-

assessment/monitoring-guidelines/): Monitoring of Radioactive Substances, determination of heavy metals in sediment, determination of chlorinated hydrocarbons in sediment, determination of PAH in sediment. EU. 2010. Guidance Document No. 25. Guidance on chemical monitoring of sediment and biota under thewater framework directive.ment, determination of PAH in sediment. EU. 2010. Guidance Document No. 25. Guidance on chemical monitoring of sediment and biota under thewater framework directive.HELCOM monitoring guidelines for sediments (https://helcom.fi/action-areas/monitoring-and-assessment/monitoring-guidelines/): Monitoring of Radioactive Substances, determination of heavy metals in sediment, determination of chlorinated hydrocarbons in sedi