HELCOM Monitoring Programme topic

Concentrations of contaminants

Programme:

Marine bird health

Contents

a.	Metadata on monitoring strategies and monitoring programmes	2
	a.1 Responsible HELCOM subsidiary body	2
	a.2 Regional Cooperation (RegionalCooperation)	3
b.	Monitoring strategies	3
	b.1 Descriptor	3
	b.2 BSAP segments	4
	b.3 Monitoring strategy description	4
	b.4 BSAP Ecological objectives	4
	b.5 Gaps in monitoring	5
c.	Monitoring programmes	5
	c.1 Purpose of monitoring	5
	c.2 Other legislation	9
	c.3 Implementation of Regional Cooperation (Regional Cooperation_implementation)	10
	c.4 Monitoring concepts	11
	c.5 Monitoring and assessment requirements	14
	c.6 Data providers and access	15
	c.7 MSFD Criteria (GES criteria)	17
٦	Deferences	2.4

a. Metadata on monitoring strategies and monitoring programmes

a.1 Responsible HELCOM subsidiary body

Please indicate the relevant expert group/network if available, otherwise the responsible HELCOM Working Group. or the responsible working group understanding is that they are the EN or EG if available, otherwise the WGs

	Permament Groups			
	Gear – Group on the Implementation of the Ecosystem Approach			
	Maritime – Maritime Working Group			
	Pressure – Working Group on Reduction of Pressures from the Baltic Sea Catchment Area			
	Response – Response Working Group			
\boxtimes	State and Conservation – Working Group on the State of the Environmental and Nature Conservation			
	Time-limited Groups			
	Agri – Group on Sustainable Agricultural Practices			
	Fish – Group on Ecosystem-based Sustainable Fisheries			
	HELCOM-VASAB MSP WG - Joint HELCOM-VASAB Maritime Spatial Planning Working Group			
	Expert Groups			
	Expert Groups AIS EWG – Expert Working Group for Mutual Exchange and Deliveries of AIS data			
	AIS EWG – Expert Working Group for Mutual Exchange and Deliveries of AIS data			
	AIS EWG – Expert Working Group for Mutual Exchange and Deliveries of AIS data EN Hazardous Substances – Expert Network on hazardous substances			
	AIS EWG – Expert Working Group for Mutual Exchange and Deliveries of AIS data EN Hazardous Substances – Expert Network on hazardous substances EN Marine Litter – Expert Network on Marine Litter			
	AIS EWG – Expert Working Group for Mutual Exchange and Deliveries of AIS data EN Hazardous Substances – Expert Network on hazardous substances EN Marine Litter – Expert Network on Marine Litter EN Noise – Expert Network on Underwater Noise			
	AIS EWG – Expert Working Group for Mutual Exchange and Deliveries of AIS data EN Hazardous Substances – Expert Network on hazardous substances EN Marine Litter – Expert Network on Marine Litter EN Noise – Expert Network on Underwater Noise ESA – Expert Network on Economic and Social Analyses			
	AIS EWG – Expert Working Group for Mutual Exchange and Deliveries of AIS data EN Hazardous Substances – Expert Network on hazardous substances EN Marine Litter – Expert Network on Marine Litter EN Noise – Expert Network on Underwater Noise ESA – Expert Network on Economic and Social Analyses EWG OWR – Expert Working Group on Oiled Wildlife Response			
	AIS EWG – Expert Working Group for Mutual Exchange and Deliveries of AIS data EN Hazardous Substances – Expert Network on hazardous substances EN Marine Litter – Expert Network on Marine Litter EN Noise – Expert Network on Underwater Noise ESA – Expert Network on Economic and Social Analyses EWG OWR – Expert Working Group on Oiled Wildlife Response EWG SHORE – Expert Working Group on Response on the Shore			

	IWGAS – Informal Working Group on Aerial Surveillance
\boxtimes	JWG Bird – HELCOM-OSPAR-ICES Joint Working Group on Seabirds
	MORS EG – Expert group on monitoring of radioactive substances in the Baltic Sea
	PRF Cooperation Platform – Cooperation Platform on Port Reception Facilities in the Baltic Sea
	SAFE NAV – Group of Experts on Safety of Navigation
	SUBMERGED – Expert Group on Environmental Risks of Hazardous Submerged Objects

a.2 Regional Cooperation (RegionalCooperation)

The monitoring of this	programme is:
------------------------	---------------

☐ Fully	coordinated
I uii	, cooraniatea

☐ Partly coordinated. Indicate missing component(s):

☑ Coordinated monitoring is under development. Indicate by which group/project and by when a recommendation on coordinated monitoring can be expected.

Common monitoring guidelines: missing.

Common quality assurance programme: missing. National QA/QC exist. Common database: missing

b. Monitoring strategies

b.1 Descriptor

The programme supports the following obligatory MSFD Monitoring Strategies. Tick one or more relevant

 ☑ D1 Biodiversity ☑ D2 Non-indigenous Species ☑ D3 Commercial fish and shellfish ☑ D4 Food webs ☑ D5 Eutrophication ☑ D6 Seafloor integrity 	
 □ D3 Commercial fish and shellfish □ D4 Food webs □ D5 Eutrophication 	
☑ D4 Food webs☐ D5 Eutrophication	
□ D5 Eutrophication	
·	
☐ D6 Seafloor integrity	
☐ D7 Hydrographical conditions	
□ D8 Contaminants	

□ D9	Contaminants in seafood					
□ D10	Marine litter					
□ D11	Energy including underwater noise					
b.2 BSAP so The sub-programm	egments me serves the following BSAP segments. Tick one or more relevant boxes.					
□Eutrophicatio	n					
⊠Hazardous sul	⊠ Hazardous substances					
⊠Biodiversity						
☐ Maritime activities						
b.3 Monito	ring strategy description					
	tegy: Monitoring is to be carried out to fulfill assessment requirements of HELCOM tives that are specified through HELCOM core indicators. The requirements on nclude number of stations, the sampling frequency and replication.					
	cological objectives nost relevant option(s). Tick one or more boxes below.					
Eutrophication Concentrations of nutrients close to natural levels						
	☐ Clear water					
	☐ Natural level of algal blooms					

☐ No introductions of alien species from ships
\square Minimum air pollution from ships
☐ Zero discharges from offshore platforms
b.5 Gaps in monitoring In relation to the GES criteria addressed, indicate when sufficient monitoring was in place or by when sufficient coverage will be in place (Coverage_GEScriteria)
☐ Adequate monitoring was in place in 2014
☐ Adequate monitoring was in place by 2018
☐ Adequate monitoring is in place by July 2020
☐ Adequate monitoring will be in place by 2024
\square Monitoring is not being put in place for this descriptor due to a low risk
☐ Monitoring for this descriptor is not relevant
Description of the implementation gaps and plans to complete the establishment and implementation of this descriptor monitoring strategy (Gaps_Plans):
Besides for the white-tailed eagle, monitoring for marine bird health currently does not exist.
c. Monitoring programmes
c.1 Purpose of monitoring
c.1a Assessment purpose in general The programme supports the assessment of:

Tick the relevant box	Tick	the	relevant	box
-----------------------	------	-----	----------	-----

Temporal trends	Spatial distribution	State classification
		\boxtimes

The **programme** supports the assessment of: (MonitoringPurpose).

Note that the answer to this question will be decisive for whether to answer upcoming questions e.g. upcoming questions on pressures should only be answered if the monitoring is defined as supporting the assessment of pressures.

Tick the relevant boxes.

¹ Sea-based 'Pressures at source' refers to monitoring pressures from sea-based activities where the monitoring is directly at the activity rather than at a distance from or time period after it is generated by the activity (e.g. D1 incidental by-catch when fishing, D2 ballast water discharges, D6 use of bottom fishing gear, D8 contaminant discharges and pollution events from a vessel or pipeline, D11 impulsive sound events from a vessel or platform).

\boxtimes						
If this is selected fill in the following questions:	If this is selected fill in the following questions:	If this is selected fill in the following questions:	If this is selected fill in the following questions:	If this is selected fill in the following questions:		
c.1b	c.1c, d	c.1c, d	c.1c, d	c.1c, d		
Give any other monitoring purpose e.g. if the programmes include supporting parameters for other monitoring programmes						

For questions 1b-1d, select when applicable for the sub-programme, the link from the Reporting on the 2020 update of Article 11 for the Marine Strategy Framework Directive (MSFD Guidance Document 17, 2020) (Features) to:

- Ecosystem components (relevant for monitoring and assessment for Article 8(1a) for D1C2-C5, D3, D4, D6C3-C5, D7C2)
- Pressures and impacts in the marine environment (relevant for monitoring and assessment for Article 8(1b) for D1C1, D2, D5, D6C1-C2, D7C1, D8, D9, D10, D11)
- Pressure inputs to the marine environment (relevant for monitoring and assessment for Article 10)
- Uses and human activities (relevant for monitoring and assessment for Article 8(1c) and 13)

c.1b • Ecosystem components (Features)

Choose only the most relevant option(s). Tick one or more boxes below.

Theme	Sub-theme	Label feature
Species	☐ Birds	☐ Grazing birds
		☐ Wading birds
		☐ Surface-feeding birds
		☐ Pelagic-feeding birds
		☐ Benthic-feeding birds
	☐ Mammals	☐ Small toothed cetaceans
		☐ Deep-diving toothed cetaceans
		☐ Baleen whales
		☐ Seals
	☐ Reptiles	☐ Turtles
	☐ Fish	☐ Coastal fish
		☐ Pelagic shelf fish
		☐ Demersal shelf fish
		☐ Deep-sea fish
		☐ Commercially exploited fish and shellfish

	☐ Cephalopods	\square Coastal/shelf cephalopods				
		☐ Deep-sea cephalopods				
Habitats	\square Benthic habitats	\square Benthic broad habitats				
		\square Other benthic habitats				
	☐ Pelagic habitats	☐ Pelagic broad habitats				
		\square Other pelagic habitats				
Ecosystems	☐ Physical and hydrological	characteristics				
	☐ Chemical characteristics					
	☐ Ecosystems, including	☐ Coastal ecosystems				
	food webs	☐ Shelf ecosystems				
		☐ Oceanic/deep-sea ecosystems				
	Pressures and impacts in the most relevant option(s). Tick	he marine environment (Features) ek one or more boxes below.				
Theme	Label: Feature					
Biological	☐ Newly introduced non	☐ Newly introduced non-indigenous species				
	☐ Established non-indige	☐ Established non-indigenous species				
	☐ Species affected by inc	cidental by-catch				
Physical and	☐ Hydrographical change	☐ Hydrographical changes				
hydrological	☐ Physical disturbance to seabed					
	☐ Physical loss of the seabed					
Substances,	☐ Eutrophication					
litter and energy	☐ Contaminants - non UPBT substances					
G,	☐ Contaminants - UPBT substances					
	☐ Contaminants – in seafood					
	□ Adverse effects on species or habitats					
	☐ Acute pollution events	☐ Acute pollution events				
	☐ Litter in the environme	ent				
	☐ Impulsive sound in wa	☐ Impulsive sound in water				
	☐ Continuous low freque	ency sound				
c.1d • Pressure inputs to the marine environment (Features)						
Theme	Label: Feature					

Biological	☐ Input or spread of non-indigenous species			
	☐ Input of microbial pathogens			
	$\hfill\Box$ Input of genetically modified species and translocation of native species			
	$\hfill\Box$ Loss of, or change to, natural biological communities due to cultivation of animal or plant species			
	$\hfill\Box$ Disturbance of species (e.g. where they breed, rest and feed) due to human presence			
	\square Extraction of, or mortality/injury to, wild species (by commercial and recreational fishing and other activities)			
Substances,	☐ Input of nutrients — diffuse sources, point sources, atmospheric deposition			
litter and energy	\square Input of organic matter — diffuse sources and point sources			
епегду	\square Input of other substances (e.g. synthetic substances, non-synthetic substances, radionuclides) — diffuse sources, point sources, atmospheric deposition, acute events			
	☐ Input of litter (solid waste matter, including micro-sized litter)			
	☐ Input of anthropogenic sound (impulsive, continuous)			
	\square Input of other forms of energy (including electromagnetic fields, light and heat)			
	☐ Input of water — point sources (e.g. brine)			

c.1e • Uses and human activities (Features)

Choose only the most relevant option(s). Tick one or more boxes below.

Physical restructuring of rivers, coastline or seabed (water management) Coastal defence and flood protection Goffshore structures (other than for oil/gas/renewables) Restructuring of seabed morphology, including dredging and depositing of materials Extraction of non-living resources Extraction of oil and gas, including infrastructure Extraction of salt Extraction of water Production of energy Non-renewable energy generation Non-renewable energy generation	Ineme	Label: Feature
rivers, coastline or seabed (water management) Coastal defence and flood protection Costal defence and flood protection Offshore structures (other than for oil/gas/renewables) Restructuring of seabed morphology, including dredging and depositing of materials Extraction of non-living resources Extraction of oil and gas, including infrastructure Extraction of salt Extraction of water Production of energy Renewable energy generation (wind, wave and tidal power), including infrastructure	•	☐ Land claim
management) Coastal deferte and flood protection	ŭ	☐ Canalisation and other watercourse modifications
□ Offshore structures (other than for oil/gas/renewables) □ Restructuring of seabed morphology, including dredging and depositing of materials Extraction of non-living resources □ Extraction of oil and gas, including infrastructure □ Extraction of salt □ Extraction of water Production of energy energy generation (wind, wave and tidal power), including infrastructure		☐ Coastal defence and flood protection
Extraction of non-living resources Extraction of oil and gas, including infrastructure Extraction of salt Extraction of water Production of energy Renewable energy generation (wind, wave and tidal power), including infrastructure	management	☐ Offshore structures (other than for oil/gas/renewables)
non-living resources □ Extraction of oil and gas, including infrastructure □ Extraction of salt □ Extraction of water Production of energy □ Renewable energy generation (wind, wave and tidal power), including infrastructure		
resources Extraction of oil and gas, including infrastructure Extraction of salt Extraction of water Production of energy Renewable energy generation (wind, wave and tidal power), including infrastructure	Extraction of	☐ Extraction of minerals (rock, metal ores, gravel, sand, shell)
□ Extraction of salt □ Extraction of water Production of energy □ Renewable energy generation (wind, wave and tidal power), including infrastructure	_	☐ Extraction of oil and gas, including infrastructure
Production of energy	resources	☐ Extraction of salt
energy infrastructure		☐ Extraction of water
	Production of	\square Renewable energy generation (wind, wave and tidal power), including
☐ Non-renewable energy generation	energy	infrastructure
		☐ Non-renewable energy generation

	☐ Transmission of electricity and communications (cables)			
Extraction of	☐ Fish and shellfish harvesting (professional, recreational)			
living resources	☐ Fish and shellfish processing			
	☐ Marine plant harvesting			
	☐ Hunting and collecting for other purposes			
Cultivation of	☐ Aquaculture — marine, including infrastructure			
living resources	☐ Aquaculture — freshwater			
	☐ Agriculture			
	☐ Forestry			
Transport	☐ Transport infrastructure			
	☐ Transport — shipping			
	☐ Transport — air			
	☐ Transport — land			
Urban and	☐ Urban uses			
industrial uses	☐ Industrial uses			
	☐ Waste treatment and disposal			
Tourism and	☐ Tourism and leisure infrastructure			
leisure	☐ Tourism and leisure activities			
Security/defence	☐ Military operations (subject to Article 2(2))			
Education and research	☐ Research, survey and educational activities			
c.2 Other leg The sub-programmone or more relevan	e links with the following other international legislation (OtherPoliciesConventions). Tick			
☐ Bathing Water Di	irective			
☐ Common Fisheries Policy and Data Collection Framework				
☐ Habitats Directive				
⊠ Birds Directive				
☐ Nitrates Directive				
□Urban Waste Wa	ter Treatment Directive			
⊠Water Framewor	k Directive			
□ OSPAR Convention	on			
☐Trilateral Wadde	n Sea Convention			
☑Other, Specify: CITES, Bonn Convention, Bern Convention				

c.3 Implementation of Regional Cooperation (RegionalCooperation_implementation) Indicate the level of implementation by selecting one of the following:

\square Agreed data collection methods
\square Common monitoring strategy (spatial and temporal design of programme)
\square Coordinated data collection (delivered separately by each country)
\Box Joint data collection (multinational delivery using same platform and/or algorithms)

c.4 Monitoring concepts

Monitoring concepts table²:

Current means of coordination	Features or Elements Elements (Features) (Features_e num)	Parameter Parameters (Parameter) (ParametersOth er)	Method MonitoringMetho d (Monitoring Method) MonitoringMetho dOther)	QA/QC (Free text)	Frequency ³ MonitoringFrequency	Spatial resolution (density) of sampling (ProgrammeDescription)	Link to HELCOM core indicators ⁴ (RelatedIndicator) (RelatedIndicator_n ame	Spatial scope (SpatialSco pe)	Monitorin g started (year) (TemporalSc ope)	CPs monitoring ⁵ (CountryCode_E num)
Semi- regional	White- tailed eagle	Breeding success, Nestling brood size, Producvity	National	National	Yearly	Selected coastal areas	Whitetailed eagle producvity	Coastal Waters	SE: 1964, 1965- 1973 PL: 2015	EE, DE, DK, FI, LV, LT, PL, RU, SE
Other	White- tailed eagle	Abundance, productivity	National, counting nests and reproducve parameters	National	Yearly	Location of known nests	Whitetailed eagle producvity	Coastal waters	1994	EE

-

² Needed codelists can be found on 2020 update of Article 11 for the Marine Strategy Framework Directive (MSFD Guidance Document 17, 2020).

³ The option "Different for each country - see MORE overview" refers to the <u>overview</u> carried out in 2013

⁴ Give the name of HELCOM core indicators that are based on the monitoring parameter.

⁵ Provide information on the Contracting Partie(s) that are monitoring the parameter.

PARAMETER

Element/Parameter pair

White-tailed eagle reproduction/ breeding success, nestling brood size and productivity

METHOD (Monitoring Details)

Element/parameter

Method: White-tailed eagles are resident and faithful to their territories throughout their lifetime. These features provide good opportunities for long-term monitoring of populations and breeding performance. Nest sites are checked annually for occupancy and the reproductive output is recorded for each occupied territory. Based on the frequency distribution in the population of occupied nests containing 0, 1, 2 or 3 nestlings, three reproductive parameters are assessed: the proportion of reproducing pairs in the population (Breeding success), the mean number of nestlings per successfully breeding pair (Nestling brood size) and the the mean number of nestlings per checked pair in the population (Productivity).

Based on data from nests inspected by climbing the nest tree or checked with Unmanned Airborne Systems (UAS), and excluding nests checked only from the ground, nestling brood size is a precise standard. Nest trees are climbed for precise assessment of reproductive parameters. Some samples are taken from the ground. In connection with these nest visits, measurements and biological samples are taken. The following parameters are usually measured from the nestlings: wing chord (for estimation of age in days), tarsus width and depth (for esmation of sex, see Helander 1981, Helander et al. 2007), weight (for nutrional status) and in some areas feather and blood samples (for chemical analyses and genetic studies). The nestlings are ringed using an international colour ringing programme for idenfication, according to Helander (2003b). Dead eggs and shell pieces are collected for measurements, investigation of contents and chemical analyses, for studies on relaonships with reproduction. Feathers shed from adults are generally collected. All samples collected in Sweden are archived in the National Specimen Data Bank.

QA/QC

Element/Parameter pair

National. Regional coordination to be developed for reporting of national metadata.

FREQUENCY

Frequency

Element/Parameter pair

Monitoring is done in the HELCOM Contracting Perties on an annual basis.

SPATIAL SCOPE

Spatial Scope

Element/Parameter pair

Coastal waters/ TW

SPATIAL RESOLUTION (DENSITY) OF SAMPLING

Spatial resolution

Element/Parameter pair

Eagles presently breed along the coasts and mainland shores of the whole Baltic Sea, and are monitored in a network of national projects with harmonized methodology. Monitoring is made for the entire population. There are sub-regions with small subpopulations: the Gulf of Finland, and especially the Kattegat where the species was brought to extinction in the 1800s but where resettlement is now in progress.

Provide considerations for the scale of aggregation of data for an indicator-based assessment Tick one or more relevant boxes below:

☐HELCOM assessment unit Level 4: Subbasins with coastal WFD division				
\square HELCOM assessment unit Level 3: Subbasins with coastal and offshore division				
☐ HELCOM assessment unit Level 2: Subbasin				
☐HELCOM assessment unit Level 1: Baltic Sea				
☐MSFD Region				
□EU				
⊠Other (specify) Coastal zones				
□Unknown				

c.5 Monitoring and assessment requirements

Monitoring requirements:

White-tailed sea eagle reproductive ability is monitored annually by assessing the frequency distribution of occupied eagle nests containing 0, 1, 2 or 3 nestlings (3 being the maximum in this species). These data are used for the assessment of the three indicators of reproductive ability: breeding success, nestling brood size and productivity. Survey techniques and sampling methods are presented in (Helander 1994b, Helander et al. 2007, 2009). For assessment of nutritional condition of nestlings, weight and winglength (for age) is required.

The core indicator 'Productivity of white-tailed eagle' describes not only biomagnification of contaminants, but also persecution, disturbance of nest sites, food availability and availability of suitable nesting sites. This indicator combines the breeding success and brood size into a single indicator and assesses the reproductive output of the population. It is a useful indicator in studies on relationships between reproduction and anthropogenic pressures and also a vital parameter in assessments of population status in management perspectives. Most Baltic Sea countries are monitoring white-tailed eagle and for assessment purposes monitoring should be done in all areas along the Baltic coast where the species occurs in adequate minumum abundance. In order to include white-tailed eagles that forage specifically in the coastal ecosystem zone and marine area, only breeding pairs in the coastal area should be considered (tentative boundary up to a maximum of 10 km inland from the coastal mean water line, in accordance with the guidelines of EC Nature 2-5, 1993). Other bird species are currently not being monitored for marine bird health.

Adequacy for assessment of GES:

Monitoring should provide adequate data and information to enable the periodic assessment of environmental status, and distance from and progress towards GES as required by MSFD under Article 9 and Article 11.

	Yes	No
Adequate data?	\boxtimes	
Established methods for assessment?		
Adequate understanding of GES?	\boxtimes	
Adequate capacity to perform assessments?		

Assessment of natural variability

Quantitative. Natural variability in reproductive output is very small for this species.

c.6 Data providers and access

From which database the data can be made available? Tick the relevant boxes below: ☐ HELCOM ☐ HELCOM PLC ☐ HELCOM MORS **COMBINE** \boxtimes Other: National data centres If the previous answer is "Other" please fill in the next questions (In case the answer is a HELCOM database, the HELCOM Secretariat will do it) **Data type** Tick the relevant boxes below: ☐ Unprocessed/raw Data ⊠ Processed Data sets ☐ Data Products ☐ Modelled data Data management: General description of data management (DataManagement, Free text) No information. What method/mechanism will be used to make the data available? Tick the relevant boxes below and provide location (DataAccess): ☐ Providing URL to view data: ☐ Providing URL to download data: ☑ Provide location of data in national data centre: Click here to enter text. ☐ Provide location of data in international data centre (e.g. RSC, ICES, EEA, EMODnet): When will the data first become available? (DataPublicationDate) Enter the date of reporting, or even a past date if desired (MM/YYYY): No information How frequently are the data expected to be updated thereafter? Tick the relevant box below: ☐ Every 6 years □Weekly ⊠Every 3 years ☐ Daily ☐ Every 2 years □Hourly ☐ Yearly □ Continually □6-monthly □One-off

\square 3-monthly	☐As needed	
□Monthly	□Other (specify)	
\square 2-weekly	□Unknown	
List providing contact poi	nts in the Contracting Parties	
No information	its in the contracting ranges	
No illiorillation		
Has the data been used o	r is it planned to be used in HELCOM assessments? Tick the relevant box below:	
⊠Yes □No		
Select if data is used in th below:	e following Baltic Sea Environment Fact Sheets (BSEF) Tick the relevant boxes	
Biodiversity		
☐ Abundance and distribu	ition of marenzelleria species	
☐ Abundance and distribu	ition of Round goby	
☐ Abundance and distribu	ition of the Zebra mussel	
\square Biopollution level index		
□Observed non-indigeno	us and cryptogenic species in the Baltic Sea	
☐ Population developmen	t of Great Cormorant	
☐ Population developmen	t of Sandwich Tern	
☐ Population developmen	t of Southern Dunlin	
⊠ Population Developmer	nt of White-tailed Sea Eagle	
☐ Temporal development of Baltic coastal fish communities and key species		
Eutrophication		
☐ Bacterioplankton growt	h	
☐ Chlorophyll-a concentrations, temporal variations and regional differences from satellite remote sensing		
☐ Cyanobacteria biomass		
☐ Cyanobacterial blooms in the Baltic Sea		
□Cyanobacteria bloom index		
☐ Impacts of invasive phytoplankton species on the Baltic Sea ecosystem in 1980-2008		
□ Nitrogen atmospheric deposition to the Baltic Sea		
□ Nitrogen emissions to the air in the Baltic Sea area		
☐ Phytoplankton biomass and species succession		

☐ Shifts in the Baltic Sea summer phytoplankton communities in 1992-2006		
☐ Spatial distribution of the winter nutrient pool		
☐Unusual phytoplankton event		
Hazardous subst	ances	
\square Atmospheric de	position of heavy metals on the Baltic Sea	
\square Atmospheric de	position of PCDD/Fs on the Baltic Sea	
\square Atmospheric em	nissions of heavy metals in the Baltic Sea region	
\square Atmospheric em	nissions of PCDD/Fs in the Baltic Sea region	
\square Cesium-137 in B	altic Sea sediments	
☐Temporal trends	in contaminants in Herring in the Baltic Sea in the period 1980-2010	
\square Emissions from	Baltic Sea shipping	
□Illegal discharge	s of oil in the Baltic Sea	
☐ Liquid discharge	s of Cs-137, Sr-90 and Co-60 into the Baltic Sea	
☐Trace metal con	centrations and trends in Baltic surface and deep waters	
Hydrography		
\square Development of	Sea Surface Temperature in the Baltic Sea	
\square Hydrography an	d Oxygen in the Deep Basins	
\square Ice season		
\square Total and region	al runoff to the Baltic Sea	
□Water Exchange	between the Baltic Sea and the North Sea, and conditions in the Deep Basins	
☐ Wave climate in the Baltic Sea		
c.7 MSFD Ci	riteria (GES criteria)	
	ost relevant option(s). Tick one or more boxes below.	
Descriptor 1	□ D1C1 – Primary:	
	The mortality rate per species from incidental by-catch is below levels which threaten the species, such that its long- term viability is ensured.	
	Member States shall establish the threshold values for the mortality rate from incidental by-catch per species, through regional or subregional cooperation.	
	☑ D1C2 – Primary:	
	The population abundance of the species is not adversely affected due to anthropogenic pressures, such that its long-term viability is ensured.	
	Member States shall establish threshold values for each species through regional or subregional cooperation, taking account of natural variation in population size and the	

mortality rates derived from D1C1, D8C4 and D10C4 and other relevant pressures. For species covered by Directive 92/43/EEC, these values shall be consistent with the Favourable Reference Population values established by the relevant Member States under Directive 92/43/EEC. ☑ D1C3 – Primary for commercially- exploited fish and cephalopods and secondary for other species: The population demographic characteristics (e.g. body size or age class structure, sex ratio, fecundity, and survival rates) of the species are indicative of a healthy population which is not adversely affected due to anthropogenic pressures. Member States shall establish threshold values for specified characteristics of each species through regional or subregional cooperation, taking account of adverse effects on their health derived from D8C2, D8C4 and other relevant pressures. ☐ D1C4 – Primary for species covered by Annexes II, IV or V to Directive 92/43/EEC and secondary for other species: The species distributional range and, where relevant, pattern is in line with prevailing physiographic, geographic and climatic conditions. Member States shall establish threshold values for each species through regional or subregional cooperation. For species covered by Directive 92/43/EEC, these shall be consistent with the Favourable Reference Range values established by the relevant Member States under Directive 92/43/EEC. ☐ D1C5 – Primary for species covered by Annexes II, IV and V to Directive 92/43/EEC and secondary for other species: The habitat for the species has the necessary extent and condition to support the different stages in the life history of the species. ☐ D1C6 – Primary The condition of the habitat type, including its biotic and abiotic structure and its functions (e.g. its typical species composition and their relative abundance, absence of particularly sensitive or fragile species or species providing a key function, size structure of species), is not adversely affected due to anthropogenic pressures. Member States shall establish threshold values for the condition of each habitat type, ensuring compatibility with related values set under Descriptors 2, 5 and 8, through regional or subregional cooperation. Descriptor 2 \square D2C1 – Primary: The number of non-indigenous species which are newly introduced via human activity into the wild, per assessment period (6 years), measured from the reference year as reported for the initial assssment under Article 8(1) of Directive 2008/56/EC, is minimised and where possible reduced to zero. Member States shall establish the threshold value for the number of new introductions of non-indigenous species, through regional or subregional cooperation. \square D2C2 — Secondary: Abundance and spatial distribution of established non-indigenous species, particularly of invasive species, contributing significantly to adverse effects on particular species

	groups or broad habitat types.
	□ D2C3 — Secondary:
	Proportion of the species group or spatial extent of the broad habitat type which is adversely altered due to non-indigenous species, particularly invasive non-indigenous species.
	Member States shall establish the threshold values for the adverse alteration to species groups and broad habitat types due to non-indigenous species, through regional or subregional cooperation.
Descriptor 3	□ D3C1 — Primary:
	The Fishing mortality rate of populations of commercially-exploited species is at or below levels which can produce the maximum sustainable yield (MSY). Appropriate scientific bodies shall be consulted in accordance with Article 26 of Regulation (EU) No 1380/2013.
	□ D3C2 — Primary:
	The Spawning Stock Biomass of populations of commercially-exploited species are above biomass levels capable of producing maximum sustainable yield. Appropriate scientific bodies shall be consulted in accordance with Article 26 of Regulation (EU) No 1380/2013.
	□ D3C3 — Primary:
	The age and size distribution of individuals in the populations of commercially-exploited species is indicative of a healthy population. This shall include a high proportion of old/large individuals and limited adverse effects of exploitation on genetic diversity.
	Member States shall establish threshold values through regional or subregional cooperation for each population of species in accordance with scientific advice obtained pursuant to Article 26 of Regulation (EU) No 1380/2013.
Descriptor 4	□ D4C1 — Primary:
	The diversity (species composition and their relative abundance) of the trophic guild is not adversely affected due to anthropogenic pressures.
	Member States shall establish threshold values through regional or subregional cooperation.
	□ D4C2 — Primary:
	The balance of total abundance between the trophic guilds is not adversely affected due to anthropogenic pressures.
	Member States shall establish threshold values through regional or subregional cooperation.
	□ D4C3 — Secondary:
	The size distribution of individuals across the trophic guild is not adversely affected due to anthropogenic pressures.
	Member States shall establish threshold values through regional or subregional cooperation.
	oximes D4C3 — Secondary (to be used in support of criterion D4C2, where necessary):

	Productivity of the trophic guild is not adversely affected due to anthropogenic pressures.
	Member States shall establish threshold values through regional or subregional cooperation.
Descriptor 5	□ D5C1 — Primary:
	Nutrient concentrations are not at levels that indicate adverse eutrophication effects.
	The threshold values are as follows:
	(a) in coastal waters, the values set in accordance with Directive 2000/60/EC;
	(b) beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation
	□ D5C2 — Primary:
	Chlorophyll a concentrations are not at levels that indicate adverse effects of nutrient enrichment.
	The threshold values are as follows:
	(c) in coastal waters, the values set in accordance with Directive 2000/60/EC;
	(d) beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation.
	□ D5C3 — Secondary:
	The number, spatial extent and duration of harmful algal bloom events are not at levels that indicate adverse effects of nutrient enrichment.
	□ D5C4 — Secondary:
	The photic limit (transparency) of the water column is not reduced, due to increases in suspended algae, to a level that indicates adverse effects of nutrient enrichment.
	The threshold values are as follows:
	(e) in coastal waters, the values set in accordance with Directive 2000/60/EC;
	(f) beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation.
	\square D5C5 — Primary (may be substituted by D5C8):
	The concentration of dissolved oxygen is not reduced, due to nutrient enrichment, to levels that indicate adverse effects on benthic habitats (including on associated biota and mobile species) or other eutrophication effects.
	The threshold values are as follows:
	(g) in coastal waters, the values set in accordance with Directive 2000/60/EC;
	(h) beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation.

	□ D5C6 — Secondary:
	The abundance of opportunistic macroalgae is not at levels that indicate adverse effects of nutrient enrichment.
	The threshold values are as follows:
	(a) in coastal waters, the values set in accordance with Directive 2000/60/EC;
	(b) should this criterion be relevant for waters beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation.
	□ D5C7 — Secondary:
	The species composition and relative abundance or depth distribution of macrophyte communities achieve values that indicate there is no adverse effect due to nutrient enrichment including via a decrease in water transparency, as follows:
	(a) in coastal waters, the values set in accordance with Directive 2000/60/EC;
	(b) should this criterion be relevant for waters beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation.
	\square D5C8 — Secondary: (except when used as a substitute for D5C5):
	The species composition and relative abundance of macrofaunal communities, achieve values that indicate that there is no adverse effect due to nutrient and organic enrichment, as follows:
	(a) in coastal waters, the values for benthic biological quality elements set in accordance with Directive 2000/60/EC;
	(b) beyond coastal waters, values consistent with those for coastal waters under Directive 2000/60/EC. Member States shall establish those values through regional or subregional cooperation.
Descriptor 6	□ D6C1 – Primary:
	Spatial extent and distribution of physical loss (permanent change) of the natural seabed.
	□ D6C2 – Primary:
	Spatial extent and distribution of physical disturbance pressures on the seabed.
	□ D6C3 – Primary:
	Spatial extent of each habitat type which is adversely affected, through change in its biotic and abiotic structure and its functions (e.g. through changes in species composition and their relative abundance, absence of particularly sensitive or fragile species or species providing a key function, size structure of species), by physical disturbance.
	Member States shall establish threshold values for the adverse effects of physical disturbance, through regional or subregional cooperation.
	□ D6C4 – Primary:
	The extent of loss of the habitat type, resulting from anthropogenic pressures, does not

	exceed a specified proportion of the natural extent of the habitat type in the assessment area.
	Member States shall establish the maximum allowable extent of habitat loss as a proportion of the total natural extent of the habitat type, through cooperation at Union level, taking into account regional or subregional specificities.
	□ D6C5 – Primary:
	The extent of adverse effects from anthropogenic pressures on the condition of the habitat type, including alteration to its biotic and abiotic structure and its functions (e.g. its typical species composition and their relative abundance, absence of particularly sensitive or fragile species or species providing a key function, size structure of species), does not exceed a specified proportion of the natural extent of the habitat type in the assessment area.
	Member States shall establish threshold values for adverse effects on the condition of each habitat type, ensuring compatibility with related values set under Descriptors 2, 5, 6, 7 and 8, through cooperation at Union level, taking into account regional or subregional specificities. Member States shall establish the maximum allowable extent of those adverse effects as a proportion of the total natural extent of the habitat type, through cooperation at Union level, taking into account regional or subregional specificities.
Descriptor 7	□ D7C1 – Secondary:
	Spatial extent and distribution of permanent alteration of hydrographical conditions (e.g. changes in wave action, currents, salinity, temperature) to the seabed and water column, associated in particular with physical loss(1) of the natural seabed.
	□ D7C2 – Secondary:
	Spatial extent of each benthic habitat type adversely affected (physical and hydrographical characteristics and associated biological communities) due to permanent alteration of hydrographical conditions.
Descriptor 8	□ D8C1 – Primary:
	Within coastal and territorial waters, the concentrations of contaminants do not exceed the following threshold values:
	(a) for contaminants set out under point 1(a) of criteria elements, the values set in accordance with Directive 2000/60/EC;
	(b) when contaminants under point (a) are measured in a matrix for which no value is set under Directive 2000/60/EC, the concentration of those contaminants in that matrix established by Member States through regional or subregional cooperation;
	(c) for additional contaminants selected under point 1(b) of criteria elements, the concentrations for a specified matrix (water, sediment or biota) which may give rise to pollution effects. Member States shall establish these concentrations through regional or subregional cooperation, considering their application within and beyond coastal and territorial waters.
	Beyond territorial waters, the concentrations of contaminants do not exceed the following threshold values:

	(a) for contaminants selected under point 2(a) of criteria elements, the values as applicable within coastal and territorial waters;
	(b) for contaminants selected under point 2(b) of criteria elements, the concentrations for a specified matrix (water, sediment or biota) which may give rise to pollution effects. Member States shall establish these concentrations through regional or subregional cooperation.
	☑ D8C2 – Secondary:
	The health of species and the condition of habitats (such as their species composition and relative abundance at locations of chronic pollution) are not adversely affected due to contaminants including cumulative and synergetic effects.
	Member States shall establish those adverse effects and their threshold values through regional or subregional cooperation.
	□ D8C3 – Primary:
	The spatial extent and duration of significant acute pollution events are minimised.
	\square D8C4 – Secondary (to be used when a significant acute pollution event has occurred):
	The adverse effects of significant acute pollution events on the health of species and on the condition of habitats (such as their species composition and relative abundance) are minimised and, where possible, eliminated.
Descriptor 9	□ D9C1 – Primary:
	The level of contaminants in edible tissues (muscle, liver, roe, flesh or other soft parts, as appropriate) of seafood (including fish, crustaceans, molluscs, echinoderms, seaweed and other marine plants) caught or harvested in the wild (excluding fin-fish from mariculture) does not exceed:
	(a) for contaminants listed in Regulation (EC) No 1881/2006, the maximum levels laid down in that Regulation, which are the threshold values for the purposes of this Decision;
	(b) for additional contaminants, not listed in Regulation (EC) No 1881/2006, threshold values, which Member States shall establish through regional or subregional cooperation.

Descriptor 10	□ D10C1 – Primary:
	The composition, amount and spatial distribution of litter on the coastline, in the surface layer of the water column, and on the seabed, are at levels that do not cause harm to the coastal and marine environment.
	Member States shall establish threshold values for these levels through cooperation at Union level, taking into account regional or subregional specificities.
	□ D10C2 — Primary:
	The composition, amount and spatial distribution of micro-litter on the coastline, in the surface layer of the water column, and in seabed sediment, are at levels that do not cause harm to the coastal and marine environment.
	Member States shall establish threshold values for these levels through cooperation at Union level, taking into account regional or subregional specificities.
	□ D10C3 — Secondary:
	The amount of litter and micro-litter ingested by marine animals is at a level that does not adversely affect the health of the species concerned. Member States shall establish threshold values for these levels through regional or subregional cooperation.
	□ D10C4 — Secondary:
	The number of individuals of each species which are adversely affected due to litter, such as by entanglement, other types of injury or mortality, or health effects. Member States shall establish threshold values for the adverse effects of litter, through regional or subregional cooperation.
Descriptor 11	□ D11C1 – Primary:
	The spatial distribution, temporal extent, and levels of anthropogenic impulsive sound sources do not exceed levels that adversely affect populations of marine animals.
	Member States shall establish threshold values for these levels through cooperation at Union level, taking into account regional or subregional specificities.
	□ D11C2 – Primary:
	The spatial distribution, temporal extent and levels of anthropogenic continuous low-frequency sound do not exceed levels that adversely affect populations of marine animals.
	Member States shall establish threshold values for these levels through cooperation at Union level, taking into account regional or subregional specificities.

d. References

Make a list of cited references and literature for further supportive information.

Core indicator Producvity of white-tailed Eagle.