



Climate Change in the Baltic Sea

2021 Fact Sheet

Climate change



BSEP n°180

**EMBARGOED**

For preview purposes only. Do not share.

Published by:
Helsinki Commission – HELCOM
Katajanokanlaituri 6 B
00160 Helsinki, Finland

www.helcom.fi

This document is part of the flagship publication series of HELCOM, the Baltic Sea Environment Proceedings (BSEP) that have been running since the entry into force of the first Helsinki Convention in 1980. Although this document has been approved for publication by the members of the Helsinki Commission, views expressed in this publication are the authors’ own and might vary from those of the Helsinki Commission or its members. Any maps that are featured in this publication are intended for illustration purposes only and do not necessarily designate the exact boundaries of sovereign states and entities.

The development of this publication was steered by the Joint HELCOM/Baltic Earth Expert Network on Climate Change (EN CLIME).

For bibliographic purposes this document should be cited as:
“Climate Change in the Baltic Sea. 2021 Fact Sheet. Baltic Sea Environment Proceedings n°180. HELCOM/Baltic Earth 2021.”

© 2021 Baltic Marine Environment Protection Commission
(Helsinki Commission – HELCOM)

All rights reserved. Information included in this publication or extracts thereof, with the exception of images and graphic elements that are not HELCOM’s or Baltic Earth’s own and identified as such, may be reproduced without prior consent on the condition that the complete reference of the publication is given as stated above.

Contributors:
Markus Ahola, Lena Bergström, Mats Blomqvist, Dieter Boedeker, Florian Börgel, Ida Carlén, Thomas Carlund, Jacob Carstensen, Jesper Philip Aagaard Christensen, Martyn Futter, Elie Gaget, Matthias Gröger, Volker Dierschke, Christian Dieterich, Morten Frederiksen, Anders Galatius, Bo Gustafsson, Claudia Frauen, Jannica Haldin, Antti Halkka, Christina Halling, Jürgen Holfort, Kari Hyytiäinen, Klaus Jürgens, Mart Jüssi, Laura Kaikkonen, Meri Kallasvuo, Markus Kankainen, Martin Karlsson, Dorte Krause-Jensen, Anders Kiessling, Erik Kjellström, Antanas Kontautas, Karol Kuliński, Sanna Kuningas, Jukka Käyhkö, Petra Kääriä, Janika Laht, Ari Laine, Gesine Lange Antti Lappalainen, Terhi Laurila, Maiju Lehtiniemi, Knut-Olof Lerche, Urmas Lips, Georg Martin, Michelle McCrackin, H.E. Markus Meier, Noora Mustamäki, Bärbel Müller-Karulis, Rahmat Naddafi, Lauri Niskanen, Antonia Nyström Sandman, Jens Olsson, Diego Pavón-Jordán, Jonas Pålsson, Mika Rantanen, Artūras Razinkovas-Baziukas, Berit Recklebe, Gregor Rehder, Jan H. Reißman, Martin Reutgård, Stuart Ross, Marta Ruiz, Anna Rutgersson, Jarkko Saarinen, Lauri Saks, Oleg Savchuk, Mikhail Sofiev, Katarzyna Spich, Jani Särkkä, Markku Viitasalo, Jouni Vilma, Joonas Virtasalo, Isa Wallin, Ralf Weisse, Johan Wikner, Wenyan Zhang, Eduardo Zorita, Örjan Östman

Acknowledgements:
Maris Arro, Paweł Banaś, Imre Banyasz, Edyta Białowąs, Penina Blankett, Matthias Brenner, Laura Briekmane, Michele Casini, Johan Dannewitz, Michael Dähne, Jacques Delsalle, Rune Dietz, Łukasz Dziemian, Anthony David Fox, Oksana Glibko, Magnus Huss, Norbert Häubner, Birgit Hünicke, Tamara Jadczyzyn, Eglė Jakubavičiūtė, Dominika Juskowska, Magdalena Kamińska, Agnes Karlsson, Marcin Kawka, Ilga Kokorite, Harri Kuosa, Joakim Lagner, Kristina Lehnert, Adam Lejk, Peter Löwe, Katarina Magnusson, Sofia Malmsten, Piotr Margorński, Johanna Mattila, Iwona Pawliczka, Liisa Pietola, Maris Plikss, Inese Pommere-Bramane, Konrad Prandecki, Marcus Reckermann, Daria Ryabchuk, Hanna Sjölund, Piotr Skowron, Miriam Sollich, Henrik Svedäng, Agata Święcka, Morten Tange, Maciej Tomczak, Lasse Tor, Emma Undeman, Jacek Walczak, Tamara, Zalewska, Sergey Zhuravlev, Marek Zieliński

We also thank the four anonymous and independent reviewers for their detailed and constructive comments that considerably helped to improve the fact sheet.

Further acknowledgements go to the FutureMARES EU project (<https://www.futuremares.eu/>)

Editors: Jannica Haldin, Petra Kääriä, H.E. Markus Meier, Jonas Pålsson

Layout: Dominik Littfass

ISSN: 0357-2994



Contents

The Baltic: A sea of change	6	Indirect parameters: Ecosystem	36
Baltic Sea Expert Network on Climate Change – EN CLIME	7	Oxygen	38
Impact map	7	Microbial community and processes	39
Confidence assessment	7	Benthic habitats	40
Parameters covered	8	Coastal and migratory fish	41
Peer review of key messages	8	Pelagic and demersal fish	42
Climate change & climate mitigation	8	Waterbirds	43
Connections between parameters	9	Marine mammals	44
Climate future of the Baltic Sea	10	Non-indigenous species	45
Projections under a medium climate scenario	10	Marine protected areas	46
		Nutrient concentrations and eutrophication	47
		Ecosystem function	48
Direct parameters	18	Indirect parameters: Human use	50
Air temperature	20	Offshore wind farms	52
Water temperature	21	Coastal protection	53
Large scale atmospheric circulation	22	Shipping	54
Sea ice	23	Tourism	55
Solar radiation	24	Fisheries	56
Salinity and saltwater inflows	25	Aquaculture	57
Stratification	26	Blue Carbon storage capacity	58
Precipitation	27	Marine and coastal ecosystem services	59
River run-off	28		
Carbonate chemistry	29		
Riverine nutrient loads and atmospheric deposition	30		
Sea level	31		
Wind	32	Glossary	62
Waves	33	Policy linkages	65
Sediment transportation	34	References	66



Climate change effects on the Baltic Sea environment are manifold. It is for example expected that water temperature and sea level will rise, and sea ice cover will decrease. This will affect ecosystems and biota; for example, range shifts are expected for a number of marine species, benthic productivity will decrease, and breeding success of ringed seals will be reduced. The impacts will hence affect the overall ecosystem function and also extend to human uses of the sea; trawling will follow the fish towards southern areas, aquaculture will likely face a shift towards species diversification, and the value of most ecosystem services is expected to change — to name a few.

This Climate Change Fact Sheet provides the latest scientific knowledge on how climate change is currently affecting the Baltic Sea and how it is expected to develop in the foreseeable future. It is aimed at guiding policy makers to take climate change into account, but also to the general public. Updated Baltic Sea Climate Change Fact Sheets are expected to be published approximately every seven years.



The Baltic: A sea of change

Introduction



Climate change impacts are evident in the Baltic Sea: water temperature is rising, ice extent is decreasing, and annual mean precipitation is increasing over the northern part of the region. All these changes affect the nature of the sea, its ecosystems, and ecosystem services, as well as the human activities depending on the sea. For example, many wintering birds have shifted their wintering range northwards, the numbers of warm water fish species (such as sticklebacks) are increasing, the risk of infection of human-pathogenic *Vibrio* spp. has increased through surface water warming, and trawl fishing now begins earlier in the year.

The Baltic Sea is facing a complex system of effects and feedbacks between climatic and non-climatic factors. Multiple environmental pressures affect the ecosystem, and climate change adds further cumulative pressures to the existing anthropogenic ones. These various climate change effects are not straightforward to understand and are difficult to distinguish from certain human pressures. Climate and other human-induced pressures vary significantly between different regions in the Baltic Sea, making it impossible to find simple management solutions that can work everywhere. In order to mitigate these negative effects, policymakers need to be aware of these differences and utilise an adaptive management approach based on the best available science.

This Fact Sheet provides the latest scientific knowledge on how climate change is affecting the Baltic Sea in a concise format. It is the first of a series of successive Baltic Sea Climate Change Fact Sheets aiming to track advances in the understanding of how climate change impacts the state of marine systems, drawing on the best available science for the region.

How climate change already has and is expected to impact the Baltic Sea is described through 34 parameters that have been identified by EN CLIME as being of relevance for science and management. These parameters constitute physiochemical parameters that are directly affected by climate change, referred to as direct parameters (page 18), as well as ecosystem and human use parameters that are indirectly affected, referred to as indirect parameters (page 36). The full list of parameters is shown in Table 1 (page 8).

The first part of this report provides summary information of climate change impacts on each parameter (pages 12-17), as well as an impact map showing the projected regional changes for a selected suite of parameters under the RCP4.5 climate scenario across the Baltic Sea. The second part of the report (pages 18-59) gives a more detailed, yet concise, overview of climate change impacts on each parameter - described as key messages.



Baltic Sea Expert Network on Climate Change – EN CLIME

In 2018, the Baltic Sea Environment Protection Commission (HELCOM) and Baltic Earth formed a joint Expert Network on Climate Change in the Baltic Sea region (EN CLIME). This Expert Network involves more than 110 scientists from around the Baltic Sea. The purpose of the network is to function as a coordinating framework and a platform to harness the expertise of leading scientists on both direct and indirect effects of climate change on the Baltic Sea environment and ecosystems and make this expertise available to and open up for closer dialogue with policy makers.

Impact map

The impact map (pages 10-11) depicts projected regional changes for some of the most relevant parameters in a particular subbasin of the Baltic Sea under the RCP4.5 scenario. While there is

also important information on the other parameters, there was a need to reduce the total 34 parameters to the presented parameters to make the map more legible. The presented parameters have 1) direct societal relevance/experience and/or relevance for other parameters, 2) medium to high confidence of the changes relative to the noise and model/expert judgement uncertainty under the RCP4.5 scenario, and 3) a hotspot sub-region in the Baltic with medium to high confidence of patterns of the regional changes.

Confidence assessment

The level of confidence of statements is shown with confidence assessments using the scale low-medium-high (Figure 1). The authors were asked to consider both the level of consensus and the amount of evidence when defining an overall confidence of a statement and to select the overall confidence by using the precautionary principle (e.g., in case the level of consensus is low and the amount of evidence medium, the overall confidence is low).




Figure 1. The overall confidence is resulting from the confidence assessment of the agreement/consensus on and evidence of the assessed data.



Parameters covered

The 34 parameters have been categorized into six different categories: Energy cycle, Water cycle, Carbon and Nutrient cycles, Sea level and wind, Biota and ecosystems, Human activities, and Services.

The following parameters were considered as important to include, but due to the lack of lead authors, they were not included in this version of the fact sheet:

- Pelagic habitats (incl. phytoplankton and zooplankton community structure, spring blooms, functional traits etc.)
- Harmful algal blooms (HABs)
- Pollution and hazardous substances
- Ecotoxicology
- Human health
- Pathogens

Table 1. Full list of EN CLIME parameters. The asterisk (*) indicates those parameters that include information on extreme events.

Direct parameters	Categorization
Air temperature*	Energy cycle
Water temperature*	Energy cycle
Large scale atmospheric circulation	Energy cycle
Sea ice*	Energy cycle
Solar radiation	Energy cycle
Salinity and saltwater inflows*	Water cycle
Stratification	Water cycle
Precipitation*	Water cycle
River run-off*	Water cycle
Carbonate chemistry	Carbon and nutrient cycles
Riverine nutrient loads and atmospheric deposition	Carbon and nutrient cycles
Sea level*	Sea level and wind
Wind*	Sea level and wind
Waves*	Sea level and wind
Sediment transportation*	Sea level and wind

Indirect parameters	Categorization
Oxygen	Carbon and nutrient cycles
Microbial community and processes	Biota and ecosystems
Benthic habitats	Biota and ecosystems
Coastal and migratory fish	Biota and ecosystems
Pelagic and demersal fish	Biota and ecosystems
Waterbirds	Biota and ecosystems
Marine mammals	Biota and ecosystems
Non-indigenous species	Biota and ecosystems
Marine protected areas (MPAs)	Biota and ecosystems
Nutrient concentrations and eutrophication	Biota and ecosystems
Ecosystem function	Biota and ecosystems
Offshore wind farms	Human activities
Coastal protection	Human activities
Shipping	Human activities
Tourism	Human activities
Fisheries	Human activities
Aquaculture	Human activities
Blue carbon storage capacity	Services
Marine and coastal ecosystem services	Services

Peer review of key messages

The key messages have been peer reviewed and improved in a two-step process. The first review round was carried out by six external scientists and the second round was carried out by the Co-chairs and HELCOM Secretariat.

Climate change & climate mitigation

The global climate is changing, and this is due to human influence in the form of greenhouse gas emissions (GHG) from fossil fuel use and land use change. The current changes in the climate systems have already had widespread impacts on human and natural systems.

According to the Intergovernmental Panel on Climate Change (IPCC)¹, human activities are estimated to have caused approximately 1.1°C of global warming above pre-industrial levels and global warming will continue during the coming decades. The pace and magnitude of warming will depend on how global greenhouse gas emissions evolve.

In order to reduce the impacts of rising temperature on Earth, all global policy actions aiming at the mitigation of greenhouse gas emissions are highly relevant. With the help of climate models and various emission scenarios, projections of global and regional climates have been performed to support policymaking such as the Paris Agreement.

Different Representative Concentration Pathways (RCPs) are used to describe different climate futures depending on the greenhouse gas emissions in the coming years. The RCPs indicate a possible range of radiative forcing (difference between solar energy absorbed by the Earth and radiated back to space) in the year 2100. The RCPs include a “mitigation” scenario which aims to keep global warming below 2°C above pre-industrial temperatures (RCP2.6) and a high emissions “worst case” scenario (RCP8.5), that corresponds to a future without climate mitigation. One intermediate scenario is the RCP4.5 which is used in the impact map of this Fact Sheet and likely results in global mean temperature rise between 2-3°C by 2100.

When the IPCC Assessments have been referred to in this Climate Change Fact Sheet, the information is based on the IPCC Assessment Report 5 (2013)², the Special Report on the Ocean and Cryosphere in Changing Climate (2019)³ and earlier publications, as the most recent Assessment Report 6 had not yet been published by the time this Fact Sheet was produced. Information about regional climate change is based upon the BACC Reports (BALTEX and Baltic Earth Assessments of Climate Change for the Baltic Sea Basin, BACC Author Team, 2008⁴; BACC II Author Team, 2015⁵; see www.baltic.earth).

Connections between parameters

Links between the different parameters have been shown in Figure 2, depicting complex interconnections between the different abiotic, ecosystem, and human dimension parameters. The colour of each arrow comes from the parameter it originates from.




Figure 2. Linkages between the different parameters that were used in the assessment of the effects of climate change in the Baltic Sea.



Climate future of the Baltic Sea

Projections under a medium climate scenario (such as RCP4.5)

The impact map depicts projected regional changes for some of the most relevant parameters in a particular subbasin of the Baltic Sea under the RCP4.5 scenario. While there is also important information on the other parameters, there was a need to reduce the total 34 parameters to the presented parameters to make the map more legible. The presented parameters have 1) direct societal relevance/experience and/or relevance for other parameters, 2) medium to high confidence of the changes relative to the noise and model/expert judgement uncertainty under the RCP4.5 scenario, and 3) a hotspot sub-region in the Baltic with medium to high confidence of patterns of the regional changes.



Bothnian Sea

Sea surface temperature would rise everywhere in the Baltic and in all seasons. Most pronounced would be summer warming in the Bothnian Bay and Bothnian Sea. Winter precipitation including high-intensity extremes would increase. Increased freshwater discharge would bring more dissolved organic carbon to the sea, affecting benthic habitats by decreasing pelagic primary production and phytoplankton sedimentation. In the Bothnian Sea, Gulf of Finland and Gulf of Riga, the decline in sea ice cover would be largest. Waves would be higher and shipping might increase if the ice cover is reduced. Food accessibility for migratory water birds would improve causing a northward shift of breeding and wintering areas towards ice free coastal areas. In the Archipelago Sea, ringed seal populations might decrease.



Baltic Sea entrance area

Sea surface temperature would rise. Mean sea level is projected to rise relative to the land, and higher storm surges would occur. Higher atmospheric pCO₂ would cause increased acidification.



Bothnian Bay

Air temperature is projected to rise, most pronounced in the northern Baltic Sea region during winter. Sea surface temperature would rise and sea ice thickness and the length of the ice season would decrease. Winter precipitation including high-intensity extremes would increase. Increased freshwater discharge would bring more dissolved organic carbon to the sea, affecting benthic habitats by decreasing pelagic primary production and phytoplankton sedimentation. Land is rising faster than the projected sea level and the mean sea level would sink relative to land.



Baltic Proper

Sea surface temperature would rise. If BSAP measures on nutrient loads were to be implemented, phosphorus concentrations and algal blooms would decrease and oxygen conditions of the deep water would improve. Without load reductions, only minor changes in nutrient concentrations are expected. The combined effects of warming and planned nutrient reductions will eventually lead to less carbon reaching the seafloor, reducing benthic animal biomass. In shallow archipelago waters, the fates of benthic animal and plant populations depend on local variations in biogeochemistry and primary productivity. In the southern Baltic, mean sea level would rise relative to the land, and higher storm surges would occur. Sediment transports would change.



Gulf of Finland

Sea surface temperature would rise and sea ice cover, ice thickness and the length of the ice season would decrease, affecting ringed seal breeding and probably causing a decline of the populations in the eastern Gulf of Finland. Likewise breeding and wintering areas of migratory water birds would be affected. Wave heights would increase and the potential for shipping would increase if the ice cover is reduced, but shipping intensity is more dependent on market development than climate change. In the eastern Gulf of Finland, mean sea level would rise relative to the land, and higher storm surges would occur.



Gulf of Riga

Sea surface temperature would rise and sea ice cover would decline, affecting ringed seal populations in the northern Gulf of Riga. Likewise, breeding and wintering areas of migratory water birds would be affected. In the southern Gulf of Riga, mean sea level would rise relative to the land, and higher storm surges would occur.

Assessment sub-basins

1. Bothnian Bay (Bothnian Bay and the Quark)
2. Bothnian Sea (Bothnian Sea and Åland Sea)
3. Gulf of Finland
4. Gulf of Riga
5. Baltic Proper (Northern Baltic Proper, Western Gotland Basin, Eastern Gotland Basin, Bornholm basin and Gdansk Basin)
6. Entrance area (Kattegat, Great Belt, the Sound, Kiel Bay, Bay of Mecklenburg and Arkona Basin)



Direct parameters

Physiochemical parameters directly affected by climate change


Categories

- Energy cycle
- Water cycle
- Carbon and nutrient cycle
- Sea level and wind
- Biota and ecosystems
- Human activities
- Services



Air temperature

Air temperature shows the clearest response to increased greenhouse gas emissions. A significant air temperature increase in the Baltic Sea region has been observed during the last century, larger than the global trend, and this increase is expected to continue. In addition, warm extremes are projected to become more pronounced.



Water temperature

The marginal seas around the globe have become warmer during the past 40 years. The sea surface temperature of the Baltic Sea has warmed more than the average for the global ocean and will continue to warm.



Large scale atmospheric circulation

The climate of the Baltic Sea region is strongly influenced by the large-scale atmospheric circulation, in particular the North Atlantic Oscillation, atmospheric blocking patterns, and Atlantic Multidecadal Oscillation are dominant patterns. As the response of these atmospheric circulation patterns to climate change differs among models, future projections are very uncertain.



Sea ice

Sea ice forms every winter, the most important factor being air temperature, but also wind, snow cover and ocean currents. Over the past 100 years, the winters have become milder, the ice season shorter and the maximum ice extent decreased. This development is expected to continue in a changing climate.



Solar radiation

The solar radiation is the engine of the climate system. The solar radiation reaching the surface strongly depends on cloudiness, and also on aerosols. There is an indication of decline in cloudiness during the past decades. For the future, there is very limited knowledge.



Salinity and saltwater inflows

Salinity affects the dynamics of ocean currents and ecosystem functioning. Salinity decreases gradually from Kattegat to the Bothnian Bay. Inflows from the North Sea sporadically renew the deep water with saline, oxygen rich water. No statistically significant trends in salinity have been found, and uncertainties of future projections are high.



Stratification

Seawater is layered (stratified) according to its density, a property governed by temperature and salinity. Over the last 40 years, stratification in the Baltic Sea has become stronger. This trend may continue in the future and cause harm to the marine ecosystem by decreasing the mixing between surface waters and deep waters.




Precipitation

Precipitation depends on the circulation, the amount of water vapor in the air, the temperature and the land-sea contrast. Annual mean precipitation has significantly increased over the northern Baltic Sea lately while in the south, changes are small – a trend that may continue in the future.



River run-off

Runoff describes the amount flowing water entering the sea. The total annual river runoff has not changed over the last 500 years, but a significant increase in winter river discharge and a decrease in spring floods has been observed lately. The total runoff to the Baltic Sea may increase with warming temperatures.




Carbonate chemistry

The carbonate system regulates seawater pH. The amount of CO₂ in the Baltic Sea surface water changes seasonally mostly due to biologically driven processes (photosynthesis and respiration), which induces seawater pH oscillations. In the long term, atmospheric CO₂ increase will raise seawater CO₂ concentration and cause pH decrease.



Riverine nutrient loads and atmospheric deposition

External nutrient inputs from land and atmosphere are the major long-term drivers of the Baltic Sea eutrophication. Substantial reductions in nutrient loads have occurred since the 1980s, however, no large-scale effects on ecosystem status can be detected yet. In the future, land-based nutrient management will have greater effect on loads than greenhouse gases emissions.



Sea level

Baltic Sea mean level responds to global sea level rise and regional land uplift and varies with season and climate. Baltic sea level is rising and will continue to rise. Storm surges are sensitive to changes in atmospheric circulation and future changes are uncertain.



Wind

The wind climate and storms over the Baltic Sea are determined by the large-scale atmospheric circulation. Storms are typically more frequent and stronger during winter. The large natural variability over the Baltic Sea masks possible past and future trends.



Waves

The wave climate in the Baltic Sea strongly depends on the wind field and shows large long-term variability. Significant trends in the wave height have not been detected. For northern and eastern parts of the Baltic a slight increase is significant and extreme wave height is projected.



Sediment transportation

Near shore sediment transport is triggered by waves and wind and leads to erosion and accumulation of sediments. Sandy beaches along the southern and eastern coastlines of the Baltic Sea are especially vulnerable and rising sea level will increase sediment transport.




Indirect parameters: Ecosystem

Ecosystem parameters indirectly affected by climate change

Categories

- Energy cycle
- Water cycle
- Carbon and nutrient cycle
- Sea level and wind
- Biota and ecosystems
- Human activities
- Services



Oxygen

Oxygen concentration is controlled by physical transport and remineralization of organic matter. Bottom water oxygen deficiency observed in a vast area of the Baltic Sea is a consequence of water column stratification and eutrophication. Thus, future oxygen availability will depend on nutrient loads, while projected warming may reinforce eutrophication.



Microbial community and processes

Bacterially-mediated processes as well as the occurrence of pathogenic Vibrios are expected to increase with current environmental changes. However, only small changes in bacterial biomass and growth were detected during the past decades. The potential genetic adaptation to climate change and lack of proper models including bacterioplankton make predictions for the future uncertain.



Benthic habitats

In the Baltic Sea, many benthic species exist at the edge of their distribution, and even small fluctuations in temperature and salinity can impact their abundance, biomass, and spatial distribution. In concurrence with trophic cascades and eutrophication, climate change might lead to major changes in biodiversity and ecosystem functions of benthic habitats.



Coastal and migratory fish

Coastal and migratory fish respond to changes in temperature, ice-cover, salinity and river-discharge. Spring and summer-spawning species (e.g. perch, cyprinids, pike) will benefit from increasing temperatures, whereas autumn-spawning (e.g. salmonids) may be disfavoured. Future actions must consider eutrophication, fishing, food-web interactions and habitat exploitation, for migratory fish also in rivers.



Pelagic and demersal fish

Fish of marine origin mainly respond to changes in temperature, salinity, water stratification and circulation influencing oxygen conditions. Actions to reduce eutrophication, anoxic conditions, and fishing, while considering food-web interactions will be important.



Waterbirds

Most obvious effects of climate change on Baltic waterbirds are range shifts in winter (migratory birds stay closer to breeding areas). Food supply (fish, bivalves) and breeding conditions are influenced in various ways.



Marine mammals

Grey and particularly ringed seal breeding success will be reduced by decreased sea ice quality and quantity. Harbour and grey seal southern Baltic distribution will be reduced by flooding of haul-outs. Changed temperature, stratification, prey distribution, quality and quantity will affect marine mammals, but aggregate effects are unpredictable.



Non-indigenous species

While shipping is the main driver of new non-indigenous species (NIS) introductions, climate change related changes in abiotic environment may support their establishment and range expansion. Increasing water temperature may favour species of warm water origin, and potential salinity decrease will benefit NIS of freshwater origin, impacting likely estuarine ecosystems.



Marine protected areas

Climate change may impact Marine protected areas (MPAs) via changes in abiotic environment causing diverse changes in ecosystem structure and functions, thus altering MPAs' conservation values. Changes are expected first in seal and water bird populations, followed by potential large-scale changes in benthic habitats if possible salinity decrease starts affecting the distribution of key species.



Nutrient concentrations and eutrophication

Nitrogen and phosphorus pools are controlled by loads from land and atmosphere and influenced by oxygen-sensitive biogeochemical processes. Future load changes will have a stronger influence on nutrients than climate change, even though projected warming will increase nutrient cycling and reduce bottom water oxygenation.



Ecosystem function

Baltic Sea ecosystems provide an array of functions related to nutrient- and carbon circulation, biomass production and regulation. Climate impacts ecosystem functions via temperature, water circulation, salinity, river-discharges, and solar-radiation. In the future, increased productivity, stronger impact of nutrients and reduced influence of predators will influence Baltic Sea ecosystem functioning.



Indirect parameters: Human use

Human use parameters indirectly affected by climate change

Categories

- Energy cycle
- Water cycle
- Carbon and nutrient cycle
- Sea level and wind
- Biota and ecosystems
- Human activities
- Services



Offshore wind farms

Wind farms are the most significant offshore structures in the Baltic Sea. Declining ice cover and rising sea level can affect offshore wind farms. Offshore wind farms affect many oceanographic processes and have a substantial effect on the structural and functional biodiversity of the benthic system. They account for 10 % of European offshore wind energy and are crucial for reaching new energy and climate targets.



Coastal protection

The shorelines of the Baltic Sea vary from bedrock-dominated stable coasts in the north to soft, sandy shores in the south, where periods of storminess cause coastal erosion. Declining ice cover and rising sea level increase the potential for coastal erosion.



Shipping

Shipping is primarily affected by ice and extreme weather, and the potential for shipping will increase if the ice cover is reduced. However, shipping intensity is more dependent on market development than climate change. Regulatory measures to decarbonise shipping are increasing and driving important adaptations across the industry.




Tourism

Climate change shapes the spatial and temporal distribution of tourism resources within and between regions. The future competitiveness of coastal and maritime tourism in the Baltic Sea region will be conditional to the adaptive capacity of the sector to climate change, changing consumer values, natural and human-made hazards, and economic and political disturbances.



Fisheries

Most notable impacts to fisheries will take place in the northern Baltic Sea. Trawl fishing season will be extended, trawling areas shifted towards the south and shallower areas, target species compositions shifted towards species preferring warmer waters, and winter-time coastal fishing decreased due to diminishing ice-cover.



Aquaculture

Baltic Sea aquaculture is dominated by open-cage rainbow trout farms with low climate impact. Cultivation of blue catch-crops, including plants and invertebrates, is increasing. Warmer conditions will promote offshore locations and species diversification. Industrial scale, land-based aquaculture farms are unlikely in rural parts due to their external resource- and infrastructure dependents.



Blue carbon storage capacity

Blue Carbon (BC) refers to the carbon marine organisms sequester in oceanic carbon sinks. Climate change effects on BC habitats, such as effects on carbon sink capacity and changed amount of macrophytes, are expected to increase in the future, with associated effects on climate change mitigation.



Marine and coastal ecosystem services

Ecosystem services in the north-most and coastal semi-enclosed areas with lower salinities will be affected first. Most ecosystem services are expected to decline, while only the cultural services, connected to recreation, could gain from the climatic changes due to longer summers and higher air and water temperatures. Other anthropogenic pressures could both offset positive and strengthen negative trends in ecosystem services supply.

Direct parameters





Air temperature

Linked parameters:
Water temperature, Large-scale atmospheric circulation, Sea ice, Solar radiation, River runoff, Riverine nutrient loads and atmospheric deposition, Waterbirds, Tourism

Links to main policies:
UN Sustainable Development Goal 13
EU Biodiversity Strategy



Description

Since air temperature shows the clearest response to the increased green-house effect, the near-surface mean air temperature is often used as the main indicator of a changing climate globally and regionally. Changes in temperature extremes may influence biological and human activities much more than changes in average temperature.



What is already happening?

Mean change: An increase in air temperature is seen during the last century, with an accelerated increase during the last decades¹⁻³. Annual mean temperature trends during 1876–2018 indicate that air temperature has increased more in the Baltic Sea region than globally. The increase is accompanied by large multi-decadal variations, in particular during winter, but the warming is seen for all seasons and is largest during spring.

Extremes: During the recent decade, record breaking heat waves have hit the region, with an increasing trend of warm spell duration^{4,5}. A decrease is seen in the length of the frost season and in the number of frost days.



What can be expected?

Mean change: Air temperatures are projected to increase more in the Baltic Sea region than the global mean. Regional scenarios project an annual mean near-surface temperature increase over the Baltic Sea of 1.4°C (1.2–1.9°C, RCP2.6) to 3.9°C (3.1–4.8°C, RCP8.5)* by the end of this century⁶, compared to 1976–2005. The air temperature increase is larger in the North than in the South because of the snow and sea-ice cover decline enhancing absorption of sunlight by soil and water².

Extremes: Larger warming is expected for cold extremes than for the mean winter temperature⁷. In summer, warm extremes are projected to become more pronounced. Warm extremes presently with a 20-year return probability will occur around once every five years in Scandinavia by 2071–2100⁸.



Knowledge gaps


The variability in temperature and temperature extremes are to a large extent determined by the large-scale circulation patterns. There is limited knowledge primarily concerning changes in large-scale atmospheric circulation patterns in a changing climate because of model differences.



Policy relevance

Higher temperatures trigger marine heatwaves, and will have direct and indirect effects on habitats, species, and populations in terrestrial and aquatic ecosystems. Higher mean temperatures and increased number of heatwaves will increase the risks of droughts and forest fires. There is a need for better urban planning, for example adapting building standards for warmer climate and increasing urban green areas. Areas such as Gotland have increased the capacity of their desalination plants, to ensure sufficient drinking water during droughts. Further measures to better manage heat and drinking water need to be implemented.

*) Changes in mean, 5th and 95th percentiles indicating the spread in an ensemble of 9 climate models.



Water temperature

Linked parameters:
Air temperature, Sea ice, Solar radiation, Stratification, Carbonate chemistry, Oxygen, Microbial community and processes, Benthic habitats, Coastal and migratory fish, Pelagic and demersal fish, Waterbirds, Non-indigenous species, Nutrient concentrations and eutrophication, Ecosystem function, Tourism, Aquaculture, Blue carbon storage capacity

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goals 13 and 14
UN Convention on Biological Diversity
EU Green Deal
EU Marine Strategy Framework Directive (MSFD)
EU Water Framework Directive (WFD)
EU Maritime Spatial Planning Directive (MSP)
EU Habitats Directive
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Biodiversity Strategy



Description

As air temperature increases, also water temperature rises¹. Starting at the surface, the heat spreads downward through different processes and may warm up even the deep water of the Baltic Sea. The ocean plays an important role for the climate because by far the largest amount of the heat from global warming is stored in the oceans. Due to their huge heat capacity, oceans respond slowly, and moderate temperature increases in the atmosphere. Oceans are also important in providing moisture to the atmosphere, the more the warmer the water is.



What is already happening?

Mean change: Marginal seas around the globe have warmed faster than the global ocean², and the Baltic Sea has warmed the most of all marginal seas². Average surface-water temperature increased by +0.59°C/decade for 1990–2018³ and between +0.03 and +0.06°C/decade for 1856–2005 in northeastern and southwestern areas, respectively⁴.

Extremes: With reference to 2020, the summer of 2018 was the warmest on instrumental record in Europe, and also the warmest summer in the past 30 years in the southern half of the Baltic Sea⁵, with surface-water temperatures 4–5°C above the 1990–2018 long-term mean. The heat wave has also been recorded in bottom temperatures⁶.



What can be expected?

Mean change: Global ocean temperatures are rising at accelerating rates^{7,8}. Scenario simulations for the Baltic Sea project a sea surface temperature increase of 1.1°C (0.8–1.6°C, RCP2.6) to 3.2°C (2.5–4.1°C, RCP8.5)* by the end of this century compared to 1976–2005^{9–12}. In all scenarios, sea surface temperature changes at the end of the century significantly exceed natural variability.

Extremes: The RCP4.5 and RCP8.5 scenarios project more tropical nights over the Baltic Sea, increasing the risk of record-breaking water temperatures¹³.



Knowledge gaps

For the projection of water temperatures in the Baltic Sea, regional climate models are needed. However, the effect of aerosols in regional climate models has not been investigated. More knowledge on natural variability of Baltic Sea temperature and its connection to large-scale patterns of climate variability is needed. The occurrence of marine heatwaves is projected to increase. However, their potential to affect the ecosystem in the Baltic Sea is not well known.



Policy relevance

Water temperature has profound effects on the marine ecosystem. Climate change mitigation is the only way to counteract temperature increase. The best adaptation response available is to reduce environmental pressures to the Baltic Sea, thus building climate change resilience. The protection of marine areas where the temperature increase is expected to be lower, so-called climate refuges, focuses on areas where climate change impacts are not contributing to multiple stressors^{14,15}. These could become a last outpost for species affected by climate change.

*) Changes in mean, 5th and 95th percentiles indicating the spread in an ensemble of 9 climate models.



Large scale atmospheric circulation

Linked parameters:
Air temperature, Solar radiation, Precipitation, Wind

Authors
Claudia Frauen, Leibniz Institute for Baltic Sea Research Warnemünde (IOW), Germany
Anna Rutgersson, Uppsala University, Sweden
Florian Börgel, Leibniz Institute for Baltic Sea Research Warnemünde (IOW), Germany
H.E. Markus Meier, Leibniz Institute for Baltic Sea Research Warnemünde (IOW), Germany
and Swedish Meteorological and Hydrological Institute (SMHI), Sweden

Energy cycle

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goal 13
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Green Deal
EU Biodiversity Strategy



Description

The climate of the Baltic Sea region is influenced by the large-scale atmospheric circulation. The variability of the circulation can be decomposed into various dedicated modes of variability:

1. The North Atlantic Oscillation (NAO) describes the intensity of the westerly flow. A positive NAO is related to mild, wet winters and increased storminess¹⁻⁸.
2. Atmospheric blocking occurs when persistent high-pressure systems interrupt the normal westerly flow over middle and high latitudes^{9,10}.
3. The Atlantic Multidecadal Oscillation (AMO) describes fluctuations in North Atlantic sea surface temperature with a 50–90 year period¹¹⁻¹⁵ affecting the large-scale atmospheric circulation¹⁵.



What is already happening?

The NAO has high interannual variability but shows no significant trend during the last century. After an increase from 1960 to 1990 (with more frequent wet and mild winters), the NAO index returned to lower values and after 1990 the blocking pattern shifted eastwards^{16,17} and the duration increased, with more stationary circulation patterns as a consequence¹⁸. However, there is low confidence in the changes concerning blocking patterns¹⁹.
The AMO warmed from the late 1970s to 2014 as part of natural variability. Recently, the AMO began transitioning to a negative phase again²⁰. Paleoclimate reconstructions and model simulations suggested that the AMO might change its dominant frequency over time^{21,22}. However, the impact of the AMO on Northern European climate is independent of its frequency^{14,15}.



What can be expected?

In the future, the NAO is very likely to continue to exhibit large natural variations, similar to those observed in the past. It is likely to become slightly more positive (more frequent wet and mild winters) on average, as a response to global warming¹⁹. Trends in the intensity and persistence of blocking remain uncertain²³. Even under weak global warming the AMO is expected to respond very sensitively, that is, a shortening of time scale and weakening in amplitude²⁴.



Knowledge gaps

While climate models are able to simulate the main features of the NAO, its future changes may be sensitive to boundary processes, like e.g. stratosphere-troposphere interactions or atmospheric response to Arctic sea ice decline, which are not yet well represented in many climate models¹⁹. Most global climate models still underestimate the frequency of blocking over the European-Atlantic sector¹⁹.



Policy relevance

The impact of anthropogenic greenhouse gas emissions might change the large-scale circulation that connects northern Europe with the North Atlantic region. Small changes in the flow would have large consequences for the climate in the Baltic Sea region, i.e., more a maritime or continental climate.



Sea ice

Linked parameters:
Air temperature, Water temperature, Solar radiation, Precipitation, Sea level, Wind, Waves, Sediment transportation, Microbial community and processes, Benthic habitats, Pelagic and demersal fish, Coastal and migratory fish, Waterbirds, Marine Mammals, Coastal protection, Offshore wind farms, Ecosystem function, Shipping, Tourism, Fisheries, Aquaculture, Blue carbon storage capacity, Marine and coastal ecosystem services

Authors
Jürgen Hoffort, Federal Maritime and Hydrographic Agency (BSH), Germany

Energy cycle

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goals 13 and 14
UN Convention on Biological Diversity
EU Green Deal
EU Marine Strategy Framework Directive (MSFD)
EU Water Framework Directive (WFD)
EU Maritime Spatial Planning Directive (MSP)
EU Habitats Directive
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Biodiversity Strategy



Description

In the northern regions of the Baltic Sea, ice is present every winter, while further south sea ice occurs only sporadically. As water effectively absorbs heat, whereas sea ice mostly reflects it, the influence of sea ice on the Baltic energy balance is high. A sea-ice cover also limits the atmosphere-ocean exchange and dampens surface waves. While air temperature has the largest influence on the formation and decay of sea ice, wind has a large influence on the spatial distribution and deformation (ridging, rafting).



What is already happening?

Mean change: During the last 100+ years, ice winters have become milder, the ice season shorter (-18 days at Kemi/Bothnian Bay and -41 days at Loviisa/Gulf of Finland)¹ and the maximum ice extent has decreased by about 30% (6,700 km² decade⁻¹). Indices based on the total winter ice volume show a decreasing trend in the period 1985-2015 (more than 10%/decade in many regions)².
Extremes: The maximum ice extent in the Baltic Sea, including Kattegat, varies from year to year between 40,000 and 420,000 km². The probability of severe ice winters has decreased, an extent larger than 300,000 km² occurred in 16% of the last 100 winters, compared to 3.3% of the last 30.³



What can be expected?

Mean change: In the future, it is very likely that the maximum sea-ice extent will decrease (by between 6,400 (RCP4.5) and 10,900 (RCP8.5) km² per decade)⁴. The thickness of level ice is also very likely to decrease, but there are still large uncertainties for the thickness of ridged ice⁵. The number of days with ice and length of the ice season are likely to decrease, but with considerable regional differences in the magnitude⁶.
Extremes: Inter-annual ice variability is likely to continue to be large, but the probability of severe to very severe winters will likely decrease⁵.



Knowledge gaps

Sea ice as a brittle material is not well represented in numerical climate models⁷. The fact that ice dynamics, like rafting and ridging, are not well-represented also leads to large uncertainties in sea-ice thickness and albedo (i.e., amount of sun light reflected/absorbed). There is only limited information about sea-ice thickness and ice categories and long data sets for these parameters are sparse.



Policy relevance

The importance of sea-ice change is higher in the northern part of the Baltic Sea, especially for ringed seals and shipping. Shipping will be affected through less restrictions on routes and less need for icebreakers, but less ice cover on average does not mean absence of severe ice winters nor of the presence of pack-ice/ridging. Diminishing ice cover also increases the risk and severity of coastal erosion in vulnerable areas. A lack of ice cover should have an influence on the planning of coastal protection, and policies for this may need to be adapted.



Solar radiation

Linked parameters:
Air temperature, Water temperature, Large-scale atmospheric circulation, Sea ice, Stratification, Precipitation, Ecosystem function, Tourism

Authors
Anna Rutgersson, Uppsala University, Sweden
Thomas Carlund, Swedish Meteorological and Hydrological Institute (SMHI), Sweden

Energy cycle

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goals 2 and 14
EU Green Deal
EU Water Framework Directive (WFD)
EU National Emissions Ceilings Directive (NECD)
EU Common Agricultural Policy (CAP)
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Biodiversity Strategy



Description

Solar radiation is the engine of the climate system. Radiation emitted by the sun varies little. Hence, apart from the variation with the time of the year and day, radiation at the surface depends largely on cloudiness. Total cloudiness comprises clouds at all levels (low, medium, and high) and is related to the general atmospheric circulation as well as the water cycle. A cloud layer often reflects 40% to 80% of incoming solar radiation. Atmospheric aerosols have a smaller, but significant effect on solar radiation, both directly and indirectly, through interaction with clouds.



What is already happening?

Mean change: Multidecadal variations in solar radiation, called “dimming” and “brightening”, have been observed in Europe and other parts of the world, especially in the northern hemisphere¹⁻³. Aerosol-induced multidecadal variations in surface solar radiation could be expected also over oceans⁴, but long-term measurements are lacking. Satellite cloudiness trends since the 1980s differ for many areas but seem to agree on a decline over the Baltic Sea region⁵. Records indicate weak but significant negative trends (0.5–1.9% per decade) for global as well as for northern mid-latitude cloudiness.



What can be expected?

Mean change: Future change is uncertain. Global climate models indicate an increase in surface solar radiation, highest over southern Europe and decreasing towards north, but still showing a slight increase over the Baltic Sea region. However, regional climate model runs could instead show a decrease in surface solar radiation over the Baltic Sea region⁶. Unknown future aerosol emissions add to the uncertainty.



Knowledge gaps

Multidecadal variations in surface solar radiation are generally not well captured by current climate model simulations^{7,8}. The extent, to which the observed surface solar radiation variations are caused by natural variation in cloudiness induced by atmospheric dynamic variability^{9,10}, anthropogenic aerosol emissions^{2,8,11,12} or perhaps other causes, is not well understood.



Policy relevance

Solar radiation influences biological activity and ecosystems, through effects on phytoplankton and algal blooms. Altered solar radiation would either increase or decrease biological activities (e.g., photosynthesis). Policy actions to reduce air pollution will impact solar radiation and thus climate change, as reduced air pollution increases the solar radiation reaching the surface. Reducing atmospheric aerosol particle concentrations is important to improve air quality and public health. Currently there is a lively debate related to geoengineering, including methods of increasing reflection of solar radiation back into space, to reduce its heating effect on a global scale.



Salinity and saltwater inflows

Linked parameters:
Stratification, Precipitation, River runoff, Carbonate chemistry, Sea level, Wind, Oxygen, Microbial community and processes, Benthic habitats, Pelagic and demersal fish, Coastal and migratory fish, Non-indigenous species, Ecosystem function, Aquaculture, Blue carbon storage capacity

Authors
H.E. Markus Meier, Leibniz Institute for Baltic Sea Research Warnemünde (IOW), Germany and Swedish Meteorological and Hydrological Institute (SMHI), Sweden
Jan H. Reilman, Federal Maritime and Hydrographic Agency (BSH), Germany

Water cycle

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goal 14
UN Convention on Biological Diversity
EU Marine Strategy Framework Directive (MSFD)
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Biodiversity Strategy



Description

Salinity is an important variable for density, which controls the dynamics of currents in the ocean. Salinity also affects Baltic Sea communities, for example species distribution. Due to freshwater supply from the Baltic Sea catchment and the limited water exchange with the global ocean, surface salinity gradates from > 20 g kg⁻¹ in Kattegat to < 2 g kg⁻¹ in the Bothnian Bay. The dynamics of the Baltic Sea are characterized by a pronounced, perennial vertical gradient in salinity.

Large, meteorologically driven saltwater inflows, so-called Major Baltic Inflows (MBIs), sporadically renew the deep water with saline, oxygen rich water and this is the only process that effectively ventilates the deep water^{1,2}.



What is already happening?

Mean change: There are no statistically significant trends in salinity, river flow or MBIs on centennial timescales since 1850, but pronounced multidecadal variability, with a period of about 30 years²⁻⁸. Model results suggest that a decade of decreasing salinity, like the 1983-1992 stagnation, appears approximately once per century due to natural variability⁹. Baltic Sea salinity is also influenced by the Atlantic Multidecadal Oscillation with a 50–90-year period¹⁰. Since the 1980s, bottom salinity has increased, and surface salinity has decreased¹¹.

Extremes: The frequency of MBIs shows no statistically significant trend during instrumental (1886–2017) and paleoclimate periods^{2,9}.



What can be expected?

Mean change: An increase in river runoff from the northern catchment area will tend to decrease salinity, but a global sea level rise will tend to increase salinity, because the water level above the sills at the Baltic Sea entrance and the saltwater imports from the Kattegat would be higher. A 0.5 m higher sea level would increase the average salinity by about 0.7 g kg⁻¹¹². Due to the large uncertainty in projected freshwater supply from the catchment area, wind and global sea level rise, salinity projections show a widespread trend, and no robust changes have been identified¹³⁻¹⁶.

Extremes: The frequency of MBIs is projected to slightly increase¹⁷.



Knowledge gaps

Due to the large natural variability and uncertain changes in the regional water cycle, including precipitation over the Baltic Sea catchment area, in wind fields and in global sea level, the confidence in future salinity projections is low¹⁴. Modelling data show that the north-south gradient has changed with an increase in runoff in the North, and a decrease in the South⁵. Not much is known about changes in salinity composition and their large decadal variability. Changes in total salt import have not been adequately investigated. Changes in the large-scale circulation in the Baltic Sea are not well understood^{18,19}.



Policy relevance

Salinity and the ventilation of the deep water with oxygen that is associated with MBIs, are important drivers of the Baltic Sea ecosystem functioning and structure, including reproduction of commercially important marine fish species, such as cod^{20,21}. The distribution of freshwater and marine species and the overall biodiversity depends strongly on salinity and oxygen concentrations²⁰. Hence, the salinity dynamics is a major factor for the implementation of marine policies²¹.



Stratification

Linked parameters:
Water temperature, Solar radiation, Salinity and saltwater inflows, Wind, Oxygen, Pelagic and demersal fish, Nutrient concentrations and eutrophication, Ecosystem function

Authors
Matthias Gröger, Leibniz Institute for Baltic Sea Research Warnemünde (IOW), Germany
H.E. Markus Meier, Leibniz Institute for Baltic Sea Research Warnemünde (IOW), Germany,
and Swedish Meteorological and Hydrological Institute (SMHI), Sweden
Umas Lips, Tallinn University of Technology, Estonia
Jan H. Reiffman, Federal Maritime and Hydrographic Agency (BSH), Germany

Water cycle ■■■

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goal 13 and 14
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Biodiversity Strategy



Description

Stratification is determined by density gradients resulting from the temperature and salinity distributions in the sea. Stratification controls vertical and horizontal circulation and transport of water masses.

In the Baltic Sea, the strongest vertical density gradients correspond to the thermocline (maximum temperature gradient) and halocline (maximum salinity gradient). The thermocline develops at 10–20 m depth during the warm season, whereas a pronounced halocline persists over the year at 60–80 m in most deep regions. Wind stress working on the sea surface can potentially homogenize the water column fully in some shallow water regions or partly in deep water regions, thus influencing stratification.



What is already happening?

● In the past, the haline stratification has been dominated by sporadic inflows from the adjacent North Sea and river discharge. Long-term trends in Baltic Sea salinity¹ or halocline depth² have not been demonstrated, but a trend towards increased horizontal sea surface salinity difference between the northern and southern Baltic Sea during 1920–2005 has been detected, resulting in increased horizontal gradients³. In addition, during 1856–2005 sea surface temperatures increased by about 0.03 and 0.06 °C decade⁻¹ in the northeastern and southwestern areas, respectively, probably resulting in increased vertical stratification⁴.

Furthermore, stratification increased in most of the Baltic Sea during 1982–2016, with the seasonal thermocline and the perennial halocline strengthening⁵.



What can be expected?

● Theoretical considerations imply that stronger stratification is favoured by increased freshwater supply to the Baltic Sea drainage basin accompanied by the supply of deep salt-rich waters from the North Sea, as well as warming of the surface layer. Thus, future development of stratification mainly depends on how much the Baltic Sea surface will warm compared to deeper layers and how freshwater supply and saltwater inflows will change.

Multi-model scenario simulations have confirmed increased vertical summer stratification due to warming³ whereas projections of salinity and related haline stratification changes are rather uncertain^{6–8}.



Knowledge gaps

The complex interplay between changes in temperature, wind and precipitation makes it difficult to project the impact of future climate on stratification. The circulation and its influence on stratification is not well understood, and the same is true for the influence of mixing processes (e.g., winter convection) on stratification. Sea surface temperature can be expected to follow air temperature, due to air-sea heat exchange, but the fate of salinity, and hence vertical salinity gradients, is uncertain. Due to the pronounced multidecadal variability in measured water temperature and salinity, projections of long-term trends based on past changes in climate cannot be made.



Policy relevance

Stratification is an important driver of ecosystem functioning and structure, controlling the vertical flux of oxygen between the well-ventilated surface waters and oxygen-poor deep waters, affecting for example benthic habitats and the reproduction of cod. In addition, an increased thermal stratification during summer can decrease vertical nutrient transport from deeper layers to the euphotic zone, thereby limiting nutrient supply and potentially affecting algal and cyanobacterial blooms, at least at the species level⁹. To counteract oxygen depletion in the deep water, various geoengineering methods such as pumping of water below the halocline reducing vertical stratification have been discussed, but their effectiveness at basin-scale has been questioned in scientific literature¹⁰.



Precipitation

Linked parameters:
Large-scale atmospheric circulation, Sea ice, Solar radiation, Salinity and saltwater inflows, River run-off, Riverine nutrient loads and atmospheric deposition, Tourism

Authors
Jukka Käyhkö, University of Turku, Finland
Erik Kjellström, Swedish Meteorological and Hydrological Institute (SMHI), Sweden

Water cycle ■■■

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goal 13
EU Maritime Spatial Planning Directive (MSP)
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Biodiversity Strategy



Description

Precipitation forms in the atmosphere when air is saturated with water vapor, and cloud droplets or ice crystals grow large enough by condensation or deposition, respectively, to precipitate from the cloud under gravity. Depending on the conditions in the cloud and along the way to the ground, the falling particles are in liquid or frozen form (drops, flakes, hail, etc.). Precipitation is strongly linked with other variables of the water cycle. As the amount of water vapour that can be held in air increases with temperature, so does precipitation. Generally, there is more precipitation in summer than in winter in large parts of the Baltic Sea region^{1,2}. Precipitation is strongly modified by the ground surface elevation and land-sea contrasts, implying that the large-scale circulation of the atmosphere including wind direction and vertical stability are important factors.



What is already happening?

● **Mean change:** During the 20th century, precipitation was subject to large variations³, which increases the difficulty in determining statistically significant trends or regime shifts. Generally, precipitation increases in winter. Sweden shows an overall wetting trend in particular since the 1950s⁴. In Finland, the overall increase detected for 1961–2010 is neither regionally consistent nor always statistically significant⁵. The same holds for the Baltic countries⁶.

● **Extremes:** Daily amounts of precipitation extremes typically range from 8 to 20 mm, being more numerous in summer⁷. Extreme precipitation intensity has been rising in the period 1960–2018. The maximum annual five consecutive days precipitation index (Rx5d) has shown a significant increase of up to 5 mm per decade over the eastern Baltic Sea catchment⁸. The change has been most pronounced in winter.



What can be expected?

● **Mean change:** Average precipitation amounts are expected to increase in the future. The relative increase will be largest in winter in the North. Most simulations show increasing summer precipitation for the northern parts, while for the intermediate and southern parts of the region, the direction of change is uncertain⁹.

● **Extremes:** Warming increases the potential for extreme precipitation due to intensification of the hydrological cycle associated with growth of atmospheric moisture content. Regional climate models indicate an overall rise in the frequency and volume of heavy precipitations in all seasons. The projected increases in northern Europe might be significant throughout all the seasons from 2050 onwards^{9,10}. Autumn is projected to see the largest increase of high precipitation days. The number of drought events per year may decrease, while their length may increase⁹.



Knowledge gaps

Different methods and data sets used in national studies in the region imply that the knowledge of the precipitation climate is not fully coherent.

Compared to traditional “high-resolution” models, the recent very high-resolution climate model projections, at 1–3 km resolution, have proven to show better agreement with observations in representing precipitation extremes, and sometimes also larger climate change signals, but these are yet to be built for the Baltic Sea region.



Policy relevance

Adaptation to changes in precipitation will have to involve consideration of both increasing precipitation with a risk for flooding and decreasing precipitation with a risk for drought. This will have implications for agricultural policies as well as urban flood and storm-water management.



Authors
Jukka Käyhkö, University of Turku, Finland
Water cycle



River run-off

Linked parameters:
Air temperature, Salinity and saltwater inflows, Precipitation, Riverine nutrient loads and atmospheric deposition, Benthic habitats, Coastal and migratory fish, Coastal protection

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goal 13
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Floods directive
EU Biodiversity Strategy



Description

Runoff describes the amount of flowing water, typically given in litres per second per square kilometre ($\text{l s}^{-1} \text{ km}^{-2}$) to allow comparisons between differently sized rivers. Runoff can also be given in millimeters per year (mm a^{-1}), allowing comparisons with precipitation and evaporation. Discharge refers to channel flow, typically given as cubic metres per second ($\text{m}^3 \text{ s}^{-1}$). Floods are extreme run-off events of water submerging usually dry land. In the Baltic Sea region, floods typically occur in the spring-time snow-melt period, or in connection to heavy/long-lasting rain. Floods are closely linked to precipitation, temperature (melting, evaporation), wind, and catchment properties (land use, topography).



What is already happening?

- Mean change:** No statistically significant change in total annual river runoff has been detected during the last centuries^{1,2}. Large decadal and regional variations occur³. In the northern Baltic Sea and the Gulf of Finland, larger river runoff is statistically associated with warmer air temperature and increased precipitation, while further south, decreased annual runoff is associated with rising air temperatures¹. Over the 20th century, winter discharge has increased, while spring floods have decreased⁴.
- Extremes:** According to an example from Sweden, there is no significant trend in daily high flow over the past 100 years⁵.



What can be expected?

- Mean change:** The total runoff to the Baltic Sea has been projected to increase from present day by 2-22% with warming temperatures^{6,7}. The increase will take place mostly in the North^{3,6,8}, with potentially decreasing total runoff in the South⁹. Winter runoff will increase due to intermittent melting⁸.
- Extremes:** Floods are projected to decrease in the North, due to repeated melting and thinner snowpack, but increase south of 60°N due to higher precipitation^{10,11}. Large spring floods will decrease by up to 20%¹⁰.



Knowledge gaps

The impacts of observed precipitation changes on stream flow are unclear¹². The effects of how climate model results are currently transferred to the hydrological model are still inadequately understood. More research is needed to quantify the accuracy and uncertainty associated with various bias correction methods⁷. Several uncertainties are associated with the impact modelling, including parameter calibration against historical data and model structure uncertainty.




Policy relevance

Seasonal runoff changes will affect sediment and nutrient loads and thereby eutrophication of the Baltic Sea. Changes in the timing of floods will influence risks for riverside settlements. The HELCOM Baltic Sea Action Plan requires nutrient load reduction from the signatory countries. However, plans by EU Member States lack ambition in nutrient reduction implementation¹³. Flood hazard mitigation requires both short-term (rescue) and long-term (planning and construction) measures. Directive 2007/60/EC on the assessment and management of flood risks requires adequate and coordinated measures to reduce flood risk. As new projections continuously become available, climate change is important to include in river runoff and flood policies.



Authors
Karol Kuliński, Institute of Oceanology Polish Academy of Sciences, Poland
Gregor Rehder, Leibniz Institute for Baltic Sea Research Warnemünde (IOW), Germany
Anna Rutgersson, Uppsala University, Sweden
Carbon and nutrient cycle



Carbonate chemistry

Linked parameters:
Water temperature, Salinity and saltwater inflows, Riverine nutrient loads and atmospheric deposition, Wind, Waves, Microbial community and processes, Benthic habitats, Non-indigenous species, Nutrient concentrations and eutrophication, Ecosystem function, Blue carbon storage capacity

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goals 12 and 14
EU Green Deal
EU Marine Strategy Framework Directive (MSFD)
EU Water Framework Directive (WFD)
EU Common Agricultural Policy (CAP)
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Biodiversity Strategy



Description

The carbonate system (CO_2 system) characterized by the thermodynamic equilibria between hydrogen ions (reported as pH) and the different CO_2 species (CO_2 , H_2CO_3 , HCO_3^- , CO_3^{2-})^{1,2} is the main determinant of the acid/base balance in seawater and of seawater pH. Ocean acidification is the decrease of seawater pH, mainly due to the rising CO_2 concentration in the atmosphere and its exchange with the surface seawater². The exchange of CO_2 between the water and the atmosphere is controlled by the air-sea surface difference in partial pressure of CO_2 ($p\text{CO}_2$) and the efficiency of the transfer processes, with wind speed being the dominating parameter³⁻⁵.



What is already happening?

- Intra-annual variation of $p\text{CO}_2$ in the Baltic Sea surface waters is controlled by biological processes (organic matter production and remineralization) and by changes in the mixed layer depth and the sea surface temperature^{1,2,6}. Eutrophication has enhanced both production and remineralization of organic matter and thus increased the amplitude of seasonal changes in $p\text{CO}_2$ and pH^{7,8}. The Baltic Sea is a CO_2 sink in summer and a source in winter^{1,6}. However, the annual net CO_2 flux is unknown. The increase in total alkalinity (a measure of the buffer capacity) entirely mitigates ocean acidification in the Gulf of Bothnia and significantly reduces it (to about $-0.0012 \text{ year}^{-1}$) in the central Baltic Sea⁹.**



What can be expected?

- Future changes in atmospheric $p\text{CO}_2$ and total alkalinity will influence sea-water $p\text{CO}_2$ and therefore pH^{2,4,7-10}. The projected runoff increase to the northern Baltic Sea may lower alkalinity and pH, due to decreased salinity⁷. However, higher atmospheric $p\text{CO}_2$ will enhance weathering on land and release alkalinity from the catchment, while eutrophication may increase internal alkalinity generation, leaving the net effect unknown^{7,8}. Even if alkalinity in the Baltic Sea should increase, a doubling of atmospheric $p\text{CO}_2$ will still result in lower pH⁷.**



Knowledge gaps

Due to the high spatial and temporal variability in seawater $p\text{CO}_2$, it is currently unclear whether the Baltic Sea as a whole is a net sink of CO_2 or a net source^{1,2,6}. Since the origin of the currently observed alkalinity increase in the Baltic Sea is unclear, it is uncertain whether this increase will continue as strongly in the future⁹. The ecosystem productivity in the period after the spring bloom (from mid-April until mid-June) is not quantitatively understood, due to an observed continuation of $p\text{CO}_2$ decrease even after the surface nitrate pool is depleted^{1,2}.



Policy relevance

The rising atmospheric CO_2 concentration is one of the main drivers shaping the structure of the marine CO_2 system and the dominant cause of ocean acidification, which may influence marine organisms, especially those building their exoskeletons out of calcium carbonate^{1,2,9}. The implementation of the Baltic Sea Action Plan resulting in comparatively low nutrient loads and favourable oxygen conditions may minimize wintertime pH reduction⁷.



Riverine nutrient loads and atmospheric deposition

Authors
Oleg Savchuk, Baltic Sea Centre, Stockholm University, Sweden
Michelle McCrackin, Baltic Sea Centre, Stockholm University, Sweden
Mikhail Sofiev, Finnish Meteorological Institute (FMI), Finland

Carbon and nutrient cycle ■■■

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goals 2 and 14
EU Green Deal
EU Water Framework Directive (WFD)
EU National Emissions Ceilings Directive (NECD)
EU Common Agricultural Policy (CAP)
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Biodiversity Strategy

Linked parameters:
Air temperature, Precipitation, River run-off, Carbonate chemistry, Microbial community and processes, Nutrient concentrations and eutrophication, Ecosystem function, Shipping



Description

External nutrient inputs from land and atmosphere are the major long-term drivers of Baltic Sea eutrophication^{1,2}. Land loads and atmospheric deposition are determined by both natural (precipitation, river runoff, temperature) and anthropogenic (demographic, agricultural, and industrial development, wastewater treatment, international shipping) factors. These factors change both over time (changes of seasonality, long-term trends, and lags due to the watershed processes) and space (north-south gradients in climate and land use, east-west gradients in socio-economic features and climate)³. Atmospheric deposition is additionally determined by long-range transport from Central, Western and Eastern Europe and, for the Gulf of Finland, from Russia^{4,5}.



What is already happening?

Substantial reductions of riverine nutrient loads have been achieved since the 1980s⁶⁻⁹. Since there are no statistically significant trends in annual river discharges¹⁰, these reductions are attributed to socio-economic development, including protective measures, rather than to climate-related effects^{8,11}. The total nitrogen deposition to the Baltic Sea has also been substantially decreasing since the 1980s, due to overall reduction of European emissions¹². However, the reduction of nitrogen emission and deposition has slowed down since the beginning of the 21st century^{13,14}. Atmospheric phosphorus deposition amounts and trends remain highly uncertain^{4,15,16}.



What can be expected?

Projections suggest that river discharge will increase in the northern Baltic Sea region, while the discharge will decrease in the southern region¹⁰, thus potentially increasing and decreasing waterborne nutrient inputs, respectively. Leaking of excessive phosphorus, accumulated in agricultural soils, will delay the effects of mitigation measures¹⁷. Simulations with a range of scenarios suggest that land-based nutrient management will have greater effect on nutrient loads than greenhouse gas emissions¹⁸⁻²². Atmospheric deposition can be affected by changes in emissions^{5,23}, for example by increased ammonia evaporation due to rising temperature^{24,25}.




Knowledge gaps

Besides common uncertainties inherent to regionalization of climate scenarios for precipitation and river runoff¹⁰, an important source of uncertainty is poor quantitative knowledge on long-term response of terrestrial biogeochemical processes, particularly changes in the soil nutrient pools, to the climate changes²⁶. Phosphorus sources and transports^{4,16,17} as well as ammonia emission and its dynamics^{12,24,25} are among the least known processes controlling the atmospheric nutrient deposition. How the anthropogenic drivers of land loads (land use, agricultural practices, wastewater treatment, net anthropogenic nutrient inputs, etc.) will change in response to both climate change and socio-economic development is highly uncertain²⁷.



Policy relevance

Reduction of nutrient inputs is considered the most important measure for mitigating Baltic Sea eutrophication, both in coastal and offshore waters¹¹. Implementation of corresponding measures within the Water Framework Directive, Baltic Sea Action Plan, Marine Strategy Framework Directive, and National Emissions Ceilings Directive has already resulted in significant decreases of land loads and atmospheric deposition. However, effects of climate change on the transfer of nutrients from land to sea have not yet been appropriately incorporated in these policies. Additionally, the ammonia (NH₃) emissions, which unlike the nitrogen oxide (NO_x) emissions have been largely disregarded, will require large reduction efforts and policy and public support²⁵.



Sea level

Authors
Christian Dieterich, Swedish Meteorological and Hydrological Institute (SMHI), Sweden
Jürgen Hoffort, Federal Maritime and Hydrographic Agency (BSH), Germany
H.E. Markus Meier, Leibniz Institute for Baltic Sea Research Warnemünde (IOW), Germany, and Swedish Meteorological and Hydrological Institute (SMHI), Sweden
Jani Särkkä, Finnish Meteorological Institute (FMI), Finland
Ralf Weisse, Helmholtz Center for Material and Coastal Research (HZG), Germany
Eduardo Zorita, Helmholtz Center for Material and Coastal Research, (HZG), Germany

Sea level and wind ■■■

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goal 13
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Biodiversity Strategy



Description

The relative sea level in the Baltic Sea rises when melt water is added to the global ocean, when the water expands by warming, or the land sinks¹. The Baltic Sea level varies between years and seasons² and is generally highest in winter³, especially in mild winters with above average winds^{4,5}. Periods of strong westerlies temporarily fill the Baltic Sea with extra water from the North Sea⁶. Increased mean sea level leads to higher storm surges. Storms trigger sea level oscillations^{7,8} across the Baltic Sea, meteotsunamis^{9,10} (sea level extremes travelling in phase with atmospheric low pressure systems), and wave set-up where breaking waves increase the sea level locally by up to half a metre¹¹.



What is already happening?

Mean change: Global mean sea level rose 1-2 mm yr⁻¹ during the 20th century¹²⁻¹⁴. Presently, rates of 3-4 mm yr⁻¹ are estimated over shorter periods¹²⁻¹⁵. For instance in Stockholm, absolute sea level rose by about 20 cm from 1886 to 2009¹⁶. Land uplift in the northern Baltic is still faster than absolute sea level rise so that, relative to land, sea level there is still falling^{14,17-19}.

Extremes: Storm surges are a threat to low-lying Baltic Sea coastlines^{9,20,21}. No long-term increasing trend has been found for the 20th century for extreme sea levels in the Baltic Sea relative to mean changes^{14,22,23}.



What can be expected?

Mean change: Global sea level rise will accelerate^{12,13,24}. Current projections estimate Baltic sea level rise to about 87% of the global rate^{25,26}. Estimates for global mean sea level rise by 2100 are 43 cm (RCP2.6) to 84 cm (RCP8.5)¹³. The likely ranges for these estimates are 29 to 56 cm (RCP2.6) and 61 to 110 cm (RCP8.5)¹³.

Extremes: How extremes will change is uncertain, as they depend on the path of future low pressure systems^{6,27}. In the southern Baltic Sea, extremes that are rare today will become more common due to mean sea level rise^{28,29}.



Knowledge gaps

Research is needed on natural variability in drivers of storm surges in the Baltic Sea^{21,30-32}. How much the Baltic sea level rises compared to the global mean²⁵ includes large uncertainties from Antarctic ice sheet melting and climate change in the Atlantic Ocean and the Baltic Sea. Sea level will rise proportionally more on the shallow shelf regions around the continents than in the deeper, open ocean³³, but the extent of this effect has not been evaluated for the Baltic Sea. Storm surges and other hazards can turn into disasters if they coincide³⁴. Little is known about the interaction of storm surges and other extreme events.



Policy relevance

Mean sea level rise and extreme events are of great importance, for example for urban planning and commercial ports, and a challenge for flood protection. Ports can adapt to mean sea level rise by building higher quays or relocating. Shipping lanes may need to be dredged less, and ships with a deeper draught can come to port. Coastal flooding can be prevented by protective structures, such as the St Petersburg Flood Prevention Facility Complex, the Stockholm Slussen (Sluice) project, and levees along the German and Polish coasts.



Wind

Linked parameters:

Large-scale atmospheric circulation, Sea ice, Salinity and saltwater inflows, Stratification, Carbonate chemistry, Sea level, Waves, Sediment transportation, Coastal protection, Offshore wind farms, Shipping, Tourism, Fisheries

Authors
Mika Rantanen and Terhi Laurila, Finnish Meteorological Institute (FMI), Finland
Erik Kjellström, Swedish Meteorological and Hydrological Institute (SMHI), Sweden
Anna Rutgersson, Uppsala University, Sweden

Sea level and wind

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Convention on Biological Diversity
EU Green Deal
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Biodiversity strategy



Description

The wind climate in the Baltic Sea region is determined by the large-scale atmospheric circulation. Typically, the strongest wind speeds are associated with the passage of strong extratropical cyclones. These systems, and thus wind extremes, are most frequent and intense in the winter half of the year. In addition, strong local winds can occur in association with thunderstorms that are most pronounced in summer.



What is already happening?

Mean change: Owing to the large climate variability in the Baltic Sea region, it is unclear whether there is an overall trend in mean wind speed. Detected trends in the wind climate differ between seasons and depend on the chosen time period for which the trend is calculated because the internal variability is large^{1,2}. For example, mean wind speeds at the Finnish and Swedish coastlines show a slightly negative trend since the 1950s^{3,4}.

Extremes: Maximum wind speeds at the Finnish coastline show a weakening trend⁴, attributed to storm tracks shifting northwards^{5,6}. Many studies show contradicting storminess trends in the Baltic region¹.



What can be expected?

Mean change: Projected changes in wind climate are highly uncertain due to large natural variability in the Baltic Sea area⁷. Climate model simulations project a slight but significant wind speed increase in autumn and a decrease in spring⁸. Some studies mention increased future wind speeds in areas no longer covered by sea ice^{7,9,10}.

Extremes: Projected changes in extreme winds are uncertain due to differences in atmospheric circulation among climate model projections⁹. It is projected that by 2100, severe wind gusts associated with thunderstorms can increase in frequency during summer¹¹.



Knowledge gaps

Changes in wind climate are among the least certain aspects of climate change in the Baltic Sea area. This is because there are several differing projections of atmospheric circulation between different climate models, reflected in a large spread of future wind speed changes. Enhancing the ensemble sizes and improving high-resolution climate models can help in extracting possible anthropogenic signals from the large natural variability.



Policy relevance

Changes in wind extremes are relevant for example for coastal infrastructure, coastal tourism, and shipping in the Baltic Sea. Storm surges, which are typically associated with high wind speed events, can cause harm to various parts of the coast, and can damage densely populated coastal cities. Knowledge of wind extremes in combination with sea ice events is central for constructing and managing offshore wind and wave energy installations. Adaptation to such events is often considered in the management of coastal infrastructures. Future infrastructure would benefit from better wind models and from considering a higher wind stress tolerance.



Waves

Linked parameters:

Sea ice, Sea level, Wind, Sediment transportation, Benthic habitats, Marine mammals, Coastal protection, Offshore wind farms, Shipping, Tourism, Fisheries, Aquaculture, Blue carbon storage capacity

Authors
Ralf Weisse, Helmholtz-Zentrum Geesthacht (HZG), Germany
Christian Dieterich, Swedish Meteorological Institute (SMHI), Sweden

Sea level and wind

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goal 13
EU Marine Strategy Framework Directive (MSFD)
EU Water Framework Directive (WFD)
EU Maritime Spatial Planning Directive (MSP)
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Habitats Directive
EU Biodiversity Strategy



Description

Wind waves are generated by the action of wind on the sea surface. In the Baltic Sea, the highest waves typically occur during long-lasting storms with high wind speeds and long fetch (the distance over which the wind blows).

The wave climate is characterized by parameters such as significant wave height, period, and mean direction. The Baltic Sea wave climate has a pronounced seasonal cycle with higher waves in winter. Breaking waves can substantially increase coastal sea level (wave set-up). Waves are the major driver of nearshore sediment transport. High storm waves are the primary determinants of the extent of erosion.



What is already happening?

Mean change: There are no significant long-term trends in wind speed and direction but considerable decadal variability¹. Correspondingly, there are no clear indications of long-term trends in wave height².

Extremes: From a long-term perspective, no robust signals of changes in Baltic wave climate can be detected².



What can be expected?

Mean change: Changes in Baltic Sea wave climate are strongly linked to changes in wind climate and are highly uncertain³⁻⁵.

There is high confidence on reduced ice cover which may increase fetch, and perhaps change the wave climate⁶.

By 2100, changes in significant wave height are projected to be around 5% higher than today, in particular in the north and east of the Baltic Sea⁵. However, such changes are superimposed by substantial multi-decadal and inter-simulation variability and are not conclusive because only one climate model was considered⁵.

Extremes: Changes in extreme wave heights result from changes in high wind speeds, which are highly uncertain¹.



Knowledge gaps


There are only a few projections of future wind-wave climate, and assessments of changes in longshore sediment transport including its spatial and temporal variability available. Larger ensembles of scenario simulations driven by many global climate models are needed. Little is known on the role of coastal processes for the development of waves, e.g., wave set-up.

Given the pronounced inter-decadal variability, detection of significant trends and attribution studies to disentangle the impact of changing climate and other drivers, together with the development of decadal predictions of wave climate, would be useful.



Policy relevance

Increase in offshore wave action will directly impact the safety of shipping, fisheries, and offshore operations. Increase in coastal wave action will affect coastal sea level and erosion and be of immediate relevance for coastal protection. Adaptation to changes in wave climate may require for example increasing demands on hull integrity for ships and maritime structures and changes to coastal protection strategies and policies. So far, this is not the case and policymakers need to take this prospective change into account, especially when developing more windfarms in the Baltic Sea region to meet renewable energy goals.



Sediment transportation

Linked parameters:
Sea ice, Sea level, Wind, Waves, Coastal protection, Tourism

Authors
Wenyan Zhang, Helmholtz-Zentrum Geesthacht (HZG), Germany
Joonas Virtasalo, Geological Survey of Finland, Finland

Sea level and wind

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goal 13
EU Maritime Spatial Planning Directive (MSP)
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Biodiversity Strategy



Description

Sediment transport in marine environment is triggered mainly by currents and waves. Its direct consequence is erosion or accretion, leading to a gradual change of coastal landform and seabed morphology.

Short-term, small-scale sediment transport is driven by a variety of local conditions including winds, water level, waves, currents, as well as the initial state of the system. Long-term, large-scale sediment transport is primarily controlled by the type of sediment and its supply, modulated by large-scale processes, notably sea level, storms, regional wind and wave pattern, and local engineering structures.



What is already happening?

Mean change: Baltic Sea coastlines currently show a gradient from a maximum land-rise of +9 mm year⁻¹ in the North to a subsidence by -2 mm year⁻¹ in the South¹. Dominance of mobile sediments makes the southern and eastern coasts vulnerable to wind-wave induced transport². Dominating westerly winds lead to mainly west-east sediment transport and an alternation of glacial till cliffs (sources), sandy beaches, and spits (sinks)².

Extremes: Many sandy beaches and moraine cliffs are frequently eroded by storm surges and subsequently transported by currents³. Land uplift exposes shallow seafloor sediment to erosion by storm waves and transportation by currents.



What can be expected?

Mean change: Global sea level rise is accelerating⁴. Consequently, sediment transportation can be expected to increase in coastal areas. The rate depends both on sea level rise and storm frequency and trajectory⁵. Due to prevailing westerly winds in the Baltic Sea region, the dominant regional transport pattern is expected to be the same, but with high local variability along coastal sections which are featured by a small incidence angle of incoming wind-waves^{6,7}.

Extremes: Coastal sediment transport by storms depends on surge and wave impact level and is likely to increase as the sea level rises⁸.



Knowledge gaps

There is a lack of comprehensive understanding of the spatial and temporal variability of sediment transportation along the Baltic Sea coastal zone. In general, primary sediment transport is driven by currents and waves produced by the prevailing westerly winds². However, the intensity of secondary transport induced by easterly and northerly winds is poorly understood⁶. Combination with changes in sea level, storm surges (including storm tracks) and sea ice further complicate the understanding of sediment transport^{5,9}. Man-made engineering structures add to the uncertainty in the prediction of sediment transport and coastal erosion patterns¹⁰.




Policy relevance

Sediment transport, especially erosion, is important to coastal planning, construction, and protection. Management strategies include 1) protection by soft or hard measures and 2) leaving some parts in an unguarded state. Soft protection includes, e.g., beach nourishment and vegetation planting in front of foredunes. Hard protection includes groynes, dykes, seawalls, revetments, artificial headlands, and breakwaters. Management efforts differ among countries and are complex when coastal protection in one place leads to morphodynamic changes that disrupt downstream areas and possibly biodiversity¹¹. There are no synergistic measures to address these effects if they occur in other countries, due to differing legislations.

Indirect parameters:
Ecosystem





Oxygen

Linked parameters:

Water temperature, Salinity and saltwater inflows, Stratification, Microbial community and processes, Benthic habitats, Pelagic and demersal fish, Non-indigenous species, Nutrient concentrations and eutrophication, Ecosystem function, Aquaculture

Authors
H.E. Markus Meier, Leibniz Institute for Baltic Sea Research Warnemünde (IOW), Germany and Swedish Meteorological and Hydrological Institute (SMHI), Sweden
Jacob Carstensen, Aarhus University, Denmark
Oleg Savchuk, Stockholm University, Baltic Nest Institute, Sweden

Carbon and nutrient cycle ■■■

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goals 2 and 14
UN Convention on Biological Diversity
EU Green Deal
EU Marine Strategy Framework Directive (MSFD)
EU Water Framework Directive (WFD)
EU Habitats Directive
EU Common Agricultural Policy (CAP)
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Biodiversity Strategy



Description

Dissolved oxygen concentration in the water column is controlled by physical transport (air-sea exchange, advection, and diffusion), water temperature and biological processes such as photosynthesis and demand for oxidation of organic matter and sulfide¹. Deoxygenation and hypoxia occur where the oxygen demand due to elevated concentrations of organic matter concentration in water and sediment cannot be compensated by ventilation¹⁻¹⁰. Hypoxic area is defined as the extent of bottom water with oxygen concentrations below a threshold, commonly set at 2 mL O₂ L⁻¹. Hypoxia is characterized by a scarcity of multicellular life¹.



What is already happening?

Despite decreasing nutrient loads after the 1980s⁵, recently calculated oxygen consumption rates are higher than earlier observed, counteracting the effect of natural ventilation of deep water⁶. Improved oxygen conditions have been observed in some coastal waters, where inputs of nutrients and organic matter have been abated¹¹. However, hypoxia remains common in other coastal areas, with unaltered or even worsening conditions^{4,6}. In 2016, the annual maximum extent of hypoxia covered an area of about 70,000 km², whereas 150 years ago the hypoxic area was presumably small³.



What can be expected?

Projected warming may enhance oxygen depletion in the Baltic Sea by reducing air-sea and vertical transports of oxygen and by reinforcing eutrophication through intensifying internal nutrient cycling and stimulating nitrogen-fixing cyanobacteria blooms¹²⁻¹⁶. However, the future development of deep-water oxygen conditions will mainly depend on the nutrient load scenario. If nutrient loads are high, the impact of warming will be considerable and negative; if low, the effect will be small¹⁵. Scenario simulations suggest that full implementation of the Baltic Sea Action Plan resulting in required load reductions will lead to a significantly improved ecosystem state of the Baltic Sea, irrespective of the driving global climate model^{12,13,15,16}.



Other drivers

Model simulations suggest that elevated historical riverine nutrient loads and atmospheric nutrient deposition since the 1950s have been the most important drivers of oxygen depletion in the Baltic Sea^{3,5,7}. The impacts of other drivers such as the observed warming or eustatic sea level rise were comparatively smaller but still made a significant contribution to, e.g., the size of hypoxic area^{3,7}. There are no statistically significant trends in stratification and saltwater inflows on centennial timescales since 1850. Thus, variations in oxygen transports have caused interannual to decadal variability in oxygen concentrations of the Baltic Sea deep water but could not explain the long-term trend^{3,7}.



Knowledge gaps

A recent assessment suggests that, in addition to internal variability, the biggest uncertainties in projections of biogeochemical cycles are caused (not listed in order of importance) by (i) poorly known current and future bioavailable nutrient loads from land and atmosphere (see also ¹⁷), (ii) differences between the projections of global and regional climate models, in particular, with respect to the global mean sea level rise, wind and regional water cycle, (iii) differing model-specific responses of the simulated biogeochemical cycles to long-term changes in external nutrient loads and climate of the Baltic Sea region, (iv) poorly known long-term pathways of future greenhouse gas emissions^{10,11}, and (v) poorly known sediment properties regarding oxygen demand and nutrient release.



Policy relevance

Oxygen conditions are indispensable prerequisites for the marine ecosystem and are closely related to nutrients. Although nutrient loads have been reduced since the 1980s⁵, the targets for the maximum allowable inputs have not yet been completely achieved¹². In addition, the system's response to changes in nutrient loads is slow, currently preventing the Baltic Sea from attaining good eutrophication status. As global warming will worsen oxygen conditions, full implementation of the load reductions of the Baltic Sea Action Plan (BSAP) is needed. Scenario simulations suggest that this will result in successful, albeit slow, mitigation^{15,16}. The results of ongoing scenario simulations have high relevance for the BSAP.



Microbial community and processes

Linked parameters:

Water temperature, Sea ice, Salinity and saltwater inflows, Carbonate chemistry, Riverine nutrient loads and atmospheric deposition, Oxygen, Nutrient concentrations and eutrophication, Ecosystem function

Authors
Klaus Jürgens, Leibniz Institute for Baltic Sea Research Warnemünde (IOW), Germany
Johan Wikner, Umeå University, Sweden

Biota and ecosystems ■■■

Links to main policies:
UN Sustainable Development Goals 2 and 14
UN Convention on Biological Diversity
EU Green Deal
EU Marine Strategy Framework Directive (MSFD)
EU Water Framework Directive (WFD)
EU Common Agricultural Policy (CAP)
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Bathing water directive
HELCOM Baltic Sea Action Plan
EU Biodiversity Strategy



Description

This overview focuses on bacterioplankton, comprising single-celled prokaryotes, i.e., small organisms that lack a nucleus (Bacteria and Archaea), in the water column, consuming organic carbon as an energy and carbon source. Benthic prokaryotes and protozoa (i.e., unicellular zooplankton and zoobenthos) are also important but not included here. Bacteria are the major transformers of carbon, nitrogen, sulphur, and trace metal cycles in aquatic environments. The supply of organic carbon mainly controls bacterial biomass production. The bacterial community composition changes along the salinity and oxygen gradients of the Baltic Sea¹. Shifts in, e.g., food sources, temperature, and oxygen concentration result in rapid bacterioplankton community changes^{2,3}, with potential impact also on overall ecosystem functions, such as respiration, carbon consumption, and biomass production.



What is already happening?

Long time-series of marine bacteria are rare in the Baltic Sea, with mostly no or weak trends. In the southern Baltic Proper, bacterioplankton biomass declined by 3.6% per year and community growth by 0.8% per year between 1988 and 2007, mainly attributed to improved water management and changes in temperature and salinity⁴. Bacterial biomass and growth in the Gulf of Bothnia showed no or only weak trends in 1999-2014⁵; also in deeper water layers⁶. Surface water warming enhanced the risk of infection with human-pathogenic *Vibrio* spp. and increased *Vibrio*-suitable areas in the Baltic Sea⁷.



What can be expected?

Continued eutrophication together with a longer algal growth season and higher sea surface temperature, will intensify bacterially mediated transformation of organic matter, CO₂-production, and oxygen consumption in the Baltic Sea^{8,9}. Counteracting this, increased riverine dissolved organic carbon (DOC) discharge due to precipitation will hamper light and thereby algal productivity, while maintaining bacterial production¹⁰. No reliable modelling of these processes is currently available to help project the net outcome for, e.g., marine oxygen consumption. Warming and extended heatwaves will increase the risk of infection of humans by pathogenic bacterioplankton like the *Vibrio*⁷.



Other drivers

Light (influenced by, e.g., cloudiness and turbidity) influences algal growth and the production of bacterial substrates. Light also cleaves refractory compounds to usable food for bacterioplankton¹¹. Environmental toxins and pharmaceuticals may also influence bacterioplankton, either by being food for bacterioplankton¹² or by hampering bacterial growth.



Knowledge gaps

Lack of long time-series of bacterial growth, abundance, and composition make the projection of long-term effects uncertain. Few biogeochemical models coupled to meteorology include microbial activity properly, making net outcomes of large-scale and long-term effects difficult to foresee. Rapid bacterial adaptation to altered conditions, occurring on both population and genetic levels, is often associated with evolutionary changes in functions, adding uncertainty. International harmonization of methodology in microbial ecology is further of importance for building reliable knowledge.



Policy relevance

Microbial mechanisms are fundamental for the carbon balance, oxygen status and CO₂ production, and crucial in understanding the effects of climate change and biogeochemical cycles in general. Efforts to reduce greenhouse gas emissions, stop clearing of forests, and re-forest agricultural land are ongoing, but insufficient. Monitoring of bathing water quality is ongoing but needs improvement. Global actions assessed to be a long-term remedy, for example binding CO₂ by fertilizing algae, would likely lead to adverse effects on Baltic Sea oxygen status. Since no means of direct human control of microbial abundance and activity is currently available, the microbial community is not managed through any policies.





Benthic habitats

Linked parameters:
Water temperature, Sea ice, Salinity and saltwater inflows, River run off, Carbonate chemistry, Sea level, Waves, Oxygen, Pelagic and demersal fish, Coastal and migratory fish, Waterbirds, Non-indigenous species, Marine protected areas, Nutrient concentrations and eutrophication, Ecosystem function, Blue carbon storage capacity

Authors
Antonia Nyström Sandman, Aquabiota Water Research, Sweden
Markku Viltasalo, Finnish Environment Institute, Finland
Jesper Philip Aagaard Christensen, Århus University, Denmark
Mats Blomqvist, Hålok AB, Sweden
Christina Halling, Swedish University of Agricultural Sciences, Sweden
Ari Laine, Parks and Wildlife Finland, Finland
Georg Martin, University of Tartu, Estonia

Biota and ecosystems

Links to main policies:
HELCOM Baltic Sea Action Plan
EU Water Framework Directive (WFD)
EU Habitats Directive (HD)
EU Birds Directive (BD)
EU Biodiversity Strategy
UN Convention on Biological Diversity (CBD)
UN Sustainable Development Goals (SDGs)
EU Marine Strategy Framework Directive (MSFD)



Description

Benthic habitats in the Baltic Sea are characterized by a mixture of species of marine and freshwater origin¹. In the deep benthic areas, communities are dominated by only a few invertebrate species, whereas in the shallow photic areas, various macroalgae and vascular plants provide food and shelter for a large number of invertebrates and fish at both hard and soft bottoms. Climate change may affect the composition, abundance, biomass and spatial distribution of benthic species and habitats, with potential loss of biodiversity and ecosystem functions as a result².



What is already happening?

Benthic soft substrate communities in large parts of the Baltic Sea have drastically changed during the past decades, with amphipods decreasing³, Baltic clam *Limecola balthica* increasing, and the non-indigenous polychaete *Marenzelleria* becoming dominant⁴. Changes have been explained to some degree by abiotic factors such as temperature, fluctuations in salinity and oxygen, and precipitation and runoff related changes in pelagic food webs^{4,5}. Decreasing amount of sea ice has consequences for stratification, nutrient dynamics, and hence benthic communities. Despite decreasing nutrient loads, hypoxic areas continue to prevail in the central Baltic Sea⁶ and increase in the coastal zone⁷, causing loss of communities and ecosystem functions⁸⁻¹¹.



What can be expected?

Many Baltic species exist on their geographical distribution limit, and small fluctuations in temperature and salinity can have a large impact on, e.g., bladderwrack, blue mussel and eelgrass¹²⁻²⁰. Increasing temperature affects species turnover rates and physiology^{12,21-24}. In coastal ecosystems, increased precipitation and runoff might cause salinity fluctuations²⁵, affecting species reproduction and survival²⁶. Sea-level rise²⁷ will change prerequisites for important environments like shallow coastal habitats. In the presently oxic areas, macrozoobenthos productivity will decrease if oxygen conditions deteriorate^{28,29}. In areas with increasing riverine load of dissolved organic carbon (DOC), pelagic primary production may decrease, affecting the benthic system^{21,25}.



Other drivers

Eutrophication has a major impact on the benthic ecosystem, mainly through enhanced primary production causing increased water turbidity and decreasing bottom-water oxygen³⁰. Nutrient input is likely to increase with increasing precipitation, especially in the northern Baltic Sea³¹, and in combination with impacts of increasing temperature, changes can be anticipated in all trophic levels. However, success of nutrient load reductions may have a larger effect on the benthic ecosystem than climate change alone^{28,32}. Also, introductions of non-indigenous species can cause changes in marine biodiversity³³ and ecosystem functions³⁴⁻³⁶. Reduction of predatory fish might affect functionality of benthic habitats through trophic cascades³⁷⁻⁴⁰.




Knowledge gaps

Salinity decline has been hypothesized to be the major driver of geographic species shifts, but according to recent regional climate modelling the magnitude of change is uncertain^{41,42}. Effects of climate change on benthic habitats are difficult to project, due to cumulative and changing impacts of stressors, as well as confounding food web interactions^{21,43}. The interactions of climate change with other stressors are not well known, nor is the capability of organisms to adapt to climate change. For instance the keystone species bladderwrack has in some studies been shown to adapt to climate change⁴⁴⁻⁴⁶, while others have suggested that the species cannot keep pace with the projected salinity change, with large effects on biodiversity and ecosystem functioning^{19,36}.



Policy relevance

The Marine Strategy Framework Directive requires assessing the status of benthic habitats^{46,47}, and the cumulative effects of climate change and other pressures, such as eutrophication, on biodiversity and ecosystem functions should be considered. According to the Habitats Directive, the extent of adverse effects cannot exceed a certain proportion of the habitats, and member states shall establish a coherent network of Natura 2000 areas to secure ecosystem structure and function⁴⁹. If climate change causes community changes, conservation targets need to be updated and the network of MPAs should be adapted to take projected changes into account⁵⁰. Also, climate change needs to be incorporated in marine spatial planning, at appropriate spatial and temporal scales⁵¹.



Coastal and migratory fish

Linked parameters:
Water temperature, Sea ice, Salinity and saltwater inflows, River run-off, Benthic habitats, Waterbirds, Marine mammals, Marine protected areas, Ecosystem function, Fisheries

Authors
Örjan Östman, Jens Olsson, Noora Mustamäki and Rahmat Naddafi, Swedish University of Agricultural Sciences (SLU), Sweden
Meri Kallassuo, Sanna Kuningas and Antti Lappalainen, Natural Resources Institute Finland (Luke), Finland
Lauri Saks, University of Tartu, Estonia

Biota and ecosystems

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goal 14
UN Convention on Biological Diversity
EU Green Deal
EU Marine Strategy Framework Directive (MSFD)
EU Water Framework Directive (WFD)
EU Habitats Directive
EU Common Fisheries Policy (CFP)
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Biodiversity Strategy



Description

Fish of freshwater origin dominate most Baltic coastal areas, some preferring warm (perch, cyprinids) and others cold waters (salmonids, burbot)¹. These species often migrate back to their natal spawning ground for spawning, resulting in many local populations that adapt to local conditions. Small scale environmental variations, local fishing pressure, habitat availability, and food web interactions influence their reproduction, recruitment, growth, and mortality.



What is already happening?

Higher water temperature has improved the reproduction of many spring and summer spawners²⁻⁹.

In contrast, the reproduction of autumn-spawners, e.g., vendace and whitefish, is disfavoured by warm winters and their distribution decreases with less ice cover and higher winter temperatures¹⁰⁻¹³.

Species preferring warm waters have become more common relative to winter-spawning species¹⁴.

Migratory anadromous species, like salmon, return earlier to rivers after warm winter/spring. However, high water temperature in autumn and winter seems to lower the survival of salmon migrating back to the sea¹⁵⁻¹⁹.



What can be expected?

Earlier spawning, faster egg, and larval development, increased larval survival of spring spawning freshwater coastal fish species^{6-9,20-22} (*).

Earlier migration from nursery habitats⁶ may influence food web interactions with negative effects on piscivorous species²³ (*).

Reproduction of autumn-spawning migratory fish is expected to decrease with increasing temperatures, and spawning areas reduced if ice cover decreases further¹¹⁻¹³.

The effect of water temperature on body growth differs among species and size-classes: growth is generally expected to increase for small but not for large fish^{3,10,16,17,21,22}.

Possible brownification of coastal waters may decrease body growth²⁴.



Other drivers

Anthropogenic pressures, such as eutrophication, fishing, and habitat exploitation, affect fish in coastal areas .

Pharmaceutical residues and plastics might negatively affect fish locally.

Increased cormorant and seal populations consume substantial amounts of coastal fish²⁵, but the impact on fish populations is disputed²⁶.

Migratory anadromous fish are affected by a similar set of pressures as coastal fish, and in rivers also by altered hydrological regimes, migration barriers caused by dams, and increased sedimentation due to land-use changes in the drainage area¹⁹.



Knowledge gaps

Indirect and interactive effects of different natural and anthropogenic pressures in combination are poorly studied. To identify causal relationships, modelling based on monitoring data in combination with experimental studies is needed.

The effects of some expected climate induced changes, e.g., shrinking ice cover and browner waters, on coastal and migratory fish stocks are poorly studied.

The importance of extreme weather events under climate change for fish population development and status is furthermore insufficiently studied. Follow-up studies after extreme weather events (like heatwaves, and ice-free winters) are of key importance for understanding the recovery and resilience of fish populations and communities.



Policy relevance

Coastal and migratory fish are key elements for Baltic Sea coastal food web structure and function, and fundamental for small scale coastal commercial and recreational fisheries. Current measures to protect and restore coastal and migratory fish populations hardly ever target and consider climate change effects. Targeted short-term actions, e.g., temporary or spatial closures, could help affected fish populations to recover from extreme weather events. Future management should include climate change effects in status assessments and management plans, targets, and measures to acknowledge and mitigate climate related effects.

*) Expected to be caused by warmer temperatures.



Pelagic and demersal fish

Authors
Örjan Östman, Jens Olsson, Noora Mustamäki and Rahmat Naddafi, Swedish University of Agricultural Sciences (SLU), Sweden
Meri Kallassuo, Sanna Kuningas and Antti Lappalainen, Natural Resources Institute Finland (Luke), Finland

Biota and ecosystems

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goal 14
UN Convention on Biological Diversity
EU Green Deal
EU Marine Strategy Framework Directive (MSFD)
EU Habitats Directive
EU Birds Directive
EU Common Fisheries Policy (CFP)
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Biodiversity Strategy

Linked parameters:
Water temperature, Sea ice, Salinity and saltwater inflows, Stratification, Oxygen, Benthic habitats, Waterbirds, Marine mammals, Marine protected areas, Ecosystem function, Fisheries



Description

Fish of marine origin, such as cod, herring, sprat, and flatfishes (flounder, plaice, turbot, and dab), dominate pelagic and demersal habitats of the Baltic Sea¹. These species occur in large, often internationally managed, stocks.

Currently, sticklebacks make a significant part of the pelagic fish biomass.

Temperature impacts the recruitment (successful reproduction and survival of the offspring), body growth and mortality of pelagic and demersal fish, resulting in changes in spatial and seasonal distributions.



What is already happening?

Increasing temperatures and hypoxic conditions have impaired reproduction, reduced feeding areas as well as quality of food, resulting in decreasing distributions of flatfish, herring and cod, and reduced growth and body condition of cod²⁻¹⁰.

Increasing temperature favours stickleback^{11,12}.

Periods with low salinity are connected to lower recruitment of several flatfishes, herring, and cod¹³⁻¹⁸, and lower abundance and lipid content of zooplankton prey for herring and sprat¹⁹⁻²², resulting in lower body growth, condition, and abundance¹⁹⁻²³.

Recruitment of sprat is higher in warmer waters after winters with low ice cover but opposite for herring^{24,25}.



What can be expected?

Increasing water temperature causes earlier spawning, shorter development, and increased recruitment of sprat^{24,26,27},

and increasing larval growth of herring, sprat, and flatfish, and body growth of adult sticklebacks^{11,26,28,29}. Herring and cod recruits may miss optimal temperature windows resulting in lowered recruitment^{25,26,28,30,31}.

Increasing temperature, especially if the halocline shifts upwards and nutrient loads are not reduced, may reduce oxygen in water and sea bottom. This will lead to reduced reproduction and feeding areas, increased food competition, and dependency on shallow areas for cod and flatfishes^{5,32}.

If salinity decreases, this may also reduce recruitment, abundance, and distribution of flatfish, sprat, and cod^{2,6,8,15,28,33}.



Other drivers

Impacts of multiple drivers on offshore fish communities are perceivable^{34,35}. High nutrient discharges have resulted in enhanced hypoxic conditions affecting many fish species negatively⁵⁻¹⁰, but also benefitting others^{36,37}.

Nutrient loads have decreased since the 1980s, but the response in nutrient concentrations is slow and also affected by runoff and climate related variables such as temperature and stratification³¹.

Fishing strongly affects cod, herring, and sprat. Harmful substances, marine litter, and pharmaceutical residues might have negative impacts on individuals while effects on populations appear to be small, yet uncertain. Food-web interactions (competition/predation/food quality) among populations are evident.

Vitamin deficiency (e.g., thiamine) may impact fish species.



Knowledge gaps

Indirect and interactive effects of climate parameters and other pressures on fish need to be better studied³⁸⁻⁴¹. To explain causal relationships, modelling of monitoring data in combination with experiments is required. Furthermore, impacts of changes related to climate, like ice cover, brownification, and acidification, are poorly studied in the Baltic Sea.

The importance of average changes relative to extreme weather events (e.g., heatwaves vs. average temperature) are poorly studied. There is a need to analyse monitoring data before, during, and after extreme events, supplemented with experiments and long-term data to understand the recovery and resilience of fish species and communities after extreme weather events.



Policy relevance

Demersal and pelagic fish are key elements for Baltic Sea offshore food web structure and function, and offshore fisheries. Management of demersal and pelagic fish, e.g., quotas, fishing closures and protected areas, takes historic changes in stock productivity into account but does not consider predicted climate change effects. Furthermore, management of these stocks needs to be adaptive to react to long-term effects of climate change. Targeted short-term actions, e.g., temporary, or spatial closures, could help affected fish populations to recover from extreme weather events. Targets and measures in future management plans need to consider long-term impact of climate change on fish populations and communities.



Waterbirds

Authors
Volker Dierschke, Gavia EcoResearch, Germany
Morten Frederiksen, Århus University, Denmark
Elie Gaget, Department of Biology, University of Turku, Finland
Diego Pavón-Jordán, Department of Terrestrial Ecology, Norwegian Institute for Nature Research (NINA), Norway

Biota and ecosystems

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goal 14
UN Convention on Biological Diversity
EU Green Deal
EU Marine Strategy Framework Directive (MSFD)
EU Maritime Spatial Planning Directive (MSP)
EU Birds Directive, EU Habitats Directive
EU Common Fisheries Policy (CFP), EU Common Agricultural Policy (CAP)
EU Strategy for the Baltic Sea Region (EUSBSR), EU Biodiversity Strategy
AEWA Agreement
Ramsar Convention



Description

In total, around 100 waterbird species use the marine area and the coastal habitats of the Baltic Sea for breeding, staging during migration, moulting, and/or wintering. They have different roles in the food web, being predators of various fish and invertebrates and foraging in different habitats as well as providers of multiple ecosystem services¹.



What is already happening?

Many waterbird species have shifted their wintering range northwards²⁻¹⁰.

Some waterbird species migrate earlier in spring^{11,12}.

Effects of warming water temperature are inconsistent, because both positive and negative effects have been found regarding foraging conditions and food quality, including invertebrate prey and prey fish species¹³⁻¹⁵.

As most Baltic waterbirds are migratory, they are affected by climate change also outside the Baltic Sea, for example during breeding in the Arctic and migration and wintering between southern Europe and western Africa¹⁶.



What can be expected?

The northward distributional shifts are expected to continue^{9,10}.

Effects on waterbird food will be manifold, but consequences are difficult to predict¹⁶.

Rising sea level and erosion are expected to influence the availability of breeding habitats^{17,18} and rising sea level may reduce breeding success due to flooding of the breeding and wintering foraging habitats.



Other drivers

In the Baltic Sea, waterbird populations are increasingly impacted by human activities during the breeding season, such as recreation^{19,20} and introduced predators (e.g., American mink²¹⁻²³) and the wintering season (hunting^{24,25}, fishing^{26,27}, ship traffic^{28,29}, and offshore wind farms^{30,31}).

Eutrophication and fishing strongly influence foraging preconditions for waterbirds^{3,32,33}.



Knowledge gaps

Food web complexity and interacting natural and anthropogenic effects make it difficult to isolate the effects of climate change on waterbird abundance. For some well-studied species, these effects are demonstrated, but in most cases there is a lack of understanding especially of how phenological mismatches will affect breeding and wintering waterbirds across functional groups and life histories.



Policy relevance

Waterbirds are an important part of the marine food web in the Baltic Sea. Changes in the phenology and distribution of waterbirds may require adapting environmental conservation policies, notably by extending and adjusting the networks of protected areas and by supporting their management^{17,34} with robustly designed monitoring of sites and populations⁹. Hunting regulations need to be adjusted in space and time to account for distributional and phenological shifts, i.e., regulations need adjustment where climate change has caused increased waterbird occurrence with therefore higher importance of the respective locations.





Marine mammals

Linked parameters:
Sea ice, Sea level, Waves, Pelagic and demersal fish, Coastal and migratory fish, Marine protected areas, Offshore wind farms, Ecosystem function, Shipping, Fisheries, Aquaculture, Marine and coastal ecosystem services

Authors
Anders Galatius, Department of Bioscience, Århus University, Denmark
Markus Ahola, Swedish Museum of Natural History, Stockholm, Sweden
Ida Carlen, Coalition Clean Baltic
Antti Halkka, University of Helsinki, Faculty of Biological and Environmental Sciences, Helsinki, Finland
Mart Jüssi, ProMare, Estonia

Biota and ecosystems

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goal 14
UN Convention on Biological Diversity
EU Green Deal
EU Marine Strategy Framework Directive (MSFD)
EU Maritime Spatial Planning Directive (MSP)
EU Habitats Directive
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Biodiversity Strategy
ASCOBANS Jastarnia Plan



Description

Three seal species and one cetacean live in the Baltic Sea: ringed seal (*Pusa hispida*), grey seal (*Halichoerus grypus*), harbour seal (*Phoca vitulina*) and the harbour porpoise (*Phocoena phocoena*).

Being at the top of the marine food web, these predators are sensitive to changes throughout the ecosystem, including those related to climate. Furthermore, the extent and quality of sea ice are important particularly to the ice-breeding ringed seals and also the facultatively ice-breeding grey seals. In some areas, seals are dependent on low-lying haul-outs (land areas for resting, breeding, foraging etc.).



What is already happening?

Both the ice cover and duration of the ice season have already been markedly reduced¹⁻⁸. The changes are most prominent in the southern areas where the duration of ice cover during breeding season of ringed and grey seals has increasingly often been either too short or completely lacking⁸. This diminishes breeding success of ringed seals.

To a lesser extent, the breeding success of grey seals is diminished, particularly in the southern areas.

See 'Sea ice and extreme events' for further quantification.



What can be expected?

Projected reduction of sea ice^{4,9-11} for ringed seal and of snow for pupping lairs will impact ringed and grey seal breeding. Disappearance of ringed seals from southern areas is possible and transfer of grey seal breeding to land sites probable.

Sea level rise^{12,13} causing flooding of haul-outs in the southern Baltic may force out breeding seals. This will likely cause reduction of harbour and grey seal occurrence to foraging individuals.

Changes in temperature and stratification, prey distribution, quality and quantity will affect all marine mammals, but aggregated effects on their abundance and distribution are unpredictable.



Other drivers

Ice-breaking and winter shipping may worsen effects of reduced ice on seal breeding^{14,15}.

Bycatch affects marine mammals¹⁶⁻¹⁸.

Anthropogenic disturbance affects seal distribution and recruitment^{19,20}

Epidemics can reduce seal abundance and possibly distribution²¹.

Ecosystem changes and overfishing influence prey availability^{22,23}

Pollutants impair marine mammals' immune function and fertility^{24,25}

Underwater noise may for all species cause injury and displacement from habitats and disturb natural behavior, and for harbour porpoise interfere with echolocation^{16,26}.



Knowledge gaps

Seal and porpoise foraging distribution and the relation of the former to haul-out sites are not well known.

While the reduced breeding success of grey seals on land relative to ice has been studied²⁷, the absolute dependency on ice for successful breeding of ringed seals has not been sufficiently assessed.

Land-breeding of grey seals is not surveyed in most Baltic countries.

Breeding success of ringed seals during favourable ice conditions, even under current conditions, is poorly known. Ringed seals are adapted to ice-breeding. Land breeding attempts are known from extremely poor ice years, but successful land breeding has not been documented for the species.

The aggregate effects of climate-related ecosystem changes on marine mammals have not been modelled.



Policy relevance

Marine mammals are top predators in the Baltic Sea and important as sentinels of ecosystem health²⁸ and as top-down regulators of the ecosystem. Direct effects of climate change are mostly impossible to address locally. Artificial lairs may mitigate breeding failure and haul-outs could potentially be artificially sustained above water.

Seasonal shipping restrictions may reduce impacts on seal breeding. There are currently no actions to directly mitigate climate change effects on marine mammals, but measures are in place that mitigate human disturbance, pollution and bycatch, and hunting is limited or prohibited.

Further mitigation of pressures will improve climate change resilience of the populations. Consideration of effects of climate change on the populations should be integrated in national management plans.



Non-indigenous species

Linked parameters:
Water temperature, Salinity and saltwater inflows, Carbonate chemistry, Oxygen, Benthic habitats, Marine protected areas, Nutrient concentrations and eutrophication, Ecosystem function, Shipping, Aquaculture



Description

Non-indigenous species, NIS, are not native to the geographic region of interest, but transferred there by human activities. Ship ballast water and hull fouling are the main vectors for their transfer¹⁻⁴. NIS are more often found in the coastal zone than the open sea⁵⁻⁸ and ports are hot spots for their introduction⁹⁻¹². Some 170 NIS are recorded from the Baltic Sea^{13,14}, with more than 70 permanently established. Most NIS show unique responses to changes in the environment, hence changes caused by climate change will be species-specific and further modified by complex interactions with native species.



What is already happening?

So far, no invasion can be confidently attributed to climate change. Environmental changes caused by climate change may increase stress on native species¹⁵⁻¹⁸ and favour some NIS, increasing their ecological impact¹⁹⁻²¹.

Climate driven shifts in species boundaries towards higher latitudes affect potential of new introductions to the Baltic Sea. Changes in salinity regime affect distribution and establishment of NIS depending on their origin and tolerance to salinity²².



What can be expected?

Higher temperature and a possible salinity decrease can increase recruitment and growth of certain NIS, e.g., dreissenid mussels, several freshwater crustaceans, and the round goby²¹⁻³⁵.

Changes may first be seen in estuaries, where the contribution of NIS is already high^{36,37}.

If oxygen deficiency increases in warmer coastal waters, it may constrain the growth of the round goby³⁸, but more tolerant species, like the polychaete worm *Marenzelleria* spp., may increase³⁹ and change sediment nutrient fluxes and resuspend contaminated sediments⁴⁰.

Warmer winters will facilitate survival for introduced warm-water species^{41,42}.



Other drivers

As the vast majority of NIS arrive with ships, the main driver of biological invasion is the occasional, unintended, and unpredictable introduction of organism into Baltic Sea ecosystem. Also, aquaculture has a significant impact on the arrival of NIS¹⁴. Eradication after introduction is mostly impossible and the main focus must be on preventing any NIS from arriving in the first place. Anthropogenic disturbances, like eutrophication and habitat degradation, interact with biological invasions by affecting the conditions for NIS establishment.



Knowledge gaps

Most NIS are ecologically unique, and it is therefore important to predict how invasive species will behave and interact in a new environment. It is important to identify the potential threats they pose to native species and ecosystem functions. Planning of management measures is challenged by the high variability in species characteristics and the unpredictable nature of new introductions.



Policy relevance

Once NIS are established, they are practically impossible to remove. Policies are thus focused on preventive measures. Targets for minimizing adverse effects of NIS on biodiversity and ecosystems have been set in the EU Marine Strategy Framework Directive, the EU Invasive Alien Species Regulation, and the HELCOM Baltic Sea Action Plan but reaching these goals will be difficult if climate change promotes successful establishment of NIS. Policy focus should be on preventing new introductions, for example by implementing the regulations related to aquaculture, Ballast Water Management Convention, and by working to manage biofouling on ship-hulls (commercial as well as recreational).



Authors
Rahmat Naddafi, Swedish University of Agricultural Sciences (SLU), Sweden
Katarzyna Spich, National Marine Fisheries Research Institute (NMFRI), Poland
Isa Wallin, Swedish University of Agricultural Sciences (SLU), Sweden
Örjan Östman, Swedish University of Agricultural Sciences (SLU), Sweden
Maiju Lehtiniemi, Finnish Environment Institute (SYKE), Finland
Ari Laine, Parks & Wildlife Finland (Metsähallitus), Finland

Biota and ecosystems

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goals 2 and 14
UN Convention on Biological Diversity
EU Green Deal
EU Marine Strategy Framework Directive (MSFD)
EU Water Framework Directive (WFD)
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Regulation on invasive alien species
EU Biodiversity Strategy



Marine protected areas

Linked parameters:

Benthic habitats, Pelagic and demersal fish, Coastal and migratory fish, Waterbirds, Marine mammals, Non-indigenous species, Coastal protection, Offshore wind farms, Shipping, Tourism, Fisheries, Aquaculture, Blue carbon storage capacity, Marine and coastal ecosystem services



Description

Marine protected areas (MPAs) are intended to conserve ecologically significant parts of the marine and coastal environment, including biological and genetic diversity and ecological functions. Biodiversity, including genetic diversity, is needed for species' adaptation and long-term survival under changing environmental conditions¹. Sufficiently sized and adequately located MPAs will likely help marine organisms adapt to climate change and increase their survival by reducing impacts of other human pressures². In 2021, HELCOM MPAs covered 13.28% of the Baltic Sea^{3,4}.

The effect of climate change can be evaluated by assessing consequences to MPA conservation values and function based on benthic habitats, fish stocks, birds, and seals.



What is already happening?

● In comparison to other marine areas, the Baltic Sea is prone to climate change related warming and oxygen depletion⁵. Until now, negative effects of climate change on the Baltic Sea ecosystem have already become apparent through pelagic regime shifts (i.e. persistent change in the ecosystem)⁵. Milder winters with shorter ice period and reduced ice cover restrict breeding habitats for the ringed seal (see "*marine mammals*").

● Simultaneously, northward distributional shifts of birds may increase the importance of MPAs as overwintering areas (see "*waterbirds*"). Habitat change through higher temperatures and oxygen depletion (related to eutrophication) may harm fish stocks and benthic communities, also impairing MPA conservation values.



What can be expected?

● If sea ice is reduced, while water level, erosion, and flooding increase, some MPAs may lose parts of their function as breeding and feeding sanctuaries for marine mammals and waterbirds^{6,7}.

● Distributional changes of biological communities caused by climate change may impair the function of MPAs and, together with other anthropogenic pressures, prevent MPAs from meeting their objectives⁸⁻¹⁴.



Other drivers

Cumulative pressures caused by a variety of human activities both inside and outside MPAs are crucial drivers of ecosystem damage and biodiversity loss in the Baltic Sea. Intensive shipping and fishing, sand and gravel extraction, offshore installations, as well as inputs of nutrients and hazardous substances from land represent major threats to the whole Baltic Sea ecosystem and its adaptability to climate change. Pressures are further exacerbated by the limited water exchange.



Knowledge gaps

There is no commonly agreed method to assess the ecological and management effectiveness of MPAs, which impedes evaluations and optimisation of MPAs as a management tool. Moreover, totally protected no-access areas, which would provide reference sites for the determination of a baseline for natural conditions, are lacking, which also complicates defining objectives of MPAs. Knowledge gaps also exist in understanding connectivity of areas, affecting the ecological coherence of the MPA network^{15,16}.



Policy relevance

Effectively managed MPAs can mitigate impacts of climate change to conserve biodiversity and healthy, resilient marine ecosystems, which also act as carbon sinks¹⁷. International and national policies would benefit from fostering a change in reasoning behind MPAs, from protecting threatened species and biotopes towards securing functional diversity and biodiversity and ensuring ecosystem services. As of 2021, HELCOM supports a network of 177 MPAs, which could act as a minimum buffer for climate change resilience. However, an expansion of the HELCOM MPA network, with climate refuges in which food web perspectives and genetic diversity are considered¹, is needed.

Authors

Ari Laine, Metsähallitus Parks & Wildlife Finland, Finland
Orjan Östman, Swedish University of Agricultural Sciences (SLU), Sweden
Dieter Boedeker, German Federal Agency for Nature Conservation (BfN), Germany
Gesine Lange, German Federal Agency for Nature Conservation (BfN), Germany

Biota and ecosystems

Links to main policies:

HELCOM Baltic Sea Action Plan
UN Sustainable Development Goal 14
UN Convention on Biological Diversity
EU Green Deal
EU Marine Strategy Framework Directive (MSFD)
EU Water Framework Directive (WFD)
EU Maritime Spatial Planning Directive (MSP)
EU Habitats Directive
EU Birds Directive
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Biodiversity Strategy



Nutrient concentrations and eutrophication

Linked parameters:

Water temperature, Stratification, Precipitation, Riverine nutrient loads and atmospheric deposition, Carbonate chemistry, Oxygen, Benthic habitats, Pelagic and demersal fish, Coastal and migratory fish, Non-indigenous species, Ecosystem function, Aquaculture, Blue carbon storage capacity, Marine and coastal ecosystem services



Description

Nitrogen and phosphorus pools are controlled by inputs from land and atmosphere and modified by biogeochemical transformations. Both these nutrients cycle intensely between the water column, biota, and bottom sediment. Nitrogen fixation and denitrification act as major biogeochemical sources and sinks, whereas phosphorus tends to accumulate in bottom sediments. Furthermore, considerable amounts of nutrients are exported to the North Sea^{1,2}.

Bottom oxygen conditions regulate denitrification rates and the distribution of phosphorus between sediment and the water. Higher nitrogen loss and phosphorus release from sediments occur when hypoxia expands^{3,4}.

Dissolved inorganic nitrogen is described in this text with the abbreviation DIN and dissolved inorganic phosphorus with the abbreviation DIP.



What is already happening?

● Climate change impacts on nitrogen and phosphorus pools cannot yet be separated from other pressures. Effects of warming and sea level rise are masked by changes in nutrient loads and bottom water oxygen levels⁴. Nitrogen concentrations have decreased in most Baltic Sea basins since 1990, but phosphorus pools have fluctuated without trend⁵.

● Eutrophication has made shallow areas with restricted water exchange more prone to hypoxic events⁶.

● Nutrients liberated from sediments during the hypoxic events fuel summer phytoplankton blooms⁷. Changes in stratification and cloud cover⁸ currently prolong the phytoplankton growth season, with earlier spring onset and extended autumn blooms⁸, without clear effect on nutrient concentrations⁸.



What can be expected?

● The development of nutrient loads will dominate future nutrient concentrations^{9,10}, with warming expected to reduce near-bottom oxygen by increasing internal nutrient cycling and by strengthening thermal stratification⁹⁻¹¹.

● A decline in the DIP pool is projected¹² (*). DIP surface concentrations in the Baltic Proper will decrease with BSAP load scenarios and slightly increase with current load scenarios while surface DIN concentrations remain unchanged under both scenarios⁹.

● In the Gulf of Finland and Bothnian Sea, it is expected that DIN levels will increase with both load scenarios and DIP changes will be similar to the Baltic Proper⁹.

● nitrogen-fixing cyanobacteria blooms are expected to expand¹³⁻¹⁵ (**).



Policy relevance

High nutrient loads cause eutrophication, which is a major problem in the Baltic Sea. Nutrient input is primarily coming from agriculture and fertilizer use on land. Eutrophication is central in the HELCOM Baltic Sea Action Plan²¹, the EU Marine Strategy Framework Directive²² and the EU Water Framework Directive²³, and all these policies aim to reduce eutrophication in the Baltic Sea even more than the already achieved reductions.

*) When climate change effects are taken into account, the Baltic Sea Action Plan (BSAP) and current load scenarios project, respectively, a 50% and 25% decline in the Baltic DIP pool until 2070-2100¹².
**) Without nutrient load reductions, nitrogen-fixing cyanobacteria blooms are expected to expand.



Ecosystem function

Linked parameters:
Water temperature, Sea ice, Solar radiation, Salinity and saltwater inflows, Stratification, Carbonate chemistry, Riverine nutrient loads and atmospheric deposition, Oxygen, Microbial community and processes, Benthic habitats, Pelagic and demersal fish, Coastal and migratory fish, Waterbirds, Marine mammals, Non-indigenous species, Nutrient concentrations and eutrophication, Coastal protection

Authors
Örjan Östman, Rahmat Naddafi, Jens Olsson,
with input by Agnes ML Karlsson and Magnus Huss,
Swedish University of Agricultural Sciences (SLU), Sweden

Biota and ecosystems

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goal 14
UN Convention on Biological Diversity
EU Green Deal
EU Marine Strategy Framework Directive (MSFD)
EU Water Framework Directive (WFD)
EU Maritime Spatial Planning Directive (MSP)
EU Habitats Directive
EU Birds Directive
EU Common Fisheries Policy (CFP)
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Biodiversity Strategy



Description

Baltic Sea ecosystems provide an array of ecosystem functions related to, for example, nutrient and carbon circulation, biomass production, and regulation.

Climate-related factors structure Baltic Sea food webs both through top-down (predation) and bottom-up (biomass production) processes¹⁻⁷ that are fundamental for ecosystem functioning.

Climate change will likely impact several processes related to food web interactions, nutrient recycling, and ecosystem properties.



What is already happening?

- Long-term eutrophication has increased primary production and during the last decades more frequent algal blooms are observed during warmer years. This causes increased decomposition and oxygen-depleted bottom sediments^{1-4,8-10}.
- Changes in ice cover, cloudiness, and wind condition in spring may have resulted in changed timing of algal blooms, affecting benthic productivity^{1,4}.
- Changes in hydroclimatic conditions in combination with fishing and eutrophication have resulted in a shift from larger to smaller zooplankton^{8,9}, and stronger impact of nutrients on ecosystem structure (bottom-up control) and reduced the regulatory capacity of predators on ecosystem structure (top-down control) in both pelagic and coastal Baltic Sea food webs^{5-7,11-13}.



What can be expected?

- Warmer water may increase pelagic and benthic primary production^{1,2,8,9,14}.
- Unless nutrient loads are reduced, oxygen levels in the water and close to the seabed will decrease¹⁵. Responses at higher trophic levels will differ among organism groups^{8,16-19}.
- If salinity decreases, this will likely affect the species composition of zooplankton and fish, and the associated functions, e.g., predation rates^{9,17,20,21}.
- If inflow of dissolved organic matter increases, this may increase benthic production, and increase bacterial production over phytoplankton production. Reduced light conditions may reduce total primary production of benthic and pelagic food webs^{22,23}.



Other drivers

- Fishing is a strong pressure on some fish species and results in reduced natural control of their prey^{5-7,11-13}.
- Nutrient concentrations are main drivers of biomass production, causing negative impacts on oxygen levels and water clarity that can severely worsen climate change-related effects on ecosystem functioning^{8,9,15,24}.
- Seals and cormorants have increased in the Baltic Sea, but food web effects are poorly known and uncertain⁶. Toxins, marine litter, pharmaceutical residues and vitamin deficiency (e.g. thiamine) have negative impacts on individuals of different functional groups, but the ecosystem effects at the Baltic Sea scale are uncertain.



Knowledge gaps

Several parameters are intercorrelated and there are potentially indirect and interactive effects of for example oxygen, salinity, and temperature on ecosystem functioning^{6,8,9,18,19,24}.

The magnitude and interactive effects of climate change relative to other human pressures are hence important to estimate⁶.

There are knowledge gaps on how changes in Baltic Sea food web structure, resilience and functioning depend on long-term changes in climate relative extreme weather events, like heatwaves. It would be important to analyse monitoring data before, during, and after extreme events, such as the heat waves or low ice cover^{25,26}.



Policy relevance

Ecosystem functions are essential processes structuring ecosystems and food webs, including key ecosystem services to human well-being. Management actions in general focus on populations (fishing/hunting, protection) or inputs (nutrients, toxic compounds) that influence ecosystem functions, but these hardly consider climate change effects. Current management plans need to consider long-term impacts of climate change on ecosystem functions, and how extreme weather events should trigger additional short-term actions to avoid ecosystem regime shifts (level of confidence: medium)^{2,3,9,14,15}. Long-term management plans and measures should consider projected changes in primary production and trophic structure of Baltic Sea ecosystems⁶.

Indirect parameters:
Human use





Offshore wind farms

Linked parameters:

Sea ice, Wind, Waves, Sediment transportation, Coastal and migratory fish, Waterbirds, Marine mammals, Marine Protected Areas, Shipping, Tourism, Fisheries, Blue carbon storage capacity

Links to main policies:
UN Sustainable Development Goals 13 and 14
EU Maritime Spatial Planning Directive (MSP)
EU Strategy for the Baltic Sea Region (EUSBSR)
HELCOM Baltic Sea Action Plan
Renewable Energy Directive (2018/2001/EU)
EU Strategy to harness the potential of offshore renewable energy for a climate neutral future (COM(2020) 741)
EU Biodiversity Strategy



Description

Wind farms are the most significant offshore structures in the Baltic Sea, and, in 2021, account for 10% of European offshore wind energy with the current 2 GW of installed capacity¹. The ambitious scenario for deployed offshore wind power capacity by 2050 has been estimated at 32 GW². The environmental impact of wind farms should be considered case specifically with great caution. They affect many oceanographic processes including downstream turbulence, wave energy, local scour, inflowing currents, and surface upwelling³. The submerged structures may locally alter the structural and functional biodiversity of the benthic system⁴. Turbines affect marine mammals through underwater noise during construction and birds and bats through physical disturbance during operation⁵⁻⁹.



What is already happening?

The world's first offshore wind farm was installed in Vindeby, Denmark, in 1991. Currently, as of 2021, offshore wind farms are found in the waters of four countries: Germany (1,074 MW), Denmark (872 MW), Sweden (192 MW) and Finland (68 MW)¹. Climate change (e.g. changes in ice conditions, wind fields, waves) does not have any major influence on the deployment of offshore structures¹⁰. Investment in offshore renewable energy has been emphasized in the European Green Deal, and a dedicated EU strategy on offshore renewable energy was published in November 2020 proposing ways forward to support the long-term sustainable development of this sector¹¹.



What can be expected?

The European Commission estimates that Europe will need 240–450 GW of offshore wind by 2050, equalling up to 30% of Europe's estimated electricity demand at the time¹². The wind energy industry argues that reaching 450 GW would require the Baltic Sea offshore wind capacity to grow to 83 GW. The latter would suggest the annual rate of consent to increase from 2.2 GW (430 km²) to 3.6 GW (720 km²) per year between 2030 and 2040. The increasing spatial demands, contrasting interests and risks to ecosystems call for environmental impact assessments and marine spatial planning to optimize the use of the sea^{13,14}.



Other drivers

Climate change mitigation is the key driver for offshore wind farm industry. However, primarily drivers other than climate change modify the deployment of these offshore structures. The key parameters regarding the location are water depth (< 50 m), wind conditions (> 7 ms⁻¹) and planning issues². Other drivers include, e.g., investments, industrial and employment dimensions, regional and international cooperation, legal framework, supply chains, technological innovations⁶, and exclusions due to military radar issues¹³.



Knowledge gaps

There is insufficient knowledge on the impact of scale of offshore structures on marine biota. Numerical modelling is not able to predict the effects of large-scale construction, potential cumulative effects of multiple farms, or far-field effects at the coast. Further expansion of offshore wind energy should only take place gradually with adequate environmental assessments. In addition, the effects of the expansion of offshore wind energy on biodiversity must be further studied through comprehensive, continuous, and close-meshed research and monitoring. Observational studies are also necessary to validate the models, and extensive site-specific data collection is necessary to compare any changes to the natural ocean state³.



Policy relevance

Offshore wind is one of the cornerstones of EU's energy and climate targets. The European Green Deal recognizes the offshore wind potential in contributing to a modern, resource efficient and competitive economy. The Commission has published an EU strategy on offshore renewable energy, inviting stakeholders to discuss and take forward the proposed policy actions¹¹. However, the EU and national governments have also committed to protecting ecosystems, which may be at risk due to increasing offshore structures. Hence, a broad political discussion is called for to balance the need for renewable energy with its environmental impacts.



Coastal protection

Linked parameters:

Sea ice, River run-off, Sea level, Wind, Waves, Sediment transportation, Waterbirds, Marine Protected Areas, Ecosystem function, Tourism

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goals 13 and 14
EU Maritime Spatial Planning Directive (MSP)
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Floods Directive
EU Biodiversity Strategy



Description

Baltic Sea coasts are under multi-stressor impact due to climate change and various human activities^{1,2}. The impact varies regionally depending on the material of the coast, and the processes in operation¹. Over decades, various coastal protection structures have been established especially along the soft sedimentary shores at the southern coastline of the Baltic Sea, including groins, bulkheads, seawalls, revetments, breakwaters, sills, and sand fences³. These structures modify natural processes and introduce a more static habitat into a formerly dynamic one, raising concerns about their long-term sustainability and ecosystem benefits⁴, while acknowledging the need for coastal flood protection.



What is already happening?

The soft sedimentary coasts in the south are facing the largest changes¹ and exhibit also the most abundant coastal protection structures. However, there is a growing understanding that “soft interventions” rather than “hard” structures may be a more satisfactory way forward. Ecosystem services provided by, e.g., intertidal wetlands can play a critical role in reducing the vulnerability of coastal communities to rising seas and coastal hazards⁵. There are examples of abandoning traditional “hard” coastal protection measures at the Baltic Sea coastline, such as sand nourishment, to enable a recovery of natural dynamics^{6,7}.



What can be expected?

Coastal protection strategies have to increasingly take into account the effects of climate change⁸. Along the low coasts of the southern Baltic Sea, sea level rise is expected to increase cliff and beach erosion and to increase the supply of sediment to the coastal zone⁹. These effects of climate change are expected to increase societal costs for coastal protection, losses of sediment for coastal rebuilding, losses of valuable natural habitats, and of economic value and property¹⁰. Therefore, there is a need for a wider use of innovative approaches such as the Systems Approach Framework (SAF) as a tool for the transition to sustainable development in coastal zone systems^{11,12}.



Other drivers

Direct human influence often has an impact on coastal processes via changes in land use and land cover, coastal and offshore infrastructure constructions, dumping of material and dredging. Regional demographic development and socio-economy influence the coastal ecosystem, as diverse societal and economic claims need to be integrated into regional spatial planning policies alongside climate change adaptation⁸.



Knowledge gaps

Changes in land use, land cover and infrastructure construction are of crucial importance as they operate reciprocally with sedimentary processes causing unexpected morphodynamic consequences. A regional sediment budget for the southern and eastern Baltic Sea is still to be constructed. This requires interdisciplinary and international collaboration¹. In many parts of the southern Baltic coastline, the key question regarding the existing coastline protection structures is their sustainability and efficiency under changing climate and consequently, their potential replacement, adjustment, or removal procedures⁴.



Policy relevance

Coastal processes and their sustainable management under climate change have extremely high policy relevance globally, and in the Baltic Sea. Coastal protection measures should be nationally or regionally incorporated into integrated coastal zone management plans including physical and ecological parameters, cost-benefit analyses, and administrative and legal structures¹³. Due to the complexity of coastal systems and the lack of precise economic valuations, both land and marine spatial planning usually neglect natural coastal protection and other important ecosystem services², calling for a policy change.



Shipping

Linked parameters:
Sea ice, Riverine nutrient loads and atmospheric deposition, Wind, Waves, Marine mammals, Non-indigenous species, Marine protected areas, Offshore wind farms, Fisheries

Authors
Anna Rutgersson, Uppsala University, Sweden
Stuart Ross, European Community Shipowners' Associations
Jonas Pålsson, Swedish Agency for Marine and Water Management, Sweden

Human activities

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goal 14
EU Green Deal
EU Marine Strategy Framework Directive (MSFD)
EU Maritime Spatial Planning Directive (MSP)
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Biodiversity Strategy
MARPOL Annex VI



Description

The Baltic Sea has been an important route for maritime trade since prehistory and is now one of the busiest maritime areas in the world. In 2017, there were 40,391 passages into the Baltic Sea¹. In 2018, approximately 8,300 vessels operated within the area². Shipping is a carbon-efficient transport medium, but still has adverse effects on air quality, eutrophication, and other aspects of the marine environment³.



What is already happening?

In recent decades, the number and size of ships in the Baltic Sea have increased. Climate has changed, with a shorter ice season and earlier ice break-up^{4,5}, facilitating shipping in usually ice-covered areas. Changes in wind field have so far been small and depend on the time period and area studied⁶. Extreme waves have not changed significantly in strength or intensity⁶. Changes in wind and waves could potentially influence safety and fuel consumption.



What can be expected?

Modelling predicts an annual shipping increase of 2.5% for cargo and 3.9% for passenger traffic in Europe⁷. Less sea ice will require less ice-breaking, but the ice will be more mobile. Wave climate in the northern and eastern Baltic Sea is estimated to become more severe, and icing by freezing sea-spray is expected to become more frequent.

Ports and shipping lanes may need to move location or increase or decrease dredging due to sea level rise and increased sedimentation from coastal erosion and river runoff.



Other drivers

Market changes and new regulations will likely modify future shipping much more than direct climate effects. Particularly, regulations to reduce emissions of CO₂, NO_x, SO_x, and particles will influence ship design and fuel use. The changing climate may influence the transportation pattern of traded goods, as some commodities may start to be produced in new locations around the world. Hence, trade flows will shift.



Knowledge gaps

There is a knowledge gap in how new regulations driven by climate change mitigation efforts will affect the fleet composition, fuel selection, and additional technological development. Thus, the response of future Baltic Sea shipping to changes in climate cannot be fully quantified.



Policy relevance

Shipping is a CO₂ effective way to move goods, but still has a substantial carbon footprint. Member States to the IMO have committed to reduce the total annual greenhouse gas emissions from international shipping by 50% by 2050 (from 2008), and phase them out entirely by 2100⁸. Amendments to the IMO environmental regulations concerning mandatory goal-based technical and operational measures to reduce carbon emissions were adopted in 2021.

Increased shipping in previously iced covered areas may increase environmental pressures, but new regulations on noise and emissions may exclude vessels from sensitive marine areas. Establishment of offshore windfarms should be taken into account in marine spatial planning. The environmental impacts of shipping need to be better compared and prioritized with industry on land, including land transportation.



Tourism

Linked parameters:
Air temperature, Water temperature, Sea ice, Solar radiation, Precipitation, Sea level, Wind, Waves, Sediment transportation, Marine Protected Areas, Coastal protection, Offshore wind farms, Marine and coastal ecosystem services

Authors
Kari Hyytiäinen, Department of Economics and Management, University of Helsinki, Finland
Jarkko Saarinen, Geography Research Unit, University of Oulu, Finland
Janika Laht, Climate Department, Ministry of the Environment, Estonia

Human activities

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goal 14
EU Green Deal
EU Marine Strategy Framework Directive (MSFD)
EU Water Framework Directive (WFD)
EU Bathing water Directive
EU Maritime Spatial Planning Directive (MSP)
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Recommendation on Integrated Coastal Zone Management
EU Biodiversity Strategy



Description

The Baltic Sea is an important region for coastal and maritime tourism. The region's tourism industry employs approximately 640,000 people and registers over 227 million overnight stays annually¹. The coastal areas of the Baltic Sea provide opportunities for a wide range of tourism forms, including spas, sunbathing and beach activities, boating, fishing, ice-skating and recreational homes. The share of international tourism is substantial, especially in cruise-ship tourism². In 2019, the Port of Helsinki was the busiest international passenger port in Europe with a total of 12.2 million passengers³.



What is already happening?

Changing climate may either increase or reduce the provision of ecosystem services and resources relevant to different forms of coastal and maritime tourism⁴. On the one hand, warmer summers attract increasing numbers of coastal tourists to northern Europe. On the other hand, fewer below-freezing days shorten the winter-sports season. The growing season of cyanobacteria has significantly prolonged during the past few decades⁵, making bathing less attractive. Introduction of non-indigenous species alter opportunities for fishing and recreation⁶. Tourists are rather flexible in substituting the place, timing, and type of holiday at short notice depending on the conditions and services at the destination.



What can be expected?

The touristic importance of higher latitude destinations (such as the Baltic Sea region) is expected to grow due to climate warming, and with a higher probability of climate extremes and health risks (such as malaria resurgence) in the currently most popular destinations in southern and central Europe⁷. On the other hand, depending on the magnitude of future mitigation efforts, the coastal areas of the Baltic Sea may suffer from even more frequent and extended blooms of cyanobacteria, with related health and image risks. The future growth of coastal and maritime tourism in the Baltic Sea region has the potential to exceed the global average.



Other drivers

The tourism industry is vulnerable to external changes and pressures, including global and regional economic and political processes⁸. The changing environmental conditions and their local effects can either promote or hamper the development potential and demand for coastal and maritime tourism in the Baltic Sea region. Although coastal tourism has long been increasing, global health crises or security issues may quickly reduce the number of visits globally, regionally, and locally depending on geographical area and customer segments that suffer the consequences. For example, the COVID-19 outbreak in 2020 led to a quick collapse of international travel.



Knowledge gaps

The potential for developing coastal and maritime tourism in the Baltic Sea region depends not only on climate change, but also on associated socio-economic developments, frequency and type of yet unknown hazards, other changes in the state of marine and coastal environments and changing customer preferences. As a result, it is difficult to project the future demand for tourism services in the Baltic Sea region, or even to assign probabilities for different future outcomes. The relative importance of various qualities of coastal and marine environments for customer destination choice is poorly understood.



Policy relevance

Baltic Sea coastal and cruise tourism is important for the socioeconomy of the region. The competitiveness of this tourism depends largely on the environmental state of the Baltic Sea and the resilience of the tourism industry to natural, social, and economic changes. To improve the future prospects of blue tourism, it is important to control pollution loads, including nutrients, litter, and oil spills. Other relevant developments include multi-stakeholder governance of coastal and marine tourism and coordinated collection of economic, ecological, cultural, and social sustainability indicators⁸. Monitoring is required both for the internal development of the sector, which consists of a large number of enterprises of different sizes, and for planning public mitigation and adaptation policies.



Fisheries

Linked parameters:
Sea ice, Sea level, Wind, Waves, Pelagic and demersal fish, Coastal and migratory fish, Marine mammals, Marine protected areas, Offshore wind farms, Ecosystem function, Shipping, Aquaculture, Marine and coastal ecosystem services

Authors
Meri Kalliasvuoto, Sanna Kuningas, Anitta Lappalainen, Natural Resources Institute Finland (Luke), Finland
Örjan Östman, Jens Olsson, Rahmat Naddafi, Lena Bergström, Swedish University of Agricultural Sciences (SLU), Sweden
Oksana Glibko, St. Petersburg State Geological Unitary Enterprise "Specialized Firm Mineral", Russia
Antanas Kontautas, Marine research institute, Klaipėda University, Lithuania

Human activities

Links to main policies:
HELCOM Baltic Sea Action Plan (BSAP)
UN Sustainable Development Goals 2 and 14
UN Convention on Biological Diversity (CBD)
EU Maritime Spatial Planning Directive (MSP)
EU Common Fisheries Policy (CFP)
EU Biodiversity Strategy



Description

The commercial fishery in the Baltic Sea includes pelagic offshore and demersal fleets that contribute to 95% of total landings, and a variety of small-scale coastal fisheries. The main species targeted are Baltic herring, sprat, cod, and flatfishes. In addition, a variety of coastal freshwater and anadromous fish species are targeted. Mid-water and bottom trawls, gillnets, and trap-nets are the main gears used¹. Recreational fishing is common in coastal areas². For certain coastal species, the recreational catch is comparable or even higher than the commercial catch^{3,4}.



What is already happening?

● In the northern Baltic Sea, trawl fishing has already seen an earlier seasonal start in some years, with better operating conditions due to a shorter period of ice cover⁵. Coastal recreational ice fishing opportunities have been reduced². In much of the Baltic Sea, small-scale wintertime coastal fishing has also suffered from competition with seals that find ice-free fishing sites easier to access⁵. The species composition targeted especially by the coastal and demersal fisheries is changing due to eutrophication and climate change^{6,7}. Also, increased effort is needed for fishing-gear maintenance, due to accumulating biofilm and filamentous algae⁵.



What can be expected?

● The potential trawling season in the northern Baltic Sea will likely be extended due to a shorter ice-covered period. The main trawling areas for pelagic species are likely to shift towards more southern, shallower areas^{8,9}. The coastal and recreational fisheries will increasingly target species that prefer warmer and more nutrient-rich waters¹⁰. Some winter-time fishing will suffer from a shortage of ice and increased conflicts with seals. The recreational fisheries may become more popular with longer seasons for boat-trips and rod-fishing.



Other drivers

● Other drivers, such as changes in society, fish stocks, fishing regulations and fish markets, are likely to have as profound effects on the fisheries sector as climate change. For example, changes in consumer demand or changes of subsidies might affect the profitability of fisheries. Other environmental issues, partly interacting with climate change, such as increasing eutrophication if nutrient reductions according to the Baltic Sea Action Plan are not achieved, changes in the regulation of harmful substances, parasite infection-rates in fish, and the dispersal of non-indigenous species, will also affect the quantity and quality of fish, and the demand for the catch.



Knowledge gaps

Scientific evidence for alteration in Baltic Sea fisheries driven by climate change is still sparse. Complicated interacting and potentially additive effects in the environment, ecosystem and society make it very challenging to predict the potential consequences of climate change on different fisheries. Therefore, conclusions are confined to the currently observed trends.



Policy relevance

Fisheries have an important role in marine economy, providing work and healthy food. Fisheries activities are regulated by the EU Common Fisheries Policy and on national level. Fish stocks' monitoring and management plans should be adaptive and adjustable to mitigate climate change effects and ensure resilience¹¹. To acknowledge the potentially negative effect on fish stocks and other factors affecting the prospects of fisheries under climate change, a precautionary approach has to be applied. Climate change is only one of many challenges facing the fisheries sector: competition with apex predators and other fisheries sectors, low profitability, conflicts over shared resources, decreasing stocks of targeted species, and harmful substances are major concerns.



Aquaculture

Linked parameters:
Water temperature, Sea ice, Salinity and saltwater inflows, Waves, Oxygen, Pelagic and demersal fish, Coastal and migratory fish, Waterbirds, Marine mammals, Non-indigenous species, Marine protected areas, Nutrient concentrations and eutrophication, Fisheries, Blue carbon storage capacity, Marine and coastal ecosystem services

Authors
Anders Kiessling and Martyn Futter, Swedish University of Agricultural Sciences (SLU), Sweden
Georg Martin, UT, University of Tartu, Estonia
Lauri Niskanen, Markus Kankainen and Jouni Vilma, Natural Resources Institute Finland (Luke), Finland
Martin Karlsson and Martin Reutgård, Ecopelag
Knut-Olof Lerche, Raisio

Human activities

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goals 2, 6, 12, and 14
UN Convention on Biological Diversity
EU Green Deal
EU Water Framework Directive (WFD)
EU Maritime Spatial Planning Directive (MSP)
EU Habitats Directive
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Biodiversity Strategy



Description

Baltic aquaculture is currently dominated by open cage rainbow trout farms and contributes <0.5% of the total nutrient load to the Baltic Sea¹. Farms are located throughout the Baltic at Åland and Åbo Archipelago (Finland), the Danish straits, and a few other scattered locations. In both Finland and Sweden, farm closure and relocations have significantly reduced local-scale farm impacts on the marine environment. Finland and Estonia are evaluating offshore locations with a first pilot farm in the Bothnian and Tagalaht bays. Extractive farming where blue mussels and macro-algae are harvested as a way to recover excessive marine nutrients for terrestrial use, is also being explored.



What is already happening?

● Summer surface-water temperatures periodically exceed the optimal for rainbow trout in the whole Baltic and especially in the northern areas² reducing physical fitness, impairing growth, and increasing mortality³. Fish species presently farmed are unlikely to be affected by changing salinity, but any increase in terrestrial nutrient loading could be negative for aquaculture. Warmer water could promote farming of more temperature resilient species, such as perch and pikeperch. ● Farming of blue mussels and macro algae is negatively affected by both warmer water and lower salinity. Increased waves and more heatwaves, as well as increased predation by fish and birds would increase mussel losses^{4,5}.



What can be expected?

● Any temperature increase, especially in combination with high algae concentrations, will further stress currently farmed organisms. A possible salinity decrease will limit mussel farming and force a shift to cultivation of freshwater tolerant plants and invertebrates. Increasing policies promoting farming in more exposed locations will raise production costs. Offshore aquaculture, especially for mussels, but also for fish, could be co-located with offshore wind farms, offering moorings at locations with high water exchange, without risk of interference with shipping and recreation⁶.



Other drivers

● Policies promoting circular production and rural development will be positive drivers for aquaculture, however, industrial sized land-based systems are not likely to be implemented in remote locations within the archipelago, due to infrastructure dependencies. Marine spatial planning priorities, consumer acceptance of farmed fish, as a complement to wild fish, as well as governmental acceptance of blue catch crops, are all important for future Baltic Sea aquaculture. Synergies between renewable energy and food production based on co-location of aquaculture with offshore energy should especially promote extractive aquaculture. Demands for resilient, resistant, local food production and the possibility of local and circular-based feed sources, should further promote all types of aquaculture.



Knowledge gaps

There are multiple knowledge gaps related to climate change effects on Baltic Sea aquaculture. Regional differences are incompletely understood. Reliable, local-scale projections of future water temperatures, salinity, occurrence and toxicity of algae blooms, and ice cover are needed for siting new farms. Use of native species tolerant of possible future conditions requires knowledge of techniques and ecosystem effects, including use of sterilized fish. New farming technologies offering an economically feasible solution for particle recapture and deep-water siting must also be developed and evaluated. Credible environmental assessment of both sediment and total nutrient budgets of offshore farms using Baltic feed sources are also needed. Furthermore, alternative, and new species, especially those on lower trophic levels, and their acceptance by consumers is not well investigated.



Policy relevance

Aquaculture has the potential to provide sustainable, climate-smart local food while counteracting marine eutrophication. Political obstacles and public perceptions are probably more difficult challenges to Baltic Sea aquaculture than the changing climate. Aquaculture using sterile fish, which express neither phenotypic nor behavioural spawning characteristics, is needed to protect Baltic Sea biodiversity. Policy support for science-based solutions incorporating technological innovation and best practices is needed, as are marine spatial planning processes that avoid environmentally sensitive sites but still allow aquaculture to develop to meet European and regional policy targets, such as the EU Blue Growth Strategy.



Blue Carbon storage capacity

Linked parameters:
Water temperature, Sea ice, Salinity and saltwater inflows, Carbonate chemistry, Sea level, Waves, Benthic habitats, Marine protected areas, Nutrient concentrations and eutrophication, Offshore wind farms, Aquaculture, Marine and coastal ecosystem services

Authors
Dorte Krause-Jensen, Århus University, Denmark

Services

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goals 13 and 14
UN Convention on Biological Diversity
EU Green Deal
EU Marine Strategy Framework Directive (MSFD)
EU Water Framework Directive (WFD)
EU Maritime Spatial Planning Directive (MSP)
EU Habitats Directive
EU Strategy for the Baltic Sea Region (EUSBSR)
EU Biodiversity Strategy



Description

Blue Carbon (BC) refers to organic carbon that is captured and stored by marine and coastal ecosystems. Vegetated coastal ecosystems, which fringe global coastlines, support disproportionately large carbon sinks and are, therefore, the focus here^{1,2}. These “BC ecosystems” are under pressure and have experienced major global losses in area^{3,4} and, hence, losses of carbon sink capacity^{5,6}. Management strategies to protect and restore them therefore contribute to mitigating climate change². This is a win-win strategy as the BC ecosystems also constitute natural coastal protection and support biodiversity and other ecosystem services^{2,4,7,8}. In the Baltic Sea, vegetated coastal ecosystems encompass tidal marsh/coastal meadows, eelgrass/sea-grass meadows, and macroalgal beds.



What is already happening?

While Blue Carbon ecosystems offer mitigation and adaptation to climate change, they are also susceptible to multiple aspects of climate change such as warming, increased frequency of heatwaves, reduced sea ice cover, changing salinity and sea level rise^{7,9,10}. Relatively few such studies are yet available for BC ecosystems in the Baltic Sea but there are examples of negative effects of, e.g., warming on Baltic marine vegetation^{11,12}. Interactions between climate change and other human-induced pressures, which are prominent in the Baltic Sea¹³, tend to aggravate such negative effects of climate change on BC ecosystems in the region^{11,14}.



What can be expected?

Climate-change related effects on Blue Carbon ecosystems in the Baltic Sea and elsewhere are expected to increase in the future, with associated impacts on their mitigation and adaptation capacity. There are, e.g., projections of negative effects of climate change on Baltic eelgrass meadows and macroalgal beds^{14,15}, and flooding over tidal marshes as the sea level rises¹⁶. However, the extent of negative effects of climate change on BC habitats will depend on management of both climate change and other pressures^{11,13}. A recent review highlights the potential for substantial recovery by 2050 in the abundance, structure, and function of marine life, including coastal vegetated ecosystems, if major pressures, including climate change, are mitigated¹⁷.(*)



Other drivers

Vegetated coastal ecosystems in the Baltic Sea and elsewhere are affected by a wide range of human-induced pressures in addition to climate change, including eutrophication, e.g., reducing water clarity, land-use changes and fisheries^{13,17}. For example, warming in interaction with eutrophication and trawling, poses key threats to Baltic eelgrass meadows¹¹, and release of the local pressures can increase resilience of the meadows against realized and further warming^{11,14}. Likewise, land-use management can help relieve the risk of coastal squeezing of tidal marshes in the face of climate change, while also supporting coastal protection^{16,18,19}. The future of BC ecosystems and their climate change mitigation capacity therefore depends on sustainable, holistic management of combined pressures.



Knowledge gaps

Knowledge gaps at the Baltic Sea scale include quantification of the role of vegetated habitats in the marine carbon cycle of the region, i.e., mapping their area and related carbon fluxes (primary production, sequestration rates, export fluxes and fate) and identifying carbon sink areas beyond these habitats. Moreover, there is a need to quantify realized changes in vegetated areas and to estimate the potential to expand vegetated areas through restoration and protection as a nature-based solution (NBS) for mitigating climate change. Identification of target areas for restoration and protection of Blue Carbon ecosystems and carbon sinks, will maximize the benefit of NBS. A recent review provides further guidance on Blue Carbon science and management²⁰ and the recently concluded Nordic Blue Carbon project provides an update on eelgrass and macroalgae in the Nordic Blue Carbon context²¹.



Policy relevance

Restoration and conservation of coastal vegetated ecosystems are direct sustainable management measures to mitigate climate change, while also stimulating biodiversity and additional ecosystem functions. Blue Carbon-strategies are, therefore, important nature-based solutions to two concurring global challenges: climate change and biodiversity loss, which are increasingly addressed in international policy, for example Blue Carbon Initiative²², IUCN²³, the Ocean Panel¹¹, the Nordic Council of Ministers²⁴ and EU initiatives on nature-based solutions (NBS²⁵ also involving the Baltic Sea²⁶. There are local-scale initiatives in the Baltic Sea to protect and restore eelgrass, implement coastal realignment programs with tidal marshes and recreate reefs. However, coordinated Baltic-scale Blue Carbon-strategies represent a yet untapped potential.

*) Several studies confirm climate-related effects on marine vegetation and associated carbon sink capacity, with the extent and direction of change differing between regions e.g., depending on latitude and interaction with other pressures.



Marine and coastal ecosystem services

Linked parameters:
Sea ice, Waterbirds, Marine mammals, Marine protected areas, Nutrient concentrations and eutrophication, Ecosystem function, Tourism, Fisheries, Aquaculture, Blue carbon storage capacity

Authors
Artūras Razinkovas-Baziukas, Marine Research Institute, Klaipėda University, Lithuania

Services

Links to main policies:
HELCOM Baltic Sea Action Plan
UN Sustainable Development Goals 2, 5, 8, 12, and 14
UN Convention on Biological Diversity
EU Green Deal
EU Water Framework Directive (WFD)
EU Maritime Spatial Planning Directive (MSP)
EU Biodiversity Strategy
EU Habitats Directive
EU Strategy for the Baltic Sea Region (EUSBSR)



Description

Ecosystem services (ES) are commonly assessed as supply (mostly related to the biophysical or ecological characteristics of the environment), demand (mostly societal drivers) and flow (actual provision and use). The values of ES supply relate directly to ecosystem components¹, which in turn, are altered by climatic changes. The assessment of ES demand is based mostly on the global societal changes like alteration of lifestyles or climate change induced alterations in ES supply at global scales (e.g., further increase of summer temperatures in Southern Europe could trigger recreational demand in the Baltic). While ES flow functionally is a result of interaction between supply and demand functions, it could, in some cases, be easily assessed (even using econometric methods).



What is already happening?

The present trends in cultural ecosystem services, mainly tourism and recreation²⁻⁴, are positive but the interplay between the different drivers makes the relation to climate change uncertain. There is a negative trend in aquaculture (a provisioning ecosystem service), as water temperature is suboptimal for rainbow trout, mussel, and algae farming, while impacts on fisheries are mixed⁵⁻⁷. The trends in the supply of regulation and maintenance ecosystem services based on the key messages on eutrophication, benthic habitats, and ecosystem function varies across different parts of the Baltic⁸⁻¹².



What can be expected?

The cultural ecosystem services (mainly relevant to tourism and recreation, including recreational boating) may benefit from a longer bathing season and increased air and water temperature in the summer. However, warming may counteract these benefits due to impact on human health. Furthermore, ice fishing possibilities along with an expected shift towards smaller fish will negatively affect recreational fisheries. The supply of provisioning ecosystem services (mainly the most valuable fish stocks in the Baltic) are expected to decrease both in quantity, but especially in quality¹³, while aquaculture is not expected to counteract such a potential development by providing more high-quality fish. Moreover, the anticipated climatic changes are expected to reduce the role of semi-enclosed areas as coastal filters¹⁴.



Other drivers

Several anthropogenic pressures such as eutrophication, pollution, microplastics, fishing as well as habitat degradation, could offset positive and strengthen negative trends in ecosystem services supply, while protection and restoration efforts could lead to their improvement. Uncertainty in the future pressures increases the uncertainty of projections for ecosystem services.



Knowledge gaps


In the terrestrial domain, there are already many scientifically sound biophysical ES models capable of producing spatial coverage and tools for biodiversity and natural resources management, but the modelling of aquatic ES hitherto is largely underdeveloped^{3,15}. Moreover, so far there is little understanding of the relationships and feedbacks between aquatic ecosystems and the services they produce, resulting in the negligible impact on the policy process of economic valuation of coastal and marine ES⁴.



Policy relevance

Marine and coastal ecosystem services are critical for both strategic and territorial planning in the Baltic Sea region, because of the societal reliance on them. Mitigation measures for the ES parameters are diverse and described in the other key messages. The essence of ES is agreed in most of the major international agreements on environmental protection, and especially highlighted in the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services' 2019 Global Assessment Report. The ES concept should be guiding for Baltic Sea policies, but as there is no commonly agreed method to calculate or assess ES, such a method needs to be developed.

Glossary, policy linkages and
references





Glossary

Glossary of words related to climate change

accretion — deposition of sediment, opposite of erosion
albedo — amount of sun light reflected by a surface or a cloud
alkalinity — the capacity of water to resist acidification and maintain a stable pH level
anoxia — oxygen-free conditions in the environment or tissues of a body of an organism
anthropogenic — human derived, for example greenhouse gases from fossil fuel use
Atlantic Multidecadal Oscillation (AMO) — describes fluctuations in North Atlantic Sea surface temperature with a 50-90-year period
atmospheric blocking — occurs when persistent high-pressure systems interrupt the normal westerly flow over middle and high latitudes
atmospheric deposition — movement of matter (e.g., nutrients, pollution) from the atmosphere to the Earth's surface
bacterioplankton — single-celled prokaryotes, i.e., small organisms that lack a nucleus (Bacteria and Archaea), in the water column, mainly consuming organic carbon as energy and carbon source
benthic — related to the bottom of the sea including the top sediment layers
biota — plant and animal life in an area, habitat, or period
blue carbon (BC) — in marine sciences, organic carbon that is captured and stored by marine and coastal ecosystems (the abbreviation clashes with black carbon in atmospheric sciences)
blue carbon ecosystems — vegetated coastal ecosystems that capture organic carbon
biogeochemical cycle — a set of processes by which a chemical element is transformed to different chemical substances through the biotic and abiotic parts of an ecosystem
biodiversity — variety of all living things on Earth
biomass production — production of organic matter

brownification — darker water due to more organic substances and iron
carbon flux — amount of carbon exchanged between carbon pools on Earth
carbon sink — accumulates carbon more than releases and thus lowers the atmospheric concentration of CO₂, for example the ocean
carbon source — releases more carbon than absorbs and thus increases the atmospheric concentration of CO₂, for example burning of fossil fuels
climate refuges — areas in which temperature increase is expected to be lower than on average, important areas for conservation
climate change — climate change means a change in average or in variation of conditions in the state of the climate over a long period of time, typically decades or longer, and can be caused by natural processes or external activities, such as changes in solar cycles, volcanic eruptions and changes in atmosphere and land use caused by humans
climate change signal — observed long-term trends and projections linked to climate change
climate model — complex mathematical representation of the climate system, used to project future climate conditions and to understand past climates
climate projection — simulation of Earth's climate far into the future, derived using climate models and assumptions of future developments of climate drivers (e.g., greenhouse gases, land use)
CO₂ species — composed of chemically identical molecular entities with CO₂, i.e., CO₂, H₂CO₃, HCO₃⁻ and CO₃²⁻
cyanobacterial blooms — blooms in the water formed by microscopic single-celled cyanobacteria (aka blue green algae), common in the Baltic Sea during summer due to high level of eutrophication and warm water
DIN — dissolved inorganic nitrogen
DIP — dissolved inorganic phosphorus
demersal — area near the sea bottom, for

example demersal fish live on or near the sea bottom
deoxygenation — the removal of oxygen atoms from an environment, substance, or molecule, i.e., decline of oxygen, the primary cause of deoxygenation in the Baltic Sea is eutrophication, deoxygenation leads to hypoxia with detrimental effects on biota
ecosystem function — physical, chemical, and biological processes that transform energy, nutrients, and organic matter in an ecosystem; capacity of an ecosystem to provide goods and services that are potentially useful to humans
ecosystem functioning — interaction between an ecosystem and its environment, for example biotic activities affect the physical and chemical conditions of their environment
ecosystem services — services and benefits to humans provided by the nature and healthy ecosystems, commonly assessed as supply (mostly related to the biophysical or ecological characteristics of the environment), demand (mostly societal drivers) and flow (actual provision and use)
erosion — geological process by which surface material (e.g., soil, rock) is worn and transported, caused by natural processes such as wind, water, and ice
estuary — partially enclosed coastal area extensively influenced by river freshwater discharge causing brackish water conditions and estuarine circulation
euphotic zone, photic zone — water layer close to the surface with sufficient amount of light for photosynthesis (i.e., transfer of CO₂, water, and sun light into chemical energy mainly by plants and zooplankton)
eutrophication — excessive growth of phytoplankton, algae, and plants, caused by excessive input of nutrients to the marine environment; excessive eutrophication causes reduced light conditions, oxygen depletion, cyanobacterial blooms, and other ecosystem changes

euxinia — anoxia with raised level of free hydrogen sulfide (H₂S) in the water
evolution — the development of new species by mutation of the genome and natural selection
exoskeleton — external skeleton that protects an organism; acidification may impair marine organisms, such as mussels, that build their exoskeleton out of calcium carbonate
extractive farming — aquaculture where for example blue mussels and macroalgae are harvested as a way to recover excessive marine nutrients for terrestrial use
fetch — the distance over which wind blows
foodweb — representation of who eats what in an ecosystem, describes the movement of energy and nutrients through an ecosystem, contains different trophic levels
genetic diversity — variation in the genetic composition among individuals, species or a community, genetic diversity is important for adaption to changing circumstances and is an important part of biodiversity
greenhouse gases (GHGs) — gases that absorb heat in the atmosphere, the main greenhouse gases are water vapor, carbon dioxide (CO₂), methane (CH₄), Nitrous oxide (N₂O), fluorinated gases and ozone
haline — saline, salty
halocline — vertical zone in the water column in which salinity changes rapidly, usually causing strong stratification of the water due to resulting different densities with higher densities for higher salinities, permanently found in large parts of the Baltic Sea and located below the mixed surface water layer
haul-out — land area for resting, breeding, foraging etc. of a seal or other marine mammal
hypoxia — low oxygen level in the environment or tissues of a body of an organism
instrumental period — time period of

1886-2017, routine weather observations at fixed sites started in the beginning of the instrumental period
keystone species — organism that has a substantially large effect on the communities in which it occurs, helps to maintain local biodiversity, for example bladder wrack (*Fucus vesiculosus*) in the Baltic Sea
macrophyte — large aquatic plant
meteotsunami — sea level extreme travelling in phase with atmospheric low-pressure systems
mixed layer — the surface water layer, which is well mixed, oxygenated, and of uniform density
Major Baltic Inflows (MBIs) — large, meteorologically driven saltwater inflows to the Baltic Sea which sporadically renew the deep water with saline, oxygen rich water; this is the only process that effectively ventilates the deep water of the Baltic Sea.
nature-based solutions (NBSs) — actions to protect, sustainably manage, and restore natural or modified ecosystems, that address societal challenges effectively and adaptively, simultaneously providing human well-being and biodiversity benefits (IUCN description)
non-indigenous species (NIS) — species not native to the geographic region of interest, but transferred there by human activities
North Atlantic Oscillation (NAO) — describes the intensity of the westerly flow. A positive NAO is related to mild, wet winters and increased storminess over northern Europe
nucleus — a separate entity in a cell surrounded by a membrane where the genome is situated, is characteristic of plant and animal cells
ocean acidification — decrease of seawater pH, due mostly to the rising CO₂ concentration in the atmosphere and its exchange with the surface seawater
open-cage farm — aquaculture in open-top cages in coastal waters or lakes
organic matter — carbon-based com-

pounds found in nature, e.g., in plants, animals, their remains and dissolved organic matter in the water
pCO₂ — partial pressure of CO₂
pH — measure of acidity or basicity of a solution, acidic solutions (values <7) have lower pH values than basic or alkaline solutions
paleoclimate period — time period before 1886 from which no instrumental climate records are available
pelagic — refers to the water column above the bottom and below the sea surface in open sea regions
phenological mismatches — a mismatch in the timing of the life cycle between different organisms and environmental features, for example hatching of juveniles during sub-optimal conditions
phenology — the study of cyclic and seasonal natural phenomena, especially in relation to climate and plant and animal life
phenotypic or behavioural spawning characteristics — changes in behavior or appearance related to spawning, e.g., muscle tissue of fish becoming lighter and of inferior quality for food consumption during spawning
photosynthesis — conversion of sun light energy to chemical energy by plants, algae, and certain bacteria, binds carbon dioxide and produces oxygen, crucial for maintenance of life on Earth
phytoplankton — photosynthesizing microscopic marine organisms, inhabit the upper water layer, foundation of the aquatic food web
piscivorous fish species — fish that eat other fish
primary production — synthesis of organic compounds from carbon dioxide by plants and other organisms capable of photosynthesis. Oxygen is an important by-product of photosynthesis.
protozoa — unicellular organisms of many species eating other small organisms or bacteria
pycnocline — vertical zone in the water column in which density changes





Glossary (continued)

rapidly caused by corresponding changes of temperature and/or salinity with lower densities to the surface and higher densities to the bottom due to stratification

radiative forcing — difference between solar energy absorbed by the Earth and radiated back to space

range shifts — changes in the distribution limits of a species

refractory compounds — chemical compounds difficult to decompose and use as food

regime shifts — large, persistent change in the structure and function of an ecosystem, e.g., in Central Baltic Sea a previously cod-dominated system changed to domination by small pelagic fish, due to overfishing, eutrophication and climate change

remineralization — breakdown or transformation of organic matter to inorganic chemical compounds

Representative Concentration Pathway (RCP) — used to describe different climate futures depending on the greenhouse gas (GHG) emissions in the coming years. The RCPs indicate a possible range of radiative forcing in the year 2100 compared to 1850.; the RCPs include a “mitigation” scenario which aims to keep global warming below 2°C above pre-industrial temperatures (RCP2.6) and a high emissions “worst case” scenario (RCP8.5) that corresponds to a future without climate mitigation; One intermediate scenario is the RCP4.5 which likely results in global mean temperature rise between 2-3°C degrees by 2100.

recruitment — successful reproduction and survival of offspring

run-off — waterflow on land when the amount of water exceeds the ability of land to absorb water, increased river run-off increases input of nutrients to the Baltic Sea

climate scenario — representation of future climate

scenario simulations — simulations of future climate using numerical climate models driven by different assumptions on the future development of greenhouse gas emissions, land use, and aerosol emissions

sediment budget — balance between sediment added and removed from a coastal system

stratification — vertical ordering of inhomogeneous sea water according to its different densities due to gravitation, different densities in sea water are caused by temperature and/or salinity variation
thermocline — vertical zone in the water column in which temperature changes rapidly usually causing stratification of the water due to resulting different densities with lower densities for higher temperatures, in most parts of the Baltic Sea seasonally found with heated surface water in summer

thermodynamic equilibrium — stationary state of a system in which all macroscopic fluxes vanish, i.e., no macroscopic heat exchange equivalent to uniform temperature, no mechanical work between macroscopic parts of the system equivalent to uniform pressure, and stable phase composition (e.g., solid-liquid) equivalent to equal chemical potentials

trophic cascade — a change of one species or trophic level of the food web (such as removal or addition of top predators) that triggers substantial changes in ecosystem structure, nutrient flows, and ecosystem functions

trophic level — a group of organisms in a food web of similar size

zooplankton — microscopic marine organisms which are not photosynthesizing, but feed on other organisms



Policy linkages

Linkages between the parameters affected by climate change and various major policies




Figure 3. Linkages between the different parameters that were used in the assessment of the effects of climate change in the Baltic Sea and major policies.





References

Introduction

- 1 IPCC, 2021: Summary for Policymakers. In: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu and B. Zhou (eds.)]. Cambridge University Press. In Press
- 2 IPCC, 2013: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp.
- 3 IPCC, 2019: IPCC Special Report on the Ocean and Cryosphere in a Changing Climate [H.-O. Pörtner, D.C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, A. Alegría, M. Nicolai, A. Okem, J. Petzold, B. Rama, N.M. Weyer (eds.)]. In press.
- 4 BACC Author Team, 2008: Assessment of Climate Change for the Baltic Sea Basin. Springer Verlag, 474 p.
- 5 BACC II Author Team, 2015: Second Assessment of Climate Change for the Baltic Sea Basin. Springer Regional Climate Studies Open Access, 501 p

Air temperature

- 1 ACC Author Team. Assessment of climate change for the Baltic Sea basin. (Springer Science & Business Media, 2008).
- 2 BACC II Author Team. Second Assessment of Climate Change for the Baltic Sea Basin. (Springer International Publishing, 2015).
- 3 Rutgersson, A., Jaagus, J., Schenk, F. & Stendel, M. Observed changes and variability of atmospheric parameters in the Baltic Sea region during the last 200 years. *Clim. Res.* 61, 177-190, <https://doi.org/10.3354/cr01244> (2014).
- 4 Huth, R., Kysely, J. & Pokorná, L. A GCM Simulation of Heat Waves, Dry Spells, and Their Relationships to Circulation. *Clim. Change* 46, 29-60, <https://doi.org/10.1023/A:1005633925903> (2000).
- 5 Kjellström, E., Nikulin, G., Hansson, U. L. F., Strandberg, G. & Ullerstig, A. 21st century changes in the European climate: uncertainties derived from an ensemble of regional climate model simulations. *Tellus A: Dynamic Meteorology and Oceanography* 63, 24-40, <https://doi.org/10.1111/j.1600-0870.2010.00475.x> (2011).
- 6 Gröger, M., Dieterich, C. & Meier, H. E. M. Is interactive air sea

coupling relevant for simulating the future climate of Europe? *Clim. Dyn.* 56, 491-514, <https://doi.org/10.1007/s00382-020-05489-8> (2021).

- 7 Nikulin, G., Kjellström, E., Hansson, U. L. F., Strandberg, G. & Ullerstig, A. Evaluation and future projections of temperature, precipitation and wind extremes over Europe in an ensemble of regional climate simulations. *Tellus A: Dynamic Meteorology and Oceanography* 63, 41-55, <https://doi.org/10.1111/j.1600-0870.2010.00466.x> (2011).
- 8 Benestad, R. E. How often can we expect a record event? *Clim. Res.* 25, 3-13 (2003).

Water temperature

- 1 Prandle, D. & Lane, A. The annual temperature cycle in shelf seas. *Cont. Shelf Res.* 15, 681-704, [https://doi.org/10.1016/0278-4343\(94\)E0029-L](https://doi.org/10.1016/0278-4343(94)E0029-L) (1995).
- 2 Belkin, I. M. Rapid warming of large marine ecosystems. *Prog. Oceanogr.* 81, 207-213, <https://doi.org/10.1016/j.pocean.2009.04.011> (2009).
- 3 Siegel, H. & Gerth, M. Sea Surface Temperature in the Baltic Sea 2018. 7pp (2019).
- 4 Kniebusch, M., Meier, H. E. M., Neumann, T. & Börgel, F. Temperature Variability of the Baltic Sea Since 1850 and Attribution to Atmospheric Forcing Variables. *J. Geophys. Res-Oceans* 124, 4168-4187, <https://doi.org/10.1029/2018jc013948> (2019).
- 5 Naumann, M. et al. Hydrographic-hydrochemical assessment of the Baltic Sea 2018. *Meereswiss. Ber., Warnemünde* 110, <https://doi.io-warnemuende.de/10.12754/msr-2019-0110> (2019).
- 6 Humborg, C. et al. High Emissions of Carbon Dioxide and Methane From the Coastal Baltic Sea at the End of a Summer Heat Wave. *Front. Mar. Sci.* 6, <https://doi.org/10.3389/fmars.2019.00493> (2019).
- 7 Balmaseda, M. A., Mogensen, K. & Weaver, A. T. Evaluation of the ECMWF ocean reanalysis system ORAS4. *Q. J. Roy. Meteor. Soc.* 139, 1132-1161, <https://doi.org/10.1002/qj.2063> (2013).
- 8 Balmaseda, M. A., Trenberth, K. E. & Källén, E. Distinctive climate signals in reanalysis of global ocean heat content. *Geophys. Res. Lett.* 40, 1754-1759, <https://doi.org/10.1002/grl.50382> (2013).
- 9 Meier, H. E. M. & Saraiva, S. Projected Oceanographical Changes in the Baltic Sea until 2100. *Oxford Research Encyclopedia of Climate Science*, <https://doi.org/10.1093/acrefore/9780190228620.013.699> (2020).
- 10 Gröger, M., Arneborg, L., Dieterich, C., Höglund, A. & Meier, H. E. M. Summer hydrographic changes in the Baltic Sea, Kattegat and Skagerrak projected in an ensemble of climate scenarios downscaled with a coupled regional ocean-sea ice-atmosphere model. *Clim. Dyn.*, <https://doi.org/10.1007/s00382-019-04908-9> (2019).
- 11 Saraiva, S. et al. Uncertainties in Projections of the Baltic Sea Ecosystem Driven by an Ensemble of Global Climate Models. *Front. Earth Sci.* 6, <https://doi.org/10.3389/feart.2018.00244> (2019).
- 12 Meier, H. E. M., Dieterich, C. & Gröger, M. Natural variability is a large source of uncertainty in future projections of hypoxia in the Baltic Sea. *Commun. Earth Environ.* 2, 50, <https://doi.org/10.1038/s43247-021-00115-9> (2021).
- 13 Meier, H. E. M. et al. Future projections of record-breaking sea

surface temperature and cyanobacteria bloom events in the Baltic Sea. *Ambio* 48, 1362-1376, <https://doi.org/10.1007/s13280-019-01235-5> (2019).

- 14 Perry, D., Hammar, L., Linderholm, H. W. & Gullström, M. Spatial risk assessment of global change impacts on Swedish seagrass ecosystems. *PLoS ONE* 15, e0225318, <https://doi.org/10.1371/journal.pone.0225318> (2020).
- 15 Queirós, A. M. et al. Solutions for ecosystem-level protection of ocean systems under climate change. *Global Change Biol.* 22, 3927-3936, <https://doi.org/10.1111/gcb.13423> (2016).

Large-scale atmospheric circulation

- 1 Hurrell, J. W., Kushnir, Y., Ottersen, G. & Visbeck, M. An overview of the North Atlantic oscillation. *Geophys. Monogr. Ser.* 134, 1-36 (2003).
- 2 Andersson, H. C. Influence of long-term regional and large-scale atmospheric circulation on the Baltic sea level. *Tellus A: Dynamic Meteorology and Oceanography* 54, 76-88, <https://doi.org/10.3402/tellusa.v54i1.12125> (2002).
- 3 Lehmann, A., Krauß, W. & Hinrichsen, H.-H. Effects of remote and local atmospheric forcing on circulation and upwelling in the Baltic Sea. *Tellus A: Dynamic Meteorology and Oceanography* 54, 299-316, <https://doi.org/10.3402/tellusa.v54i3.12138> (2002).
- 4 Kauker, F. & Meier, H. E. M. Modeling decadal variability of the Baltic Sea: 1. Reconstructing atmospheric surface data for the period 1902-1998. *J. Geophys. Res-Oceans* 108, <https://doi.org/10.1029/2003JC001797> (2003).
- 5 Meier, H. E. M. & Kauker, F. Modeling decadal variability of the Baltic Sea: 2. Role of freshwater inflow and large-scale atmospheric circulation for salinity. *J. Geophys. Res-Oceans* 108, <https://doi.org/10.1029/2003JC001799> (2003).
- 6 Omstedt, A. & Chen, D. Influence of atmospheric circulation on the maximum ice extent in the Baltic Sea. *J. Geophys. Res-Oceans* 106, 4493-4500, <https://doi.org/10.1029/1999JC000173> (2001).
- 7 Tinz, B. On the relation between annual maximum extent of ice cover in the Baltic Sea and sea level pressure as well as air temperature field. *Geophysica* 32, 319-341 (1996).
- 8 Zorita, E. & Laine, A. Dependence of salinity and oxygen concentrations in the Baltic Sea on large-scale atmospheric circulation. *Clim. Res.* 14, 25-41, <https://doi.org/10.3354/cr014025> (2000).
- 9 Rex, D. F. Blocking Action in the Middle Troposphere and its Effect upon Regional Climate. Part II: The climatology of blocking action. *Tellus A: Dynamic Meteorology and Oceanography* 2, 275-301, <https://doi.org/10.3402/tellusa.v2i4.8603> (1950).
- 10 Rex, D. F. Blocking Action in the Middle Troposphere and its Effect upon Regional Climate. Part I: An aerological study of blocking action. *Tellus A: Dynamic Meteorology and Oceanography* 2, 196-211, <https://doi.org/10.3402/tellusa.v2i3.8546> (1950).
- 11 Enfield, D. B., Mestas-Núñez, A. M. & Trimble, P. J. The Atlantic Multidecadal Oscillation and its relation to rainfall and river flows in the continental U.S. *Geophys. Res. Lett.* 28, 2077-2080, <https://doi.org/10.1029/2000gl012745> (2001).
- 12 Kerr, R. A. A North Atlantic Climate Pacemaker for the Centuries. *Science* 288, 1984, <https://doi.org/10.1126/science.288.5473.1984> (2000).

- 13 Knight, J. R., Folland, C. K. & Scaife, A. A. Climate impacts of the Atlantic Multidecadal Oscillation. *Geophys. Res. Lett.* 33, <https://doi.org/10.1029/2006gl026242> (2006).
- 14 Börgel, F., Frauen, C., Neumann, T., Schimanke, S. & Meier, H. E. M. Impact of the Atlantic Multidecadal Oscillation on Baltic Sea Variability. *Geophys. Res. Lett.* 45, 9880-9888, <https://doi.org/10.1029/2018GL078943> (2018).
- 15 Börgel, F., Frauen, C., Neumann, T. & Meier, H. E. M. The Atlantic Multidecadal Oscillation controls the impact of the North Atlantic Oscillation on North European climate. *Environ. Res. Lett.*, <https://doi.org/10.1088/1748-9326/aba925> (2020).
- 16 Davini, P., Cagnazzo, C., Gualdi, S. & Navarra, A. Bidimensional Diagnostics, Variability, and Trends of Northern Hemisphere Blocking. *J. Climate* 25, 6496-6509, <https://doi.org/10.1175/JCLI-D-12-00032.1> (2012).
- 17 Croci-Maspoli, M., Schwierz, C. & Davies, H. C. A Multifaceted Climatology of Atmospheric Blocking and Its Recent Linear Trend. *J. Climate* 20, 633-649, <https://doi.org/10.1175/JCLI4029.1> (2007).
- 18 Mokhov, I. I. et al. Blockings in the Northern hemisphere and Euro-Atlantic region: Estimates of changes from reanalysis data and model simulations. *Dokl. Earth Sci.* 449, 430-433, <https://doi.org/10.1134/S1028334X13040144> (2013).
- 19 IPCC. Climate Change 2013 – The Physical Science Basis: Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. 1535pp (Cambridge University Press, 2014).
- 20 Frajka-Williams, E., Beaulieu, C. & Duchez, A. Emerging negative Atlantic Multidecadal Oscillation index in spite of warm subtropics. *Sci. Rep.-UK* 7, 11224, <https://doi.org/10.1038/s41598-017-11046-x> (2017).
- 21 Knudsen, M. F., Seidenkrantz, M.-S., Jacobsen, B. H. & Kuijpers, A. Tracking the Atlantic Multidecadal Oscillation through the last 8,000 years. *Nat. Commun.* 2, 178, <https://doi.org/10.1038/ncomms1186> (2011).
- 22 Wang, J. et al. Internal and external forcing of multidecadal Atlantic climate variability over the past 1,200 years. *Nat. Geosci.* 10, 512-517, <https://doi.org/10.1038/ngeo2962> (2017).
- 23 Stendel, M., Francis, J., White, R., Williams, P. D. & Woollings, T. In *Climate Change: Observed Impacts on Planet Earth* (ed Trevor M Letcher) Ch. 15, (Elsevier, 2020).
- 24 Wu, S., Liu, Z.-Y., Cheng, J. & Li, C. Response of North Pacific and North Atlantic decadal variability to weak global warming. *Adv. Clim. Chang. Res.* 9, 95-101, <https://doi.org/10.1016/j.accre.2018.03.001> (2018).

Sea ice

- 1 Haapala, J. J., Ronkainen, I., Schmelzer, N. & Sztobryn, M. in *Second Assessment of Climate Change for the Baltic Sea Basin* (ed BACC II Author Team) 145-153 (Springer International Publishing, 2015).
- 2 Schwegmann, S. & Holfort, J. Regional distributed trends of sea ice volume in the Baltic Sea for the 30-year period 1982 to 2019. *Meteorol. Z.* 30, 33-43, <https://doi.org/10.1127/metz/2020/0986> (2020).
- 3 Seinä, A. & Palosuo, E. The classification of the maximum annual extent of ice cover in the Baltic Sea 1720–1995. *Meri* 27, 79-91 (1996).
- 4 Luomaranta, A. et al. Multimodel estimates of the changes in





- the Baltic Sea ice cover during the present century. *Tellus A: Dynamic Meteorology and Oceanography* 66, 22617, <https://doi.org/10.3402/tellusa.v66.22617> (2014).
- 5 Höglund, A., Pemberton, P., Hordoir, R. & Schimanke, S. Ice conditions for maritime traffic in the Baltic Sea in future climate. *Boreal Environ. Res.* 22, 245-265 (2017).
- 6 Schmelzer, N. & Holfort, J. Climatological Ice Atlas for the Western and Southern Baltic Sea:(1961-2010): Digital Supplement: Comparison of Ice Conditions in the 30-year Periods 1961-1990, 1971-2000, 1981-2010. (Bundesamt für Seeschifffahrt und Hydrographie (BSH), 2012).
- 7 Dansereau, V., Weiss, J., Saramito, P. & Lattes, P. A Maxwell elasto-brittle rheology for sea ice modelling. *Cryosphere* 10, 1339-1359, <https://doi.org/10.5194/tc-10-1339-2016> (2016).

Solar radiation

- 1 Wild, M. et al. From Dimming to Brightening: Decadal Changes in Solar Radiation at Earth's Surface. *Science* 308, 847, <https://doi.org/10.1126/science.1103215> (2005).
- 2 Wild, M. Enlightening Global Dimming and Brightening. *Bull. Amer. Meteor. Soc.* 93, 27-37, <https://doi.org/10.1175/bams-d-11-00074.1> (2012).
- 3 Wild, M. et al. The Global Energy Balance Archive (GEBA) version 2017: a database for worldwide measured surface energy fluxes. *Earth Syst. Sci. Data* 9, 601-613, <https://doi.org/10.5194/essd-9-601-2017> (2017).
- 4 Wild, M. Decadal changes in radiative fluxes at land and ocean surfaces and their relevance for global warming. *Wires. Clim. Change* 7, 91-107, <https://doi.org/10.1002/wcc.372> (2016).
- 5 Karlsson, K.-G. & Devasthale, A. Inter-comparison and evaluation of the four longest satellite-derived cloud climate data records: CLARA-A2, ESA Cloud CCI V3, ISCCP-HGM, and PAT-MOS-x. *Remote Sens.-Basel* 10, 1567, <https://doi.org/10.3390/rs10101567> (2018).
- 6 Bartók, B. et al. Projected changes in surface solar radiation in CMIP5 global climate models and in EURO-CORDEX regional climate models for Europe. *Clim. Dyn.* 49, 2665-2683, <https://doi.org/10.1007/s00382-016-3471-2> (2017).
- 7 Allen, R. J., Norris, J. R. & Wild, M. Evaluation of multidecadal variability in CMIP5 surface solar radiation and inferred underestimation of aerosol direct effects over Europe, China, Japan, and India. *J. Geophys. Res.-Atmos.* 118, 6311-6336, <https://doi.org/10.1002/jgrd.50426> (2013).
- 8 Storelvmo, T. et al. Lethargic Response to Aerosol Emissions in Current Climate Models. *Geophys. Res. Lett.* 45, 9814-9823, <https://doi.org/10.1029/2018GL078298> (2018).
- 9 Stanhill, G., Achiman, O., Rosa, R. & Cohen, S. The cause of solar dimming and brightening at the Earth's surface during the last half century: Evidence from measurements of sunshine duration. *J. Geophys. Res.-Atmos.* 119, 9102-9110, <https://doi.org/10.1002/2013JD021308> (2014).
- 10 Parding, K., Olseth, J. A., Dagestad, K. F. & Liepert, B. G. Decadal variability of clouds, solar radiation and temperature at a high-latitude coastal site in Norway. *Tellus B: Chemical and Physical Meteorology* 66, 25897, <https://doi.org/10.3402/tellusb.v66.25897> (2014).
- 11 Ruckstuhl, C. et al. Aerosol and cloud effects on solar brightening and the recent rapid warming. *Geophys. Res. Lett.* 35, L12708, <https://doi.org/10.1029/2008GL034228> (2008).

- 12 Philipona, R., Behrens, K. & Ruckstuhl, C. How declining aerosols and rising greenhouse gases forced rapid warming in Europe since the 1980s. *Geophys. Res. Lett.* 36, <https://doi.org/10.1029/2008GL036350> (2009).

Salinity and saltwater inflows

1. Schinke, H., & Matthäus, W. (1998). On the causes of major Baltic inflows—an analysis of long time series. *Continental Shelf Research*, 18(1), 67-97.
2. Mohrholz, V. (2018). Major Baltic Inflow Statistics – Revised. *Frontiers in Marine Science* 5, 384. <https://doi.org/10.3389/fmars.2018.00384>.
3. Fonselius, S., & Valderrama, J. (2003). One hundred years of hydrographic measurements in the Baltic Sea. *Journal of Sea Research*, 49(4), 229-241.
4. BACC II Author Team (2015). Second assessment of climate change for the Baltic Sea Basin. Regional Climate Studies. Cham: Springer. <https://doi.org/10.1007/978-3-319-16006-1>.
5. Kniebusch, M., H. E. M. Meier, and H. Radtke, 2019: Changing salinity gradients in the Baltic Sea as a consequence of altered freshwater budgets. *Geophysical Research Letters*, 46, 9739–9747. <https://doi.org/10.1029/2019GL083902>
6. Meier, H. E. M., and Kauker, F. (2003). Modeling decadal variability of the Baltic Sea: 2. Role of freshwater inflow and large-scale atmospheric circulation for salinity. *Journal of Geophysical Research - Oceans* 108(C11), <https://doi.org/10.1029/2003JC001799>
7. Meier, H. E. M., Eilola, K., Almroth-Rosell, E., Schimanke, S., Kniebusch, M., Höglund, A., ... & Saraiva, S. (2019). Disentangling the impact of nutrient load and climate changes on Baltic Sea hypoxia and eutrophication since 1850. *Climate Dynamics*, 53(1-2), 1145-1166.
8. Radtke, H., Brunnabend, S. E., Gräwe, U., & Meier, H. E. (2020). Investigating interdecadal salinity changes in the Baltic Sea in a 1850–2008 hindcast simulation. *Climate of the Past*, 16(4), 1617-1642.
9. Schimanke, S. and H. E. M. Meier, 2016: Decadal to centennial variability of salinity in the Baltic Sea. *Journal of Climate*, 29(20), 7173-7188. <http://dx.doi.org/10.1175/JCLI-D-15-0443.1>
10. Börgel, F., C. Frauen, T. Neumann, S. Schimanke, and H. E. M. Meier, 2018: Impact of the Atlantic Multidecadal Oscillation on Baltic Sea variability. *Geophysical Research Letter*, 45(18), 9880-9888, <https://doi.org/10.1029/2018GL078943>.
11. Liblik, T. and Lips, U. 2019. Stratification Has Strengthened in the Baltic Sea – An Analysis of 35 Years of Observational Data. *Frontiers in Earth Science*, 7. doi:10.3389/feart.2019.00174
12. Meier, H. E. M., A. Höglund, E. Almroth-Rosell, and K. Eilola, 2017: Impact of accelerated future global mean sea level rise on hypoxia in the Baltic Sea. *Climate Dynamics*, 49, 163-172, <https://doi.org/10.1007/s00382-016-3333-y>
13. Saraiva, S., Meier, H. E. M., Andersson, H. C., Höglund, A., Dieterich, C., Gröger, M., Hordoir, R., and Eilola, K. (2019). Uncertainties in projections of the Baltic Sea ecosystem driven by an ensemble of global climate models. *Frontiers in Earth Science*, 6:244, <https://doi.org/10.3389/feart.2018.00244>.
14. Meier, H. E. M., M. Edman, K. Eilola, M. Placke, T. Neumann, H. Andersson, S.-E. Brunnabend, C. Dieterich, C. Frauen, R. Friedland, M. Gröger, B. G. Gustafsson, E. Gustafsson, A. Isaev, M. Kniebusch, I. Kuznetsov, B. Müller-Karulis, A. Omstedt, V.

- Ryabchenko, S. Saraiva, and O. P. Savchuk, 2018: Assessment of eutrophication abatement scenarios for the Baltic Sea by multi-model ensemble simulations. *Frontiers in Marine Science*, 5:440, <https://doi.org/10.3389/fmars.2018.00440>
15. Meier, H. E. M., M. Edman, K. Eilola, M. Placke, T. Neumann, H. Andersson, S.-E. Brunnabend, C. Dieterich, C. Frauen, R. Friedland, M. Gröger, B. G. Gustafsson, E. Gustafsson, A. Isaev, M. Kniebusch, I. Kuznetsov, B. Müller-Karulis, M. Naumann, A. Omstedt, V. Ryabchenko, S. Saraiva, and O. P. Savchuk, 2019: Assessment of uncertainties in scenario simulations of biogeochemical cycles in the Baltic Sea. *Frontiers in Marine Science*, 6:46, <https://doi.org/10.3389/fmars.2019.00046>
16. Schimanke, S., C. Dieterich, and H. E. M. Meier, 2014: An algorithm based on SLP- fluctuations to identify major Baltic inflow events. *Tellus A*, 66, 23452, <http://dx.doi.org/10.3402/tellusa.v66>
17. Gröger, M., L. Arneborg, C. Dieterich, A. Höglund, and H. E. M. Meier, 2019: Hydrographic changes in the North Sea and Baltic Sea projected in an ensemble of climate scenarios downscaled with a coupled regional ocean-sea ice-atmosphere model. *Climate Dynamics*, <https://doi.org/10.1007/s00382-019-04908-9>
18. Placke, M., Meier, H. E. M., and T. Neumann, 2021: Sensitivity of the Baltic Sea overturning circulation to long-term atmospheric and hydrological changes. *Journal of Geophysical Research – Oceans*, provisionally accepted.
19. Vuorinen, I., J. Hänninen, M. Rajasilta, P. Laine, J. Eklund, F. Montesino-Pouzols, F. Corona, K. Junker, H. E. M. Meier, and J. W. Dippner, 2015: Scenario simulations of future salinity and Ecological Consequences in the Baltic Sea and adjacent North Sea areas - implications for environmental monitoring. *Ecological Indicators*, 50, 196 - 205.
20. HELCOM (2018): State of the Baltic Sea – Second HELCOM holistic assessment 2011-2016. Baltic Sea Environment Proceedings 155, ISSN 0357-2994. Available at: www.helcom.fi/baltic-sea-trends/holistic-assessments/state-of-the-baltic-sea-2018/reports-and-materials/.

Stratification

- 1 Kniebusch, M., Meier, H. E. M. & Radtke, H. Changing Salinity Gradients in the Baltic Sea As a Consequence of Altered Freshwater Budgets. *Geophys. Res. Lett.* 46, 9739-9747, <https://doi.org/10.1029/2019GL083902> (2019).
- 2 Väli, G., Meier, H. E. M. & Elken, J. Simulated halocline variability in the Baltic Sea and its impact on hypoxia during 1961-2007. *J. Geophys. Res.-Oceans* 118, 6982-7000, <https://doi.org/10.1002/2013JC009192> (2013).
- 3 Gröger, M., Arneborg, L., Dieterich, C., Höglund, A. & Meier, H. E. M. Summer hydrographic changes in the Baltic Sea, Kattegat and Skagerrak projected in an ensemble of climate scenarios downscaled with a coupled regional ocean–sea ice–atmosphere model. *Clim. Dyn.*, <https://doi.org/10.1007/s00382-019-04908-9> (2019).
- 4 Kniebusch, M., Meier, H. E. M., Neumann, T. & Börgel, F. Temperature Variability of the Baltic Sea Since 1850 and Attribution to Atmospheric Forcing Variables. *J. Geophys. Res.-Oceans* 124, 4168-4187, <https://doi.org/10.1029/2018jc013948> (2019).
- 5 Liblik, T. & Lips, U. Stratification Has Strengthened in the Baltic Sea – An Analysis of 35 Years of Observational Data. *Front.*

- Earth Sci.* 7, <https://doi.org/10.3389/feart.2019.00174> (2019).
- 6 Meier, H. E. M. in Second Assessment of Climate Change for the Baltic Sea Basin (ed BACC II Author Team) 243-252 (Springer International Publishing, 2015).
- 7 Saraiva, S. et al. Uncertainties in Projections of the Baltic Sea Ecosystem Driven by an Ensemble of Global Climate Models. *Front. Earth Sci.* 6, <https://doi.org/10.3389/feart.2018.00244> (2019).
- 8 Meier, H. E. M., Dieterich, C. & Gröger, M. Natural variability is a large source of uncertainty in future projections of hypoxia in the Baltic Sea. *Commun. Earth Environ.* 2, 50, <https://doi.org/10.1038/s43247-021-00115-9> (2021).
- 9 Lips, I. & Lips, U. Abiotic factors influencing cyanobacterial bloom development in the Gulf of Finland (Baltic Sea). *Hydrobiologia* 614, 133-140, <https://doi.org/10.1007/s10750-008-9449-2> (2008).
- 10 Conley, D. J. et al. Tackling Hypoxia in the Baltic Sea: Is Engineering a Solution? *Environ. Sci. Technol.* 43, 3407-3411, <https://doi.org/10.1021/es8027633> (2009).

Precipitation

- 1 HELCOM. Climate Change in the Baltic Sea Area - HELCOM Thematic Assessment in 2007. 54pp (Helsinki, Finland, 2007).
- 2 Kriaučiūnienė, J. et al. Variability in temperature, precipitation and river discharge in the Baltic States. *Boreal Environ. Res.* 17, 150-162 (2012).
- 3 BACC II Author Team. Second Assessment of Climate Change for the Baltic Sea Basin. (Springer International Publishing, 2015).
- 4 Chen, D. et al. Hydroclimate changes over Sweden in the twentieth and twenty-first centuries: a millennium perspective. *Geogr. Ann. A*, 1-29, <https://doi.org/10.1080/04353676.2020.1841410> (2020).
- 5 Aalto, J., Pirinen, P. & Jylhä, K. New gridded daily climatology of Finland: Permutation-based uncertainty estimates and temporal trends in climate. *J. Geophys. Res.-Atmos.* 121, 3807-3823, <https://doi.org/10.1002/2015JD024651> (2016).
- 6 Jaagus, J., Briede, A., Rimkus, E. & Sepp, M. Changes in precipitation regime in the Baltic countries in 1966–2015. *Theor. Appl. Climatol.* 131, 433-443, <https://doi.org/10.1007/s00704-016-1990-8> (2018).
- 7 Cardell, M. F., Amengual, A., Romero, R. & Ramis, C. Future extremes of temperature and precipitation in Europe derived from a combination of dynamical and statistical approaches. *Int. J. Climatol.* 40, 4800-4827, <https://doi.org/10.1002/joc.6490> (2020).
- 8 European Environment Agency. Heavy precipitation in Europe. Indicator Assessment CLIM 004. IND-92-en, https://www.eea.europa.eu/ds_resolveuid/998fbe113cc84e9a978b-ecd87079f874, (last access: 24-03-2021).
- 9 Christensen, O. B. & Kjellström, E. Projections for Temperature, Precipitation, Wind, and Snow in the Baltic Sea Region until 2100. *Oxford Research Encyclopedia of Climate Science* Oxford University Press, <https://doi.org/10.1093/acrefore/9780190228620.013.695> (2018).
- 10 Rajczak, J. & Schär, C. Projections of Future Precipitation Extremes Over Europe: A Multimodel Assessment of Climate Simulations. *J. Geophys. Res.-Atmos.* 122, 10,773-710,800, <https://doi.org/10.1002/2017JD027176> (2017).



River run-off

- 1 Hansson, D., Eriksson, C., Omstedt, A. & Chen, D. Reconstruction of river runoff to the Baltic Sea, AD 1500–1995. *Int. J. Climatol.* 31, 696–703, <https://doi.org/10.1002/joc.2097> (2011).
- 2 Meier, H. E. M. et al. Disentangling the impact of nutrient load and climate changes on Baltic Sea hypoxia and eutrophication since 1850. *Clim. Dyn.* 53, 1145–1166, <https://doi.org/10.1007/s00382-018-4296-y> (2019).
- 3 Jaagus, J., Sepp, M., Tamm, T., Järvet, A. & Möisja, K. Trends and regime shifts in climatic conditions and river runoff in Estonia during 1951–2015. *Earth Syst. Dynam.* 8, 963–976, <https://doi.org/10.5194/esd-8-963-2017> (2017).
- 4 Sarauskienė, D., Kriauciuniene, J., Reiha, A. & Klavins, M. Flood pattern changes in the rivers of the Baltic countries. *J. Environ. Eng. Landsc. Manag.* 23, 28–38, <https://doi.org/10.3846/16486897.2014.937438> (2015).
- 5 Arheimer, B. & Lindström, G. Climate impact on floods: changes in high flows in Sweden in the past and the future (1911–2100). *Hydrol. Earth Syst. Sci.* 19, 771–784, <https://doi.org/10.5194/hess-19-771-2015> (2015).
- 6 Saraiva, S. et al. Baltic Sea ecosystem response to various nutrient load scenarios in present and future climates. *Clim. Dyn.* 52, 3369–3387, <https://doi.org/10.1007/s00382-018-4330-0> (2019).
- 7 Saraiva, S. et al. Uncertainties in Projections of the Baltic Sea Ecosystem Driven by an Ensemble of Global Climate Models. *Front. Earth Sci.* 6, <https://doi.org/10.3389/feart.2018.00244> (2019).
- 8 Stonevičius, E., Rimkus, E., Štaras, A., Kažys, J. & Valiuškevičius, G. Climate change impact on the Nemunas River basin hydrology in the 21st century. *Boreal Environ. Res.* 22, 49–65 (2017).
- 9 Šarauskienė, D. et al. Projection of Lithuanian river runoff, temperature and their extremes under climate change. *Hydrol. Res.* 49, 344–362, <https://doi.org/10.2166/nh.2017.007> (2017).
- 10 Roudier, P. et al. Projections of future floods and hydrological droughts in Europe under a +2°C global warming. *Clim. Change* 135, 341–355, <https://doi.org/10.1007/s10584-015-1570-4> (2016).
- 11 Veijalainen, N., Lotsari, E., Vehviläinen, B., Alho, P. & Käyhkö, J. in 6th Alexander von Humboldt International Conference on Climate Change (2010).
- 12 Hisdal, H. et al. Has streamflow changed in the Nordic countries? Report No. NVE-1/2010, 28pp (Norwegian Water Resources and Energy Directorate (NVE), Norway, 2010).
- 13 European Court of Auditors. Combating eutrophication in the Baltic Sea: further and more effective action needed. 71pp (Publications Office of the European Union, Luxembourg, 2016).

Carbonate chemistry

- 1 BACC II Author Team. Second Assessment of Climate Change for the Baltic Sea Basin. (Springer International Publishing, 2015).
- 2 Kuliński, K., Schneider, B., Szymczycha, B. & Stokowski, M. Structure and functioning of the acid–base system in the Baltic Sea. *Earth Syst. Dynam.* 8, 1107–1120, <https://doi.org/10.5194/esd-8-1107-2017> (2017).
- 3 Norman, M., Rutgersson, A. & Sahlée, E. Impact of improved air–sea gas transfer velocity on fluxes and water chemistry in a

- Baltic Sea model. *Journal of Marine Systems* 111–112, 175–188, <https://doi.org/10.1016/j.jmarsys.2012.10.013> (2013).
- 4 Omstedt, A. et al. Biogeochemical control of the coupled CO₂–O₂ system of the Baltic Sea: a review of the results of Baltic-C. *Ambio* 43, 49–59, <https://doi.org/10.1007/s13280-013-0485-4> (2014).
 - 5 Parard, G., Rutgersson, A., Raj Parampil, S. & Charantonis, A. A. The potential of using remote sensing data to estimate air–sea CO₂ exchange in the Baltic Sea. *Earth Syst. Dynam.* 8, 1093–1106, <https://doi.org/10.5194/esd-8-1093-2017> (2017).
 - 6 Kuliński, K. & Pempkowiak, J. The carbon budget of the Baltic Sea. *Biogeosciences* 8, 3219–3230, <https://doi.org/10.5194/bg-8-3219-2011> (2011).
 - 7 Gustafsson, E. & Gustafsson, B. G. Future acidification of the Baltic Sea – A sensitivity study. *Journal of Marine Systems* 211, 103397, <https://doi.org/10.1016/j.jmarsys.2020.103397> (2020).
 - 8 Omstedt, A. et al. Future changes in the Baltic Sea acid–base (pH) and oxygen balances. *Tellus B: Chemical and Physical Meteorology* 64, 19586, <https://doi.org/10.3402/tellusb.v64i0.19586> (2012).
 - 9 Müller, J. D., Schneider, B. & Rehder, G. Long-term alkalinity trends in the Baltic Sea and their implications for CO₂-induced acidification. *Limnol. Oceanogr.* 61, 1984–2002, <https://doi.org/10.1002/lno.10349> (2016).
 - 10 Kuznetsov, I. & Neumann, T. Simulation of carbon dynamics in the Baltic Sea with a 3D model. *Journal of Marine Systems* 111–112, 167–174, <https://doi.org/10.1016/j.jmarsys.2012.10.011> (2013).

Riverine nutrient loads and atmospheric deposition

- 1 Zillén, L., Conley, D. J., Andrén, T., Andrén, E. & Björck, S. Past occurrences of hypoxia in the Baltic Sea and the role of climate variability, environmental change and human impact. *Earth Sci. Rev.* 91, 77–92, <https://doi.org/10.1016/j.earscirev.2008.10.001> (2008).
- 2 Gustafsson, B. G. et al. Reconstructing the development of Baltic Sea eutrophication 1850–2006. *Ambio* 41, 534–548, <https://doi.org/10.1007/s13280-012-0318-x> (2012).
- 3 HELCOM. Sources and pathways of nutrients to the Baltic Sea. 48pp (Helsinki, Finland, 2018).
- 4 Ruoho-Airola, T., Eilola, K., Savchuk, O. P., Parviainen, M. & Tarvainen, V. Atmospheric Nutrient Input to the Baltic Sea from 1850 to 2006: A Reconstruction from Modeling Results and Historical Data. *Ambio* 41, 549–557, <https://doi.org/10.1007/s13280-012-0319-9> (2012).
- 5 Engardt, M., Simpson, D., Schwikowski, M. & Granat, L. Deposition of sulphur and nitrogen in Europe 1900–2050. Model calculations and comparison to historical observations. *Tellus B: Chemical and Physical Meteorology* 69, 1328945, <https://doi.org/10.1080/16000889.2017.1328945> (2017).
- 6 Savchuk, O. P., Eilola, K., Gustafsson, B. G., Medina, M. R. & Ruoho-Airola, T. Long term reconstruction of nutrient loads to the Baltic Sea, 1850–2006. *Baltic Nest Institute Techn. Rep.* 6 (2012).
- 7 Savchuk, O. P. Large-Scale Nutrient Dynamics in the Baltic Sea, 1970–2016. *Front. Mar. Sci.* 5, 95, <https://doi.org/10.3389/fmars.2018.00095> (2018).
- 8 HELCOM. Input of nutrients by the seven biggest rivers in the Baltic Sea region. 31pp (Helsinki, Finland, 2018).

- 9 HELCOM PLC waterborne database: http://nest.su.se/HELCOM_plc
- 10 BACC II Author Team. Second Assessment of Climate Change for the Baltic Sea Basin. (Springer International Publishing, 2015).
- 11 HELCOM. HELCOM thematic assessment of eutrophication 2011–2016., 83pp (Helsinki, Finland, 2018).
- 12 Bartnicki, J. 53–57 (Springer International Publishing).
- 13 Colette, A. et al. Air pollution trends in the EMEP region between 1990 and 2012. 105pp (Norwegian Institute for Air Research, Kjeller, Norway, 2016).
- 14 Gauss, M., Bartnicki, J. & Klein, H. Atmospheric nitrogen deposition to the Baltic Sea. 9pp (https://emep.int/publ/HELCOM/2018/B_BSEFS_N_dep_v2.pdf EMEP MSC-W, 2018).
- 15 HELCOM. Updated Fifth Baltic Sea Pollution Load Compilation (PLC-5.5). 143pp (Helsinki, Finland, 2015).
- 16 Kanakidou, M., Myriokefalitakis, S. & Tsigaridis, K. Aerosols in atmospheric chemistry and biogeochemical cycles of nutrients. *Environ. Res. Lett.* 13, 063004, <https://doi.org/10.1088/1748-9326/aabcbd> (2018).
- 17 McCrackin, M. L. et al. A Century of Legacy Phosphorus Dynamics in a Large Drainage Basin. *Global Biogeochem. Cy.* 32, 1107–1122, <https://doi.org/10.1029/2018GB005914> (2018).
- 18 Meier, H. E. M. et al. Disentangling the impact of nutrient load and climate changes on Baltic Sea hypoxia and eutrophication since 1850. *Clim. Dyn.* 53, 1145–1166, <https://doi.org/10.1007/s00382-018-4296-y> (2019).
- 19 Saraiva, S. et al. Baltic Sea ecosystem response to various nutrient load scenarios in present and future climates. *Clim. Dyn.* 52, 3369–3387, <https://doi.org/10.1007/s00382-018-4330-0> (2019).
- 20 Saraiva, S. et al. Uncertainties in Projections of the Baltic Sea Ecosystem Driven by an Ensemble of Global Climate Models. *Front. Earth Sci.* 6, <https://doi.org/10.3389/feart.2018.00244> (2019).
- 21 Meier, H. E. M. et al. Correction to: Disentangling the impact of nutrient load and climate changes on Baltic Sea hypoxia and eutrophication since 1850. *Clim. Dyn.* 53, 1167–1169, <https://doi.org/10.1007/s00382-018-4483-x> (2019).
- 22 Bartosova, A. et al. Future socioeconomic conditions may have a larger impact than climate change on nutrient loads to the Baltic Sea. *Ambio* 48, 1325–1336, <https://doi.org/10.1007/s13280-019-01243-5> (2019).
- 23 Simpson, D. et al. Impacts of climate and emission changes on nitrogen deposition in Europe: a multi-model study. *Atmos. Chem. Phys.* 14, 6995–7017, <https://doi.org/10.5194/acp-14-6995-2014> (2014).
- 24 Skjøth, C. A. & Geels, C. The effect of climate and climate change on ammonia emissions in Europe. *Atmos. Chem. Phys.* 13, 117–128, <https://doi.org/10.5194/acp-13-117-2013> (2013).
- 25 Sutton, M. A. et al. Towards a climate-dependent paradigm of ammonia emission and deposition. *Philos. T. Roy. Soc. B.* 368, 20130166, <https://doi.org/10.1098/rstb.2013.0166> (2013).
- 26 Meier, H. E. M. et al. Assessment of Uncertainties in Scenario Simulations of Biogeochemical Cycles in the Baltic Sea. *Front. Mar. Sci.* 6, <https://doi.org/10.3389/fmars.2019.00046> (2019).
- 27 Zandersen, M. et al. Shared socio-economic pathways extended for the Baltic Sea: exploring long-term environmental problems. *Reg. Environ. Change* 19, 1073–1086, <https://doi.org/10.1007/s10113-018-1453-0> (2019).

Sea level

- 1 Hünicke, B. et al. in Second Assessment of Climate Change for the Baltic Sea Basin (ed BACC II Author Team) 155–185 (Springer International Publishing, 2015).
- 2 Chen, D. & Omstedt, A. Climate-induced variability of sea level in Stockholm: influence of air temperature and atmospheric circulation. *Adv. Atmos. Sci.* 22, 655–664, <https://doi.org/10.1007/BF02918709> (2005).
- 3 Samuelsson, M. & Stigebrandt, A. Main characteristics of the long-term sea level variability in the Baltic sea. *Tellus A* 48, 672–683, <https://doi.org/10.1034/j.1600-0870.1996.t01-4-00006.x> (1996).
- 4 Andersson, H. C. Influence of long-term regional and large-scale atmospheric circulation on the Baltic sea level. *Tellus A* 54, 76–88, <https://doi.org/10.3402/tellusa.v54i1.12125> (2002).
- 5 Karabil, S., Zorita, E. & Hünicke, B. Contribution of atmospheric circulation to recent off-shore sea-level variations in the Baltic Sea and the North Sea. *Earth Syst. Dynam.* 9, 69–90, <https://doi.org/10.5194/esd-9-69-2018> (2018).
- 6 Lehmann, A., Getzlaff, K. & Harlaß, J. Detailed assessment of climate variability in the Baltic Sea area for the period 1958 to 2009. *Clim. Res.* 46, 185–196, <https://doi.org/10.3354/cr00876> (2011).
- 7 Weisse, R. & Weidemann, H. Baltic Sea extreme sea levels 1948–2011: Contributions from atmospheric forcing. *Proc. IUTAM* 25, 65–69, <https://doi.org/10.1016/j.piutam.2017.09.010> (2017).
- 8 Wübbler, C. & Krauss, W. The two dimensional seiches of the Baltic Sea. *Oceanol. Acta* 2, 435–446 (1979).
- 9 Wolski, T. et al. Extreme sea levels at selected stations on the Baltic Sea coast**This work was financed by the Polish National Centre for Science research project No. 2011/01/B/ST10/06470. *Oceanologia* 56, 259–290, <https://doi.org/10.5697/oc.56-2.259> (2014).
- 10 Wiśniewski, B. & Wolski, T. Physical aspects of extreme storm surges and falls on the Polish coast. *Oceanologia* 53, 373–390, <https://doi.org/10.5697/oc.53-1-TI.373> (2011).
- 11 Eelsalu, M., Soomere, T., Pindsoo, K. & Lagemaa, P. Ensemble approach for projections of return periods of extreme water levels in Estonian waters. *Cont. Shelf Res.* 91, 201–210, <https://doi.org/10.1016/j.csr.2014.09.012> (2014).
- 12 Church, J. A. et al. in Climate Change 2013 – The Physical Science Basis: Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change (eds T.F. Stocker et al.) 1137–1216 (Cambridge University Press, 2013).
- 13 Oppenheimer, M. et al. in IPCC Special Report on the Ocean and Cryosphere in a Changing Climate (eds H.-O. Pörtner et al.) (2019).
- 14 Weisse, R. et al. Sea Level Dynamics and Coastal Erosion in the Baltic Sea Region. *Earth Syst. Dynam. Discuss.*, 1–40, <https://doi.org/10.5194/esd-2021-6> (2021).
- 15 Balmaseda, M. A., Trenberth, K. E. & Köllén, E. Distinctive climate signals in reanalysis of global ocean heat content. *Geophys. Res. Lett.* 40, 1754–1759, <https://doi.org/10.1002/grl.50382> (2013).
- 16 Hammarklint, T. Swedish Sea Level Series - A Climate Indicator. 5pp (Swedish Meteorological and Hydrological Institute (SMHI), 2009).
- 17 Hill, E. M., Davis, J. L., Tamisiea, M. E. & Lidberg, M. Combination of geodetic observations and models for glacial isostatic adjustment fields in Fennoscandia. *J. Geophys. Res.-Sol. Ea.*



- 115, <https://doi.org/10.1029/2009jb006967> (2010).
- 18 Richter, A., Groh, A. & Dietrich, R. Geodetic observation of sea-level change and crustal deformation in the Baltic Sea region. *Physics and Chemistry of the Earth, Parts A/B/C* 53-54, 43-53, <https://doi.org/10.1016/j.pce.2011.04.011> (2012).
- 19 Vestøl, O., Ågren, J., Steffen, H., Kierulf, H. & Tarasov, L. NKG-2016LU: a new land uplift model for Fennoscandia and the Baltic Region. *J. Geod.* 93, 1759-1779, <https://doi.org/10.1007/s00190-019-01280-8> (2019).
- 20 Meier, H. E. M., Broman, B. & Kjellström, E. Simulated sea level in past and future climates of the Baltic Sea. *Clim. Res.* 27, 59-75, <https://doi.org/10.3354/cr027059> (2004).
- 21 Dieterich, C., Gröger, M., Arneborg, L. & Andersson, H. C. Extreme sea levels in the Baltic Sea under climate change scenarios – Part 1: Model validation and sensitivity. *Ocean Sci.* 15, 1399-1418, <https://doi.org/10.5194/os-15-1399-2019> (2019).
- 22 Madsen, K. S., Høyer, J. L., Suursaar, Ü., She, J. & Knudsen, P. Sea Level Trends and Variability of the Baltic Sea From 2D Statistical Reconstruction and Altimetry. *Front. Earth Sci.* 7, <https://doi.org/10.3389/feart.2019.00243> (2019).
- 23 Ribeiro, A., Barbosa, S. M., Scotto, M. G. & Donner, R. V. Changes in extreme sea-levels in the Baltic Sea. *Tellus A* 66, 20921, <https://doi.org/10.3402/tellusa.v66.20921> (2014).
- 24 Bamber, J. L., Oppenheimer, M., Kopp, R. E., Aspinall, W. P. & Cooke, R. M. Ice sheet contributions to future sea-level rise from structured expert judgment. *P. Natl. Acad. Sci. USA* 116, 11195, <https://doi.org/10.1073/pnas.1817205116> (2019).
- 25 Grinsted, A. in *Second Assessment of Climate Change for the Baltic Sea Basin Regional Climate Studies* (ed BACC II Author Team) 253-263 (Springer International Publishing, 2015).
- 26 Pellikka, H., Särkkä, J., Johansson, M. & Pettersson, H. Probability distributions for mean sea level and storm contribution up to 2100 AD at Forsmark. 49pp (Swedish Nuclear Fuel and Waste Management Company (SKB), 2020).
- 27 Suursaar, Ü. & Sooäär, J. Decadal variations in mean and extreme sea level values along the Estonian coast of the Baltic Sea. *Tellus A* 59, 249-260, <https://doi.org/10.1111/j.1600-0870.2006.00220.x> (2007).
- 28 Andersson, H. C. Stigande havsnivåer och ökad översvämningsrisk Report No. MSB 1243, 31pp (Swedish Meteorological and Hydrological Institute, 2018).
- 29 Hieronymus, M. & Kalén, O. Sea-level rise projections for Sweden based on the new IPCC special report: The ocean and cryosphere in a changing climate. *Ambio*, <https://doi.org/10.1007/s13280-019-01313-8> (2020).
- 30 Södling, J. & Nerheim, S. Statistisk metodik för beräkning av extrema havsvattenstånd. 44pp (SMHI, Norrköping, Sweden, 2017).
- 31 Fredriksson, C., Tajvidi, N., Hanson, H. & Larson, M. Statistical analysis of extreme sea water levels at the falsterbo peninsula, south Sweden. *Vatten, J. Water Manage. Res.* 72, 129-142 (2016).
- 32 Lang, A. & Mikolajewicz, U. The long-term variability of extreme sea levels in the German Bight. *Ocean Sci.* 15, 651-668 (2019).
- 33 Bingham, R. J. & Hughes, C. W. Local diagnostics to estimate density-induced sea level variations over topography and along coastlines. *J. Geophys. Res-Oceans* 117, <https://doi.org/10.1029/2011jc007276> (2012).
- 34 Zscheischler, J. et al. Future climate risk from compound events. *Nat. Clim. Change* 8, 469-477, <https://doi.org/10.1038/s41558-018-0156-3> (2018).

Wind

- 1 Feser, F. et al. Storminess over the North Atlantic and north-western Europe—A review. *Q. J. Roy. Meteor. Soc.* 141, 350-382, <https://doi.org/10.1002/qj.2364> (2015).
- 2 Feser, F. et al. 11. Hurricane Gonzalo and its extratropical transition t a strong European storm. *Bull. Amer. Meteor. Soc.* 96, S51-S55 (2015).
- 3 Minola, L., Azorin-Molina, C. & Chen, D. Homogenization and assessment of observed near-surface wind speed trends across Sweden, 1956–2013. *J. Climate* 29, 7397-7415, <https://doi.org/10.1175/JCLI-D-15-0636.1> (2016).
- 4 Laapas, M. & Venäläinen, A. Homogenization and trend analysis of monthly mean and maximum wind speed time series in Finland, 1959–2015. *Int. J. Climatol.* 37, 4803-4813, <https://doi.org/10.1002/joc.5124> (2017).
- 5 Rutgersson, A., Jaagus, J., Schenk, F. & Stendel, M. Observed changes and variability of atmospheric parameters in the Baltic Sea region during the last 200 years. *Clim. Res.* 61, 177-190, <https://doi.org/10.3354/cr01244> (2014).
- 6 Rutgersson, A. et al. in *Second Assessment of Climate Change for the Baltic Sea Basin Regional Climate Studies* (ed BACC II Author Team) 69-97 (Springer International Publishing, 2015).
- 7 Christensen, O. B., Kjellström, E. & Zorita, E. in *Second Assessment of Climate Change for the Baltic Sea Basin* (ed Bacc I. I. Author Team The) 217-233 (Springer International Publishing, 2015).
- 8 Ruosteenoja, K., Vihma, T. & Venäläinen, A. Projected Changes in European and North Atlantic Seasonal Wind Climate Derived from CMIP5 Simulations. *J. Climate* 32, 6467-6490, <https://doi.org/10.1175/JCLI-D-19-0023.1> (2019).
- 9 Räisänen, J. (Oxford University Press, 2017).
- 10 Meier, H. E. M. et al. Quality assessment of atmospheric surface fields over the Baltic Sea from an ensemble of regional climate model simulations with respect to ocean dynamics. *Oceanologia* 53, 193-227, <https://doi.org/10.5697/oc.53-1-TI.193> (2011).
- 11 Rädler, A. T., Groenemeijer, P. H., Faust, E., Sausen, R. & Púčik, T. Frequency of severe thunderstorms across Europe expected to increase in the 21st century due to rising instability. *npj Clim. Atmos. Sci.* 2, 30, <https://doi.org/10.1038/s41612-019-0083-7> (2019).

Waves

- 1 Rutgersson, A. et al. in *Second Assessment of Climate Change for the Baltic Sea Basin Regional Climate Studies* (ed BACC II Author Team) 69-97 (Springer International Publishing, 2015).
- 2 Hünicke, B. et al. in *Second Assessment of Climate Change for the Baltic Sea Basin* (ed BACC II Author Team) 155-185 (Springer International Publishing, 2015).
- 3 Christensen, O. B., Kjellström, E. & Zorita, E. in *Second Assessment of Climate Change for the Baltic Sea Basin* (ed Bacc I. I. Author Team The) 217-233 (Springer International Publishing, 2015).
- 4 Meier, H. E. M. in *Second Assessment of Climate Change for the Baltic Sea Basin* (ed BACC II Author Team) 243-252 (Springer International Publishing, 2015).
- 5 Groll, N., Grabemann, I., Hünicke, B. & Meese, M. Baltic Sea wave conditions under climate change scenarios. *Boreal Environ. Res.* 22, 1-12 (2017).

- 6 Tobin, I. et al. Climate change impacts on the power generation potential of a European mid-century wind farms scenario. *Environ. Res. Lett.* 11, 034013, <https://doi.org/10.1088/1748-9326/11/3/034013> (2016).

Sediment transportation

- 1 Harff, J. et al. in *Coastline changes: interrelation of climate and geological processes* Vol. 426 (eds J. Harff, W. W. Hay, & D.M. Tetzlaff) 133-142 (SPECIAL PAPERS-GEOLOGICAL SOCIETY OF AMERICA, 2007).
- 2 Harff, J. et al. in *Coastline Changes of the Baltic Sea from South to East: Past and Future Projection* (eds Jan Harff, Kazimierz Furmańczyk, & Hans von Storch) 15-35 (Springer International Publishing, 2017).
- 3 Łabuz, T. A. in *Second Assessment of Climate Change for the Baltic Sea Basin* (ed BACC II Author Team) 381-396 (Springer International Publishing, 2015).
- 4 Dangendorf, S. et al. Persistent acceleration in global sea-level rise since the 1960s. *Nat. Clim. Change* 9, 705-710, <https://doi.org/10.1038/s41558-019-0531-8> (2019).
- 5 Zhang, W., Schneider, R., Harff, J., Hünicke, B. & Fröhle, P. in *Coastline Changes of the Baltic Sea from South to East: Past and Future Projection* (eds Jan Harff, Kazimierz Furmańczyk, & Hans von Storch) 107-135 (Springer International Publishing, 2017).
- 6 Musielak, S., Furmańczyk, K. & Bugajny, N. in *Coastline Changes of the Baltic Sea from South to East: Past and Future Projection* (eds Jan Harff, Kazimierz Furmańczyk, & Hans von Storch) 69-85 (Springer International Publishing, 2017).
- 7 Soomere, T., Viška, M. & Pindsoo, K. in *Coastline Changes of the Baltic Sea from South to East: Past and Future Projection* (eds Jan Harff, Kazimierz Furmańczyk, & Hans von Storch) 327-361 (Springer International Publishing, 2017).
- 8 Zhang, W., Harff, J. & Schneider, R. Analysis of 50-year wind data of the southern Baltic Sea for modelling coastal morphological evolution – a case study from the Darss-Zingst Peninsula. *Oceanologia* 53, 489-518, <https://doi.org/10.5697/oc.53-1-TI.489> (2011).
- 9 Tõnisson, H., Orviku, K., Lapinskis, J., Gulbinskas, S. & Žaromskis, R. in *Coastal Erosion and Protection in Europe* (eds Enzo Pranzini & Allan Williams) Ch. 4, 34pp (Routledge, 2013).
- 10 Dudzińska-Nowak, J. in *Coastline Changes of the Baltic Sea from South to East: Past and Future Projection* (eds Jan Harff, Kazimierz Furmańczyk, & Hans von Storch) 219-255 (Springer International Publishing, 2017).
- 11 BACC II Author Team. *Second Assessment of Climate Change for the Baltic Sea Basin*. (Springer International Publishing, 2015).

Oxygen

- 1 Conley, D. J. et al. Hypoxia-Related Processes in the Baltic Sea. *Environ. Sci. Technol.* 43, 3412-3420, <https://doi.org/10.1021/es802762a> (2009).
- 2 BACC II Author Team. *Second Assessment of Climate Change for the Baltic Sea Basin*. (Springer International Publishing, 2015).
- 3 Carstensen, J., Andersen, J. H., Gustafsson, B. G. & Conley, D. J. Deoxygenation of the Baltic Sea during the last century. *P. Natl. Acad. Sci. USA* 111, 5628-5633, <https://doi.org/10.1073/pnas.1323156111> (2014).

- 4 Conley, D. J. et al. Hypoxia Is Increasing in the Coastal Zone of the Baltic Sea. *Environ. Sci. Technol.* 45, 6777-6783, <https://doi.org/10.1021/es201212r> (2011).
- 5 Gustafsson, B. G. et al. Reconstructing the development of Baltic Sea eutrophication 1850-2006. *Ambio* 41, 534-548, <https://doi.org/10.1007/s13280-012-0318-x> (2012).
- 6 Meier, H. E. M., Väli, G., Naumann, M., Eilola, K. & Frauen, C. Recently accelerated oxygen consumption rates amplify deoxygenation in the Baltic Sea. *J. Geophys. Res-Oceans* 123, 3227-3240, <https://doi.org/10.1029/2017JC013686> (2018).
- 7 Meier, H. E. M. et al. Disentangling the impact of nutrient load and climate changes on Baltic Sea hypoxia and eutrophication since 1850. *Clim. Dyn.* 53, 1145-1166, <https://doi.org/10.1007/s00382-018-4296-y> (2019).
- 8 Meier, H. E. M. et al. Correction to: Disentangling the impact of nutrient load and climate changes on Baltic Sea hypoxia and eutrophication since 1850. *Clim. Dyn.* 53, 1167-1169, <https://doi.org/10.1007/s00382-018-4483-x> (2019).
- 9 Savchuk, O. P. in *Chemical Structure of Pelagic Redox Interfaces* 137-160 (Berlin, Heidelberg: Springer, 2010).
- 10 Savchuk, O. P. Large-Scale Nutrient Dynamics in the Baltic Sea, 1970-2016. *Front. Mar. Sci.* 5, 95, <https://doi.org/10.3389/fmars.2018.00095> (2018).
- 11 Andersen, J. H. et al. Long-term temporal and spatial trends in eutrophication status of the Baltic Sea. *Biol. Rev.* 92, 135-149, <https://doi.org/10.1111/bvr.12221> (2017).
- 12 Meier, H. E. M. et al. Future projections of record-breaking sea surface temperature and cyanobacteria bloom events in the Baltic Sea. *Ambio* 48, 1362-1376, <https://doi.org/10.1007/s13280-019-01235-5> (2019).
- 13 Meier, H. E. M. et al. Assessment of Eutrophication Abatement Scenarios for the Baltic Sea by Multi-Model Ensemble Simulations. *Front. Mar. Sci.* 5, <https://doi.org/10.3389/fmars.2018.00440> (2018).
- 14 Meier, H. E. M. et al. Assessment of Uncertainties in Scenario Simulations of Biogeochemical Cycles in the Baltic Sea. *Front. Mar. Sci.* 6, <https://doi.org/10.3389/fmars.2019.00046> (2019).
- 15 Saraiva, S. et al. Baltic Sea ecosystem response to various nutrient load scenarios in present and future climates. *Clim. Dyn.* 52, 3369-3387, <https://doi.org/10.1007/s00382-018-4330-0> (2019).
- 16 Saraiva, S. et al. Uncertainties in Projections of the Baltic Sea Ecosystem Driven by an Ensemble of Global Climate Models. *Front. Earth Sci.* 6, <https://doi.org/10.3389/feart.2018.00244> (2019).
- 17 Carstensen, J. et al. Factors regulating the coastal nutrient filter in the Baltic Sea. *Ambio* 49, 1194-1210, <https://doi.org/10.1007/s13280-019-01282-y> (2020).

Microbial community and –processes

- 1 Herlemann, D. P. R. et al. Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea. *The ISME Journal* 5, 1571-1579, <https://doi.org/10.1038/ismej.2011.41> (2011).
- 2 Shen, D., Jürgens, K. & Beier, S. Experimental insights into the importance of ecologically dissimilar bacteria to community assembly along a salinity gradient. *Environ. Microbiol.* 20, 1170-1184, <https://doi.org/10.1111/1462-2920.14059> (2018).
- 3 Herlemann, D. P. R., Manecki, M., Dittmar, T. & Jürgens, K.



Differential responses of marine, mesohaline and oligohaline bacterial communities to the addition of terrigenous carbon. *Environ. Microbiol.* 19, 3098-3117, <https://doi.org/10.1111/1462-2920.13784> (2017).

4 Hoppe, H. G., Giesenhausen, H. C., Koppe, R., Hansen, H. P. & Gocke, K. Impact of change in climate and policy from 1988 to 2007 on environmental and microbial variables at the time series station Boknis Eck, Baltic Sea. *Biogeosciences* 10, 4529-4546, <https://doi.org/10.5194/bg-10-4529-2013> (2013).

5 Ahlgren, J., Grimvall, A., Omstedt, A., Rolff, C. & Wikner, J. Temperature, DOC level and basin interactions explain the declining oxygen concentrations in the Bothnian Sea. *J. Mar. Syst.* 170, 22-30, <https://doi.org/10.1016/j.jmarsys.2016.12.010> (2017).

6 Huseby, S. Pelagic biology: bacterioplankton (in Swedish). (The Swedish Agency for Marine and Water Management and The Swedish Environmental Protection Agency, 2016).

7 Huseby, S. & Wikner, J. Bacterioplankton growth, <https://helcom.fi/wp-content/uploads/2020/06/BSEFS-Bacterioplankton-growth-.pdf> (last access: 22-06-2021).

8 Semenza, J. C. et al. Environmental Suitability of *Vibrio* Infections in a Warming Climate: An Early Warning System. *Environ. Health Perspect.* 125, 107004, <https://doi.org/10.1289/EHP2198> (2017).

9 Wohlers, J. et al. Changes in biogenic carbon flow in response to sea surface warming. *P. Natl. Acad. Sci. USA* 106, 7067, <https://doi.org/10.1073/pnas.0812743106> (2009).

10 Andersson, A. et al. Projected future climate change and Baltic Sea ecosystem management. *Ambio* 44, 345-356, <https://doi.org/10.1007/s13280-015-0654-8> (2015).

11 Wikner, J. & Andersson, A. Increased freshwater discharge shifts the trophic balance in the coastal zone of the northern Baltic Sea. *Global Change Biol.* 18, 2509-2519, <https://doi.org/10.1111/j.1365-2486.2012.02718.x> (2012).

12 Bertilsson, S. et al. Photochemically induced changes in bioavailable carbon and nitrogen pools in a boreal watershed. *Aquat. Microb. Ecol.* 19, 47-56, <https://doi.org/10.3354/ame019047> (1999).

13 Rodríguez, J. et al. Effects of Organic Pollutants on Bacterial Communities Under Future Climate Change Scenarios. *Front. Microbiol.* 9, <https://doi.org/10.3389/fmicb.2018.02926> (2018).

Benthic habitats

1 Remane, A. & Schlieper, C. Biology of Brackish Water. *Int. Rev. Hydrobiol.* 59, 586 (1974).

2 Olsson, J., Bergström, L. & Gårdmark, A. Top-Down Regulation, Climate and Multi-Decadal Changes in Coastal Zoobenthos Communities in Two Baltic Sea Areas. *PLoS ONE* 8, e64767, <https://doi.org/10.1371/journal.pone.0064767> (2013).

3 Wiklund, A.-K. E., Sundelin, B. & Rosa, R. Population decline of amphipod *Monoporeia affinis* in Northern Europe: consequence of food shortage and competition? *J. Exp. Mar. Biol. Ecol.* 367, 81-90, <https://doi.org/10.1016/j.jembe.2008.08.018> (2008).

4 Rousi, H. et al. Long-term changes in coastal zoobenthos in the northern Baltic Sea: the role of abiotic environmental factors. *ICES J. Mar. Sci.* 70, 440-451, <https://doi.org/10.1093/icesjms/fss197> (2013).

5 Eriksson Wiklund, A. K., Dahlgren, K., Sundelin, B. & Anders-

son, A. Effects of warming and shifts of pelagic food web structure on benthic productivity in a coastal marine system. *Mar. Ecol. Prog. Ser.* 396, 13-25, <https://doi.org/10.3354/meps08290> (2009).

6 Carstensen, J., Andersen, J. H., Gustafsson, B. G. & Conley, D. J. Deoxygenation of the Baltic Sea during the last century. *P. Natl. Acad. Sci. USA* 111, 5628-5633, <https://doi.org/10.1073/pnas.1323156111> (2014).

7 Conley, D. J. et al. Hypoxia Is Increasing in the Coastal Zone of the Baltic Sea. *Environ. Sci. Technol.* 45, 6777-6783, <https://doi.org/10.1021/es201212r> (2011).

8 Laine, A. O., Sandler, H., Andersin, A.-B. & Stigzelius, J. Long-term changes of macrozoobenthos in the Eastern Gotland Basin and the Gulf of Finland (Baltic Sea) in relation to the hydrographical regime. *J. Sea Res.* 38, 135-159, [https://doi.org/10.1016/S1385-1101\(97\)00034-8](https://doi.org/10.1016/S1385-1101(97)00034-8) (1997).

9 Altieri, A. H. & Gedan, K. B. Climate change and dead zones. *Global Change Biol.* 21, 1395-1406, <https://doi.org/10.1111/gcb.12754> (2015).

10 Carstensen, J. et al. Hypoxia in the Baltic Sea: Biogeochemical cycles, benthic fauna, and management. *Ambio* 43, 26-36, <https://doi.org/10.1007/s13280-013-0474-7> (2014).

11 Elmgren, R. Man's Impact on the Ecosystem of the Baltic Sea: Energy Flows Today and at the Turn of the Century. *Ambio* 18, 326-332 (1989).

12 Takolander, A., Leskinen, E. & Cabeza, M. Synergistic effects of extreme temperature and low salinity on foundational macroalga *Fucus vesiculosus* in the northern Baltic Sea. *J. Exp. Mar. Biol. Ecol.* 495, 110-118, <https://doi.org/10.1016/j.jembe.2017.07.001> (2017).

13 Westerborn, M., Mustonen, O., Jaatinen, K., Kilpi, M. & Norkko, A. Population Dynamics at the Range Margin: Implications of Climate Change on Sublittoral Blue Mussels (*Mytilus trossulus*). *Front. Mar. Sci.* 6, <https://doi.org/10.3389/fmars.2019.00292> (2019).

14 Dippner, J. W., Hänninen, J., Kuosa, H. & Vuorinen, I. The influence of climate variability on zooplankton abundance in the Northern Baltic Archipelago Sea (SW Finland). *ICES J. Mar. Sci.* 58, 569-578, <https://doi.org/10.1006/jmsc.2001.1048> (2001).

15 Westerborn, M., Kilpi, M. & Mustonen, O. Blue mussels, *Mytilus edulis*, at the edge of the range: population structure, growth and biomass along a salinity gradient in the north-eastern Baltic Sea. *Mar. Biol.* 140, 991-999, <https://doi.org/10.1007/s00227-001-0765-6> (2002).

16 Kautsky, N. Quantitative studies on gonad cycle, fecundity, reproductive output and recruitment in a baltic *Mytilus edulis* population. *Mar. Biol.* 68, 143-160, <https://doi.org/10.1007/BF00397601> (1982).

17 Kovtun, A. et al. Influence of abiotic environmental conditions on spatial distribution of charophytes in the coastal waters of West Estonian Archipelago, Baltic Sea. *J. Coastal Res.*, 412-416 (2011).

18 Ehlers, A., Worm, B. & Reusch, T. B. H. Importance of genetic diversity in eelgrass *Zostera marina* for its resilience to global warming. *Mar. Ecol. Prog. Ser.* 355, 1-7, <https://doi.org/10.3354/meps07369> (2008).

19 Jonsson, P. R. et al. High climate velocity and population fragmentation may constrain climate-driven range shift of the key habitat former *Fucus vesiculosus*. *Divers. Distrib.* 24, 892-905, <https://doi.org/10.1111/ddi.12733> (2018).

20 Kotta, J. et al. Integrating experimental and distribution data

to predict future species patterns. *Sci. Rep.-UK* 9, 1821, <https://doi.org/10.1038/s41598-018-38416-3> (2019).

21 Andersson, A. et al. Projected future climate change and Baltic Sea ecosystem management. *Ambio* 44, 345-356, <https://doi.org/10.1007/s13280-015-0654-8> (2015).

22 Beaugrand, G. The North Sea regime shift: Evidence, causes, mechanisms and consequences. *Prog. Oceanogr.* 60, 245-262, <https://doi.org/10.1016/j.pocean.2004.02.018> (2004).

23 Hiebenthal, C., Philipp, E. E. R., Eisenhauer, A. & Wahl, M. Effects of seawater pCO₂ and temperature on shell growth, shell stability, condition and cellular stress of Western Baltic Sea *Mytilus edulis* (L.) and *Arctica islandica* (L.). *Mar. Biol.* 160, 2073-2087, <https://doi.org/10.1007/s00227-012-2080-9> (2013).

24 Graiff, A., Liesner, D., Karsten, U. & Bartsch, I. Temperature tolerance of western Baltic Sea *Fucus vesiculosus* – growth, photosynthesis and survival. *J. Exp. Mar. Biol. Ecol.* 471, 8-16, <https://doi.org/10.1016/j.jembe.2015.05.009> (2015).

25 Wikner, J. & Andersson, A. Increased freshwater discharge shifts the trophic balance in the coastal zone of the northern Baltic Sea. *Global Change Biol.* 18, 2509-2519, <https://doi.org/10.1111/j.1365-2486.2012.02718.x> (2012).

26 Kotta, J., Kotta, I., Simm, M. & Pöllupüü, M. Separate and interactive effects of eutrophication and climate variables on the ecosystem elements of the Gulf of Riga. *Estuar. Coast. Shelf Sci.* 84, 509-518, <https://doi.org/10.1016/j.ecss.2009.07.014> (2009).

27 Grinstead, A. in Second Assessment of Climate Change for the Baltic Sea Basin Regional Climate Studies (ed BACC II Author Team) 253-263 (Springer International Publishing, 2015).

28 Ehrnsten, E., Bauer, B. & Gustafsson, B. G. Combined Effects of Environmental Drivers on Marine Trophic Groups – A Systematic Model Comparison. *Front. Mar. Sci.* 6, <https://doi.org/10.3389/fmars.2019.00492> (2019).

29 Weigel, B. et al. Long-term progression and drivers of coastal zoobenthos in a changing system. *Mar. Ecol. Prog. Ser.* 528, 141-159, <https://doi.org/10.3354/meps11279> (2015).

30 Breitburg, D. et al. Declining oxygen in the global ocean and coastal waters. *Science* 359, eaam7240, <https://doi.org/10.1126/science.aam7240> (2018).

31 Räike, A., Taskinen, A. & Knuuttila, S. Nutrient export from Finnish rivers into the Baltic Sea has not decreased despite water protection measures. *Ambio* 49, 460-474, <https://doi.org/10.1007/s13280-019-01217-7> (2020).

32 Ehrnsten, E., Norkko, A., Müller-Karulis, B., Gustafsson, E. & Gustafsson, B. G. The meagre future of benthic fauna in a coastal sea—Benthic responses to recovery from eutrophication in a changing climate. *Global Change Biol.* 26, 2235-2250, <https://doi.org/10.1111/gcb.15014> (2020).

33 Holopainen, R. et al. Impacts of changing climate on the non-indigenous invertebrates in the northern Baltic Sea by end of the twenty-first century. *Biol. Invasions* 18, 3015-3032, <https://doi.org/10.1007/s10530-016-1197-z> (2016).

34 Norkko, J. et al. A welcome can of worms? Hypoxia mitigation by an invasive species. *Global Change Biol.* 18, 422-434, <https://doi.org/10.1111/j.1365-2486.2011.02513.x> (2012).

35 Sandman, A. N., Näslund, J., Gren, I.-M. & Norling, K. Effects of an invasive polychaete on benthic phosphorus cycling at sea basin scale: An ecosystem disservice. *Ambio* 47, 884-892, <https://doi.org/10.1007/s13280-018-1050-y> (2018).

36 Kotta, J. et al. Novel crab predator causes marine ecosystem regime shift. *Sci. Rep.-UK* 8, 4956, <https://doi.org/10.1038/s41598-018-23282-w> (2018).

37 Casini, M. et al. Trophic cascades promote threshold-like shifts in pelagic marine ecosystems. *P. Natl. Acad. Sci. USA* 106, 197, <https://doi.org/10.1073/pnas.0806649105> (2009).

38 Casini, M. et al. Multi-level trophic cascades in a heavily exploited open marine ecosystem. *P. Roy. Soc. B-Biol. Sci.* 275, 1793-1801, <https://doi.org/10.1098/rspb.2007.1752> (2008).

39 Östman, Ö. et al. Top-down control as important as nutrient enrichment for eutrophication effects in North Atlantic coastal ecosystems. *J. Appl. Ecol.* 53, 1138-1147, <https://doi.org/10.1111/1365-2664.12654> (2016).

40 Ljunggren, L. et al. Recruitment failure of coastal predatory fish in the Baltic Sea coincident with an offshore ecosystem regime shift. *ICES J. Mar. Sci.* 67, 1587-1595, <https://doi.org/10.1093/icesjms/fsq109> (2010).

41 Saraiva, S. et al. Uncertainties in Projections of the Baltic Sea Ecosystem Driven by an Ensemble of Global Climate Models. *Front. Earth Sci.* 6, <https://doi.org/10.3389/feart.2018.00244> (2019).

42 Meier, H. E. M., Dieterich, C. & Gröger, M. Natural variability is a large source of uncertainty in future projections of hypoxia in the Baltic Sea. *Commun. Earth Environ.* 2, 50, <https://doi.org/10.1038/s43247-021-00115-9> (2021).

43 Alheit, J. et al. Synchronous ecological regime shifts in the central Baltic and the North Sea in the late 1980s. *ICES J. Mar. Sci.* 62, 1205-1215, <https://doi.org/10.1016/j.icesjms.2005.04.024> (2005).

44 Rothäusler, E., Rugiu, L. & Jormalainen, V. Forecast climate change conditions sustain growth and physiology but hamper reproduction in range-margin populations of a foundation rockweed species. *Mar. Environ. Res.* 141, 205-213, <https://doi.org/10.1016/j.marenvres.2018.09.014> (2018).

45 Rugiu, L., Manninen, I., Rothäusler, E. & Jormalainen, V. Tolerance and potential for adaptation of a Baltic Sea rockweed under predicted climate change conditions. *Mar. Environ. Res.* 134, 76-84, <https://doi.org/10.1016/j.marenvres.2017.12.016> (2018).

46 Rugiu, L., Manninen, I., Rothäusler, E. & Jormalainen, V. Tolerance to climate change of the clonally reproducing endemic Baltic seaweed, *Fucus radicans*: is phenotypic plasticity enough? *J. Phycol.* 54, 888-898, <https://doi.org/10.1111/jpy.12796> (2018).

47 European Commission. DIRECTIVE 2008/56/EC OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL of 17 June 2008, establishing a framework for community action in the field of marine environmental policy (Marine Strategy Framework Directive) (Text with EEA relevance). (2008).

48 European Commission. COMMISSION DECISION (EU) 2017/848 of 17 May 2017 laying down criteria and methodological standards on good environmental status of marine waters and specifications and standardised methods for monitoring and assessment, and repealing Decision 2010/477/EU., (2017).

49 European Commission. COUNCIL DIRECTIVE 92/43/EEC of 21 May 1992 on the conservation of natural habitats and of wild fauna and flora., (1992).

50 Nyström Sandman, A. et al. Grön infrastruktur i havet : landskapsperspektiv i förvaltningen av Sveriges marina områden. Report No. Naturvårdsverket 6930, 130pp (2020).

51 Frazão Santos, C. et al. Integrating climate change in ocean planning. *Nat. Sustain.* 3, 505-516, <https://doi.org/10.1038/s41893-020-0513-x> (2020).





Coastal and migratory fish

- 1 HELCOM. State of the Baltic Sea - Second HELCOM holistic assessment 2011-2016. 155pp (Helsinki, Finland, 2018).
- 2 Böhling, P. et al. Variations in Year-Class Strength of Different Perch (*Perca fluviatilis*) Populations in the Baltic Sea with Special Reference to Temperature and Pollution. *Can. J. Fish. Aquat. Sci.* 48, 1181-1187, <https://doi.org/10.1139/f91-142> (1991).
- 3 Karäs, P. & Thoresson, G. An application of a bioenergetics model to Eurasian perch (*Perca fluviatilis* L.). *J. Fish Biol.* 41, 217-230, <https://doi.org/10.1111/j.1095-8649.1992.tb02652.x> (1992).
- 4 Lehtonen, H. & Lappalainen, J. The effects of climate on the year-class variations of certain freshwater fish species. *Canadian Special Publication of Fisheries and Aquatic Sciences* 121, 37-44 (1995).
- 5 Karäs, P. Recruitment of perch (*Perca fluviatilis* L.) from Baltic coastal waters. *Arch. Hydrobiol.* 138, 99-121, <https://doi.org/10.1127/archiv-hydrobiol/138/1996/99> (1996).
- 6 Kjellman, J., Lappalainen, J. & Urho, L. Influence of temperature on size and abundance dynamics of age-0 perch and pikeperch. *Fish. Res.* 53, 47-56, [https://doi.org/10.1016/S0165-7836\(00\)00265-4](https://doi.org/10.1016/S0165-7836(00)00265-4) (2001).
- 7 Heikinheimo, O., Pekcan-Hekim, Z. & Raitaniemi, J. Spawning stock-recruitment relationship in pikeperch *Sander lucioperca* (L.) in the Baltic Sea, with temperature as an environmental effect. *Fish. Res.* 155, 1-9, <https://doi.org/10.1016/j.fishres.2014.02.015> (2014).
- 8 Kokkonen, E., Heikinheimo, O., Pekcan-Hekim, Z. & Vainikka, A. Effects of water temperature and pikeperch (*Sander lucioperca*) abundance on the stock-recruitment relationship of Eurasian perch (*Perca fluviatilis*) in the northern Baltic Sea. *Hydrobiologia* 841, 79-94, <https://doi.org/10.1007/s10750-019-04008-z> (2019).
- 9 Pekcan-Hekim, Z. et al. Climate Warming and Pikeperch Year-Class Catches in the Baltic Sea. *Ambio* 40, 447-456, <https://doi.org/10.1007/s13280-011-0143-7> (2011).
- 10 Candolin, U. & Voigt, H.-R. Population growth correlates with increased fecundity in three-spined stickleback populations in a human-disturbed environment. *Aquat. Sci.* 82, 21, <https://doi.org/10.1007/s00027-020-0695-3> (2020).
- 11 Kallio-Nyberg, I., Veneranta, L., Saloniemi, I., Jokikokko, E. & Leskelä, A. Different growth trends of whitefish (*Coregonus lavaretus*) forms in the northern Baltic Sea. *J. Appl. Ichthyol.* 35, 683-691, <https://doi.org/10.1111/jai.13898> (2019).
- 12 Veneranta, L., Hudd, R. & Vanhatalo, J. Reproduction areas of sea-spawning coregonids reflect the environment in shallow coastal waters. *Mar. Ecol. Prog. Ser.* 477, 231-250, <https://doi.org/10.3354/meps10169> (2013).
- 13 Bergenius, M. A., Gårdmark, A., Ustups, D., Kaljuste, O. & Aho, T. Fishing or the environment—what regulates recruitment of an exploited marginal vendace (*Coregonus albula* (L.)) population? *Adv. Limnol.* 64, 57-70, <https://doi.org/10.1127/1612-166X/2013/0064-0029> (2013).
- 14 Veneranta, L., Urho, L., Koho, J. & Hudd, R. Spawning and hatching temperatures of whitefish (*Coregonus lavaretus* (L.)) in the Northern Baltic Sea. *Adv. Limnol.* 64, 39-55, <https://doi.org/10.1127/1612-166X/2013/0064-0019> (2013).
- 15 Olsson, J., Bergström, L. & Gårdmark, A. Abiotic drivers of coastal fish community change during four decades in the Bal-

- tic Sea. *ICES J. Mar. Sci.* 69, 961-970, <https://doi.org/10.1093/icesjms/fss072> (2012).
- 16 Dahl, J. et al. The timing of spawning migration: implications of environmental variation, life history, and sex. *Can. J. Zool.* 82, 1864-1870, <https://doi.org/10.1139/z04-184> (2004).
 - 17 Kallio-Nyberg, I., Jutila, E., Saloniemi, I. & Jokikokko, E. Association between environmental factors, smolt size and the survival of wild and reared Atlantic salmon from the Simojoki River in the Baltic Sea. *J. Fish Biol.* 65, 122-134, <https://doi.org/10.1111/j.1095-8649.2004.00435.x> (2004).
 - 18 Kallio-Nyberg, I., Jutila, E., Jokikokko, E. & Saloniemi, I. Survival of reared Atlantic salmon and sea trout in relation to marine conditions of smolt year in the Baltic Sea. *Fish. Res.* 80, 295-304, <https://doi.org/10.1016/j.fishres.2006.03.026> (2006).
 - 19 Friedland, K. D. et al. Post-smolt survival of Baltic salmon in context to changing environmental conditions and predators. *ICES J. Mar. Sci.* 74, 1344-1355, <https://doi.org/10.1093/icesjms/fsw178> (2017).
 - 20 Tamario, C., Sunde, J., Petersson, E., Tibblin, P. & Forsman, A. Ecological and Evolutionary Consequences of Environmental Change and Management Actions for Migrating Fish. *Front. Ecol. Evol.* 7, <https://doi.org/10.3389/fevo.2019.00271> (2019).
 - 21 Härmä, M., Lappalainen, A. & Urho, L. Reproduction areas of roach (*Rutilus rutilus*) in the northern Baltic Sea: potential effects of climate change. *Can. J. Fish. Aquat. Sci.* 65, 2678-2688, <https://doi.org/10.1139/f08-167> (2008).
 - 22 Dainys, J. et al. Predicted climate change effects on European perch (*Perca fluviatilis* L.) - A case study from the Curonian Lagoon, south-eastern Baltic. *Estuar. Coast. Shelf Sci.* 221, 83-89, <https://doi.org/10.1016/j.ecss.2019.03.020> (2019).
 - 23 Östman, Ö., Boström, M. K., Bergström, U., Andersson, J. & Lunneryd, S.-G. Estimating Competition between Wildlife and Humans—A Case of Cormorants and Coastal Fisheries in the Baltic Sea. *PLoS ONE* 8, e83763, <https://doi.org/10.1371/journal.pone.0083763> (2014).
 - 24 Huss, M., Lindmark, M., Jacobson, P., van Dorst, R. M. & Gårdmark, A. Experimental evidence of gradual size-dependent shifts in body size and growth of fish in response to warming. *Global Change Biol.* 25, 2285-2295, <https://doi.org/10.1111/gcb.14637> (2019).
 - 25 van Dorst, R. M. et al. Warmer and browner waters decrease fish biomass production. *Global Change Biol.* 25, 1395-1408, <https://doi.org/10.1111/gcb.14551> (2019).
 - 26 Hansson, S. et al. Competition for the fish – fish extraction from the Baltic Sea by humans, aquatic mammals, and birds. *ICES J. Mar. Sci.* 75, 999-1008, <https://doi.org/10.1093/icesjms/fsx207> (2018).

Pelagic and demersal fish

- 1 ICES. Baltic Sea Ecosystem - Fisheries Overview. Report of the ICES Advisory Committee ICES Advice 2019, section 4.2, 29pp, <https://doi.org/10.17895/ices.advice.5566> (2019).
- 2 Rau, A., Lewin, W.-C., Zettler, M. L., Gogina, M. & von Dorrien, C. Abiotic and biotic drivers of flatfish abundance within distinct demersal fish assemblages in a brackish ecosystem (western Baltic Sea). *Estuar. Coast. Shelf Sci.* 220, 38-47, <https://doi.org/10.1016/j.ecss.2019.02.035> (2019).
- 3 Dinesen, G. E. et al. Cod and climate: a systems approach for sustainable fisheries management of Atlantic cod (*Gadus*

- morhua*) in coastal Danish waters. *J. Coast. Conserv.* 23, 943-958, <https://doi.org/10.1007/s11852-019-00711-0> (2019).
- 4 Andersen, N. G., Lundgren, B., Neuenfeldt, S. & Beyer, J. E. Diel vertical interactions between Atlantic cod *Gadus morhua* and sprat *Sprattus sprattus* in a stratified water column. *Mar. Ecol. Prog. Ser.* 583, 195-209 (2017).
 - 5 Orio, A. et al. Spatial contraction of demersal fish populations in a large marine ecosystem. *J. Biogeogr.* 46, 633-645, <https://doi.org/10.1111/jbi.13510> (2019).
 - 6 Solberg, I., Røstad, A. & Kaartvedt, S. Ecology of overwintering sprat (*Sprattus sprattus*). *Prog. Oceanogr.* 138, 116-135, <https://doi.org/10.1016/j.pocean.2015.08.003> (2015).
 - 7 Neuenfeldt, S. The influence of oxygen saturation on the distributional overlap of predator (cod, *Gadus morhua*) and prey (herring, *Clupea harengus*) in the Bornholm Basin of the Baltic Sea. *Fish. Oceanogr.* 11, 11-17, <https://doi.org/10.1046/j.1365-2419.2002.00183.x> (2002).
 - 8 Neuenfeldt, S. & Beyer, J. E. Oxygen and salinity characteristics of predator-prey distributional overlaps shown by predatory Baltic cod during spawning. *J. Fish Biol.* 62, 168-183, <https://doi.org/10.1046/j.1095-8649.2003.00013.x> (2003).
 - 9 Neuenfeldt, S., Andersen, K. H. & Hinrichsen, H. H. Some Atlantic cod *Gadus morhua* in the Baltic Sea visit hypoxic water briefly but often. *J. Fish Biol.* 75, 290-294, <https://doi.org/10.1111/j.1095-8649.2009.02281.x> (2009).
 - 10 Limburg, K. E. & Casini, M. Otolith chemistry indicates recent worsened Baltic cod condition is linked to hypoxia exposure. *Biol. Lett.* 15, 20190352, <https://doi.org/10.1098/rsbl.2019.0352> (2019).
 - 11 Lefébure, R., Larsson, S. & Byström, P. A temperature-dependent growth model for the three-spined stickleback *Gasterosteus aculeatus*. *J. Fish Biol.* 79, 1815-1827, <https://doi.org/10.1111/j.1095-8649.2011.03121.x> (2011).
 - 12 Lefébure, R., Larsson, S. & Byström, P. Temperature and size-dependent attack rates of the three-spined stickleback (*Gasterosteus aculeatus*); are sticklebacks in the Baltic Sea resource-limited? *J. Exp. Mar. Biol. Ecol.* 451, 82-90, <https://doi.org/10.1016/j.jembe.2013.11.008> (2014).
 - 13 Nissling, A. & Westin, L. Salinity requirements for successful spawning of Baltic and Belt Sea cod and the potential for cod stock interactions in the Baltic Sea. *Mar. Ecol. Prog. Ser.* 152, 261-271 (1997).
 - 14 Nissling, A., Westin, L. & Hjerne, O. Reproductive success in relation to salinity for three flatfish species, dab (*Limanda limanda*), plaice (*Pleuronectes platessa*), and flounder (*Pleuronectes flesus*), in the brackish water Baltic Sea. *ICES J. Mar. Sci.* 59, 93-108, <https://doi.org/10.1006/jmsc.2001.1134> (2002).
 - 15 Peterreit, C., Hinrichsen, H. H., Franke, A. & Köster, F. W. Floating along buoyancy levels: Dispersal and survival of western Baltic fish eggs. *Prog. Oceanogr.* 122, 131-152, <https://doi.org/10.1016/j.pocean.2014.01.001> (2014).
 - 16 Illing, B., Moyano, M., Hufnagl, M. & Peck, M. A. Projected habitat loss for Atlantic herring in the Baltic Sea. *Mar. Environ. Res.* 113, 164-173, <https://doi.org/10.1016/j.marenvres.2015.12.007> (2016).
 - 17 Voss, R. et al. Egg mortality: predation and hydrography in the central Baltic. *ICES J. Mar. Sci.* 68, 1379-1390, <https://doi.org/10.1093/icesjms/fsr061> (2011).
 - 18 Hinrichsen, H. H., Hüsey, K. & Huwer, B. Spatio-temporal variability in western Baltic cod early life stage survival mediated by egg buoyancy, hydrography and hydrodynamics. *ICES J.*

- Mar. Sci.* 69, 1744-1752, <https://doi.org/10.1093/icesjms/fss137> (2012).
- 19 Rajasilta, M. et al. Influence of environmental conditions, population density, and prey type on the lipid content in Baltic herring (*Clupea harengus membras*) from the northern Baltic Sea. *Can. J. Fish. Aquat. Sci.* 76, 576-585, <https://doi.org/10.1139/cjfas-2017-0504> (2018).
 - 20 Flinkman, J., Aro, E., Vuorinen, I. & Viitasalo, M. Changes in northern Baltic zooplankton and herring nutrition from 1980s to 1990s: top-down and bottom-up processes at work. *Mar. Ecol. Prog. Ser.* 165, 127-136 (1998).
 - 21 Rönkkönen, S., Ojaveer, E., Raid, T. & Viitasalo, M. Long-term changes in Baltic herring (*Clupea harengus membras*) growth in the Gulf of Finland. *Can. J. Fish. Aquat. Sci.* 61, 219-229, <https://doi.org/10.1139/f03-167> (2004).
 - 22 Casini, M., Bartolino, V., Molinero, J. C. & Kornilovs, G. Linking fisheries, trophic interactions and climate: threshold dynamics drive herring *Clupea harengus* growth in the central Baltic Sea. *Mar. Ecol. Prog. Ser.* 413, 241-252 (2010).
 - 23 Köster, F. W. et al. Eastern Baltic cod recruitment revisited—dynamics and impacting factors. *ICES J. Mar. Sci.* 74, 3-19, <https://doi.org/10.1093/icesjms/fsw172> (2016).
 - 24 MacKenzie, B. R. & Köster, F. W. Fish production and climate: sprat in the Baltic Sea. *Ecology* 85, 784-794, <https://doi.org/10.1890/02-0780> (2004).
 - 25 Polte, P. et al. Reduced Reproductive Success of Western Baltic Herring (*Clupea harengus*) as a Response to Warming Winters. *Front. Mar. Sci.* 8, <https://doi.org/10.3389/fmars.2021.589242> (2021).
 - 26 Nissling, A. in *Biology of the Baltic Sea*. (eds Hans Kautsky & Pauli Snoeijs) 115-123 (Springer Netherlands).
 - 27 Baumann, H. et al. Recruitment variability in Baltic Sea sprat (*Sprattus sprattus*) is tightly coupled to temperature and transport patterns affecting the larval and early juvenile stages. *Can. J. Fish. Aquat. Sci.* 63, 2191-2201, <https://doi.org/10.1139/f06-112> (2006).
 - 28 Nissling, A., Johansson, U. & Jacobsson, M. Effects of salinity and temperature conditions on the reproductive success of turbot (*Scophthalmus maximus*) in the Baltic Sea. *Fish. Res.* 80, 230-238, <https://doi.org/10.1016/j.fishres.2006.04.005> (2006).
 - 29 Hakala, T. et al. Temporal and spatial variation in the growth rates of Baltic herring (*Clupea harengus membras* L.) larvae during summer. *Mar. Biol.* 142, 25-33, <https://doi.org/10.1007/s00227-002-0933-3> (2003).
 - 30 Wieland, K., Jarre-Teichmann, A. & Horbowa, K. Changes in the timing of spawning of Baltic cod: possible causes and implications for recruitment. *ICES J. Mar. Sci.* 57, 452-464, <https://doi.org/10.1006/jmsc.1999.0522> (2000).
 - 31 Dodson, J. J. et al. Environmental determinants of larval herring (*Clupea harengus*) abundance and distribution in the western Baltic Sea. *Limnol. Oceanogr.* 64, 317-329, <https://doi.org/10.1002/lno.11042> (2019).
 - 32 Saraiva, S. et al. Baltic Sea ecosystem response to various nutrient load scenarios in present and future climates. *Clim. Dyn.* 52, 3369-3387, <https://doi.org/10.1007/s00382-018-4330-0> (2019).
 - 33 Schaber, M., Hinrichsen, H.-H. & Gröger, J. Seasonal changes in vertical distribution patterns of cod (*Gadus morhua*) in the Bornholm Basin, central Baltic Sea. *Fish. Oceanogr.* 21, 33-43, <https://doi.org/10.1111/j.1365-2419.2011.00607.x> (2012).





- 34 Bauer, B. et al. Food web and fisheries in the future Baltic Sea. *Ambio*, <https://doi.org/10.1007/s13280-019-01229-3> (2019).
- 35 Niiranen, S. et al. Combined effects of global climate change and regional ecosystem drivers on an exploited marine food web. *Global Change Biol.* 19, 3327-3342, <https://doi.org/10.1111/gcb.12309> (2013).
- 36 Österblom, H. et al. Human-induced Trophic Cascades and Ecological Regime Shifts in the Baltic Sea. *Ecosystems* 10, 877-889, <https://doi.org/10.1007/s10021-007-9069-0> (2007).
- 37 Aneer, G. Some Speculations about the Baltic Herring (*Clupea harengus membras*) in Connection with the Eutrophication of the Baltic Sea. *Can. J. Fish. Aquat. Sci.* 42, s83-s90, <https://doi.org/10.1139/f85-264> (1985).
- 38 Möllmann, C. et al. Reorganization of a large marine ecosystem due to atmospheric and anthropogenic pressure: a discontinuous regime shift in the Central Baltic Sea. *Global Change Biol.* 15, 1377-1393, <https://doi.org/10.1111/j.1365-2486.2008.01814.x> (2009).
- 39 Pekcan-Hekim, Z. et al. The role of climate and fisheries on the temporal changes in the Bothnian Bay foodweb. *ICES J. Mar. Sci.* 73, 1739-1749, <https://doi.org/10.1093/icesjms/fsw032> (2016).
- 40 Sparrevojn, C. R., Lindegren, M. & Mackenzie, B. R. Climate-induced response of commercially important flatfish species during the 20th century. *Fish. Oceanogr.* 22, 400-408, <https://doi.org/10.1111/fog.12030> (2013).
- 41 Jarre-Teichmann, A. et al. Stock-recruitment relationships for cod (*Gadus morhua callarias* L.) in the central Baltic Sea incorporating environmental variability. *Archive of Fishery and Marine Research* 48, 97-123 (2000).
- Estuary. *PeerJ* 5, e3604, <https://doi.org/10.7717/peerj.3604> (2017).
- 9 Fox, A. D., Nielsen, R. D. & Petersen, I. K. Climate-change not only threatens bird populations but also challenges our ability to monitor them. *Ibis* 161, 467-474, <https://doi.org/10.1111/ibi.12675> (2019).
- 10 Pavón-Jordán, D. et al. Habitat- and species-mediated short- and long-term distributional changes in waterbird abundance linked to variation in European winter weather. *Divers. Distrib.* 25, 225-239, <https://doi.org/10.1111/ddi.12855> (2019).
- 11 Vähätalo, A. V., Rainio, K., Lehikoinen, A. & Lehikoinen, E. Spring arrival of birds depends on the North Atlantic Oscillation. *J. Avian Biol.* 35, 210-216, <https://doi.org/10.1111/j.0908-8857.2004.03199.x> (2004).
- 12 Rainio, K., Laaksonen, T., Ahola, M., Vähätalo, A. V. & Lehikoinen, E. Climatic responses in spring migration of boreal and arctic birds in relation to wintering area and taxonomy. *J. Avian Biol.* 37, 507-515, <https://doi.org/10.1111/j.0908-8857.2006.03740.x> (2006).
- 13 Lehikoinen, A., Kilpi, M. & Öst, M. Winter climate affects subsequent breeding success of common eiders. *Global Change Biol.* 12, 1355-1365, <https://doi.org/10.1111/j.1365-2486.2006.01162.x> (2006).
- 14 Waldeck, P. & Larsson, K. Effects of winter water temperature on mass loss in Baltic blue mussels: Implications for foraging sea ducks. *J. Exp. Mar. Biol. Ecol.* 444, 24-30, <https://doi.org/10.1016/j.jembe.2013.03.007> (2013).
- 15 Larsson, K. et al. Effects of an extensive Prynmesium polylepis bloom on breeding eiders in the Baltic Sea. *J. Sea Res.* 88, 21-28, <https://doi.org/10.1016/j.seares.2013.12.017> (2014).
- 16 Fox, A. D. et al. Current and Potential Threats to Nordic Duck Populations — A Horizon Scanning Exercise. *Annal. Zool.* 52, 193-220, 128 (2015).
- 17 Clausen, K. K., Stjernholm, M. & Clausen, P. Grazing management can counteract the impacts of climate change-induced sea level rise on salt marsh-dependent waterbirds. *J. Appl. Ecol.* 50, 528-537, <https://doi.org/10.1111/1365-2664.12043> (2013).
- 18 Clausen, K. K. & Clausen, P. Forecasting future drowning of coastal waterbird habitats reveals a major conservation concern. *Biol. Conserv.* 171, 177-185, <https://doi.org/10.1016/j.biocon.2014.01.033> (2014).
- 19 Mikola, J., Miettinen, M., Lehikoinen, E. & Lehtilä, K. The effects of disturbance caused by boating on survival and behaviour of velvet scoter *Melanitta fusca* ducklings. *Biol. Conserv.* 67, 119-124, [https://doi.org/10.1016/0006-3207\(94\)90356-5](https://doi.org/10.1016/0006-3207(94)90356-5) (1994).
- 20 Berndt, R., Hein, K., Koop, B. & Lunk, S. Die Vögel der Insel Fehmarn. Husum Druck-u. Verlagsgesellschaft (2005).
- 21 Hario, M. Mink predation on black guillemots at Söderskär in 1994–1999. *Suomen Riista* 48, 18-26 (2002).
- 22 Nordström, M. et al. Effects of feral mink removal on seabirds, waders and passerines on small islands in the Baltic Sea. *Biol. Conserv.* 109, 359-368, [https://doi.org/10.1016/S0006-3207\(02\)00162-3](https://doi.org/10.1016/S0006-3207(02)00162-3) (2003).
- 23 Kube, J., Brenning, U., Kruch, W. & Nehls, H. Bestandsentwicklung von bodenbrütenden Küstenvögeln auf Inseln in der Wismar-Bucht (südwestliche Ostsee): Lektionen aus 50 Jahren Prädatorenmanagement. *Vogelwelt* 126, 299-320 (2005).
- 24 Herrmann, C. et al. in *HELCOM Red List of Baltic Sea species in danger of becoming extinct* Vol. No. 140 Baltic Sea Environment Proceedings (eds Tytti Kontula & Jannica Haldin) 74-89

Waterbirds

- 1 Green, A. J. & Elmsberg, J. Ecosystem services provided by waterbirds. *Biol. Rev.* 89, 105-122, <https://doi.org/10.1111/brv.12045> (2014).
- 2 MacLean, I. M. D. et al. Climate change causes rapid changes in the distribution and site abundance of birds in winter. *Global Change Biol.* 14, 2489-2500, <https://doi.org/10.1111/j.1365-2486.2008.01666.x> (2008).
- 3 Skov, H. et al. Waterbird populations and pressures in the Baltic Sea. Vol. 550 (Nordic Council of Ministers, 2011).
- 4 Aarvak, T., Jostein Ølen, I., Krasnov, Y. V., Gavrilov, M. V. & Shavynkin, A. A. The European wintering population of Steller's Eider *Polysticta stelleri* reassessed. *Bird Conserv. Int.* 23, 337-343, <https://doi.org/10.1017/S0959270912000251> (2013).
- 5 Lehikoinen, A. et al. Rapid climate driven shifts in wintering distributions of three common waterbird species. *Global Change Biol.* 19, 2071-2081, <https://doi.org/10.1111/gcb.12200> (2013).
- 6 Pavón-Jordán, D. et al. Climate-driven changes in winter abundance of a migratory waterbird in relation to EU protected areas. *Divers. Distrib.* 21, 571-582, <https://doi.org/10.1111/ddi.12300> (2015).
- 7 Nilsson, L. & Haas, F. Distribution and numbers of wintering waterbirds in Sweden in 2015 and changes during the last fifty years. *Ornis Svecica* 26, 3-54-53-54 (2016).
- 8 Marchowski, D., Jankowiak, Ł., Wysocki, D., Ławicki, Ł. & Girjatowicz, J. Ducks change wintering patterns due to changing climate in the important wintering waters of the Odra River

- (Helsinki Commission, Helsinki, Finland, 2013).
- 25 Luigujõe, L., Kuresoo, A. & Rattiste, K. Population status of the Bewick's Swan (*Cygnus columbianus bewickii*) and protection proposals in Estonia. *Hirundo* 26, 53-79 (2013).
- 26 Žydelis, R. et al. Bycatch in gillnet fisheries – An overlooked threat to waterbird populations. *Biol. Conserv.* 142, 1269-1281, <https://doi.org/10.1016/j.biocon.2009.02.025> (2009).
- 27 Sonntag, N., Schwemmer, H., Fock, H. O., Bellebaum, J. & Garthe, S. Seabirds, set-nets, and conservation management: assessment of conflict potential and vulnerability of birds to bycatch in gillnets. *ICES J. Mar. Sci.* 69, 578-589, <https://doi.org/10.1093/icesjms/fss030> (2012).
- 28 Schwemmer, P., Mendel, B., Sonntag, N., Dierschke, V. & Garthe, S. Effects of ship traffic on seabirds in offshore waters: implications for marine conservation and spatial planning. *Ecol. Appl.* 21, 1851-1860, <https://doi.org/10.1890/10-0615.1> (2011).
- 29 Fließbach, K. L. et al. A Ship Traffic Disturbance Vulnerability Index for Northwest European Seabirds as a Tool for Marine Spatial Planning. *Front. Mar. Sci.* 6, <https://doi.org/10.3389/fmars.2019.00192> (2019).
- 30 Dierschke, V., Furness, R. W. & Garthe, S. Seabirds and offshore wind farms in European waters: Avoidance and attraction. *Biol. Conserv.* 202, 59-68, <https://doi.org/10.1016/j.biocon.2016.08.016> (2016).
- 31 Fox, A. D. & Petersen, I. K. Offshore wind farms and their effects on birds. *Dan. ornithol. foren. tidsskr.* 113, 86-101 (2019).
- 32 Hansson, S. & Rudstam, L. G. Eutrophication and Baltic fish communities. *Ambio* 19, 123-125 (1990).
- 33 Rönkä, M. T. H., Saari, C. L. V., Lehikoinen, E. A., Suomela, J. & Häkkinen, K. Environmental changes and population trends of breeding waterfowl in northern Baltic Sea. *Annal. Zool.* 42, 587-602 (2005).
- 34 Pavón-Jordán, D. et al. Positive impacts of important bird and biodiversity areas on wintering waterbirds under changing temperatures throughout Europe and North Africa. *Biol. Conserv.* 246, 108549, <https://doi.org/10.1016/j.biocon.2020.108549> (2020).

Marine mammals

- 1 Halkka, A. Changing climate and the Baltic region biota Doctoral dissertation thesis, University of Helsinki, (2020).
- 2 Jevrejeva, S. et al. Baltic Sea ice seasons in the twentieth century. *Clim. Res.* 25, 217-227, <https://doi.org/10.3354/cr025217> (2004).
- 3 Vihma, T. & Haapala, J. Geophysics of sea ice in the Baltic Sea: A review. *Prog. Oceanogr.* 80, 129-148, <https://doi.org/10.1016/j.pocean.2009.02.002> (2009).
- 4 Luomaranta, A. et al. Multimodel estimates of the changes in the Baltic Sea ice cover during the present century. *Tellus A: Dynamic Meteorology and Oceanography* 66, 22617, <https://doi.org/10.3402/tellusa.v66.22617> (2014).
- 5 Merkouriadi, I. & Leppäranta, M. Long-term analysis of hydrography and sea-ice data in Tvärminne, Gulf of Finland, Baltic Sea. *Clim. Change* 124, 849-859, <https://doi.org/10.1007/s10584-014-1130-3> (2014).
- 6 Haapala, J. J., Ronkainen, I., Schmelzer, N. & Sztobryn, M. in *Second Assessment of Climate Change for the Baltic Sea Basin* (ed BACC II Author Team) 145-153 (Springer International

- Publishing, 2015).
- 7 Laakso, L. et al. 100 years of atmospheric and marine observations at the Finnish Utö Island in the Baltic Sea. *Ocean Sci.* 14, 617-632, <https://doi.org/10.5194/os-14-617-2018> (2018).
- 8 von Schuckmann, K. et al. Copernicus Marine Service Ocean State Report, Issue 4. *J. Oper. Oceanogr.* 13, S1-S172, <https://doi.org/10.1080/1755876X.2020.1785097> (2020).
- 9 Meier, H. E. M. Baltic Sea climate in the late twenty-first century: a dynamical downscaling approach using two global models and two emission scenarios. *Clim. Dyn.* 27, 39-68, <https://doi.org/10.1007/s00382-006-0124-x> (2006).
- 10 Meier, H. E. M., Döscher, R. & Halkka, A. Simulated distributions of Baltic Sea-ice in warming climate and consequences for the winter habitat of the Baltic ringed seal. *Ambio* 33, 249-256, <https://doi.org/10.1579/0044-7447-33.4.249> (2004).
- 11 Meier, H. E. M. in *Second Assessment of Climate Change for the Baltic Sea Basin* (ed BACC II Author Team) 243-252 (Springer International Publishing, 2015).
- 12 Grinstead, A. in *Second Assessment of Climate Change for the Baltic Sea Basin Regional Climate Studies* (ed BACC II Author Team) 253-263 (Springer International Publishing, 2015).
- 13 IPCC. IPCC Special Report on the Ocean and Cryosphere in a Changing Climate. (2019).
- 14 Wilson, S. C. et al. Assessment of impacts and potential mitigation for icebreaking vessels transiting pupping areas of an ice-breeding seal. *Biol. Conserv.* 214, 213-222, <https://doi.org/10.1016/j.biocon.2017.05.028> (2017).
- 15 Wilson, S. C., Crawford, I., Trukhanova, I., Dmitrieva, L. & Goodman, S. J. Estimating risk to ice-breeding pinnipeds from shipping in Arctic and sub-Arctic seas. *Mar. Policy* 111, 103694, <https://doi.org/10.1016/j.marpol.2019.103694> (2020).
- 16 Nabe-Nielsen, J., Sibly, R. M., Tougaard, J., Teilmann, J. & Sveegaard, S. Effects of noise and by-catch on a Danish harbour porpoise population. *Ecol. Model.* 272, 242-251, <https://doi.org/10.1016/j.ecolmodel.2013.09.025> (2014).
- 17 Vanhatalo, J., Vetemaa, M., Herrero, A., Aho, T. & Tiilikainen, R. By-Catch of Grey Seals (*Halichoerus grypus*) in Baltic Fisheries—A Bayesian Analysis of Interview Survey. *PLoS ONE* 9, e113836, <https://doi.org/10.1371/journal.pone.0113836> (2014).
- 18 Authier, M. et al. Workshop on fisheries Emergency Measures to minimize BYCatch of short-beaked common dolphins in the Bay of Biscay and harbour porpoise in the Baltic Sea (WKEM-BYC). 354 (2020).
- 19 Andersen, S. M., Teilmann, J., Dietz, R., Schmidt, N. M. & Miller, L. A. Behavioural responses of harbour seals to human-induced disturbances. *Aquat. Conserv.* 22, 113-121, <https://doi.org/10.1002/aqc.1244> (2012).
- 20 Paterson, W. D. et al. Post disturbance haulout behaviour of harbour seals. *Aquatic Conservation* 29, 144-156 (2019).
- 21 Härkönen, T. et al. The 1988 and 2002 phocine distemper virus epidemics in European harbour seals. *Dis. Aquat. Organ.* 68, 115-130, <https://doi.org/10.3354/dao068115> (2006).
- 22 DeMaster, D. P., Fowler, C. W., Perry, S. L. & Richlen, M. F. Predation and Competition: The Impact of Fisheries on Marine-Mammal Populations Over the next one Hundred Years. *J. Mammal.* 82, 641-651, [https://doi.org/10.1644/1545-1542\(2001\)082<0641:PACTIO>2.0.CO;2](https://doi.org/10.1644/1545-1542(2001)082<0641:PACTIO>2.0.CO;2) (2001).
- 23 O'Shea, T. J. & Odell, D. K. Large-Scale Marine Ecosystem Change and the Conservation of Marine Mammals. *J. Mammal.* 89, 529-533, <https://doi.org/10.1644/07-MAMM-S-416R.1>



- (2008).
- 24 Sonne, C. et al. Health effects from contaminant exposure in Baltic Sea birds and marine mammals: A review. *Environ. Int.* 139, 105725, <https://doi.org/10.1016/j.envint.2020.105725> (2020).
- 25 Murphy, S. et al. Reproductive Failure in UK Harbour Porpoises *Phocoena phocoena*: Legacy of Pollutant Exposure? *PLoS ONE* 10, e0131085, <https://doi.org/10.1371/journal.pone.0131085> (2015).
- 26 Southall, B. L. et al. Overview. *Aquat. Mamm.* 33, 411 (2007).
- 27 Jüssi, M., Härkönen, T., Helle, E. & Jüssi, I. Decreasing Ice Coverage Will Reduce the Breeding Success of Baltic Grey Seal (Halichoerus grypus) Females. *Ambio* 37, 80-85, 86 (2008).
- 28 Hazen, E. L. et al. Marine top predators as climate and ecosystem sentinels. *Front. Ecol. Environ.* 17, 565-574, <https://doi.org/10.1002/fee.2125> (2019).

Non-indigenous species

- 1 Murphy, K. R., Ritz, D. & Hewitt, C. L. Heterogeneous zooplankton distribution in a ship's ballast tanks. *J. Plankton Res.* 24, 729-734, <https://doi.org/10.1093/plankt/24.7.729> (2002).
- 2 Gollasch, S. The Importance of Ship Hull Fouling as a Vector of Species Introductions into the North Sea. *Biofouling* 18, 105-121, <https://doi.org/10.1080/08927010290011361> (2002).
- 3 Gollasch, S., Lenz, J., Dammer, M. & Andres, H.-G. Survival of tropical ballast water organisms during a cruise from the Indian Ocean to the North Sea. *J. Plankton Res.* 22, 923-937, <https://doi.org/10.1093/plankt/22.5.923> (2000).
- 4 Olenin, S., Gollasch, S., Jonušas, S. & Rimkutė, I. En-Route Investigations of Plankton in Ballast Water on a Ship's Voyage from the Baltic Sea to the Open Atlantic Coast of Europe. *Int. Rev. Hydrobiol.* 85, 577-596, [https://doi.org/10.1002/1522-2632\(200011\)85:5/6<577::AID-IROH577>3.0.CO;2-C](https://doi.org/10.1002/1522-2632(200011)85:5/6<577::AID-IROH577>3.0.CO;2-C) (2000).
- 5 Zaiko, A., Lehtiniemi, M., Narščius, A. & Olenin, S. Assessment of bioinvasion impacts on a regional scale: a comparative approach. *Biol. Invasions* 13, 1739-1765, <https://doi.org/10.1007/s10530-010-9928-z> (2011).
- 6 Nehring, S. Four arguments why so many alien species settle into estuaries, with special reference to the German river Elbe. *Helgoland Marine Research* 60, 127, <https://doi.org/10.1007/s10152-006-0031-x> (2006).
- 7 Reise, K., Olenin, S. & Thieltges, D. W. Are aliens threatening European aquatic coastal ecosystems? *Helgoland Marine Research* 60, 77, <https://doi.org/10.1007/s10152-006-0024-9> (2006).
- 8 Olenin, S. & Leppäkoski, E. Non-native animals in the Baltic Sea: alteration of benthic habitats in coastal inlets and lagoons. *Hydrobiologia* 393, 233-243, <https://doi.org/10.1023/A:1003511003766> (1999).
- 9 HELCOM. BALSAM PROJECT – WORK PACKAGE 4: Testing Monitoring Methods for Non-Indigenous Species in Ports. 57pp (Helsinki, Finland, 2015).
- 10 Normant, M. et al. in BALSAM PROJECT – WORK PACKAGE 4: Testing Monitoring Methods for Non-Indigenous Species in Ports (eds M. Ruiz, J. Karhu, & H. Backer) 31-55 (Baltic Marine Environment Protection Commission, 2015).
- 11 Kukliński, P. & Witalis, B. Wpływ wód balastowych na bioróżnorodność organizmów poroślowych portu Gdynia. (2011).

- 12 Minchin, J. D. et al. Rapid expansion and facilitating factors of the Ponto-Caspian invader *Dikerogammarus villosus* within the eastern Baltic Sea. *Aquat. Invasions* 14, 165-181, <https://doi.org/10.3391/ai.2019.14.2.02> (2019).
- 13 AquaNIS. Information system on aquatic non-indigenous and cryptogenic species, <http://www.corpi.ku.lt/databases/index.php/aquanis/>, (last access: 09.02.2021).
- 14 Ojaveer, H. et al. Dynamics of biological invasions and pathways over time: a case study of a temperate coastal sea. *Biol. Invasions* 19, 799-813, <https://doi.org/10.1007/s10530-016-1316-x> (2017).
- 15 Stachowicz, J. J., Terwin, J. R., Whitlatch, R. B. & Osman, R. W. Linking climate change and biological invasions: Ocean warming facilitates nonindigenous species invasions. *Proceedings of the National Academy of Sciences* 99, 15497, <https://doi.org/10.1073/pnas.242437499> (2002).
- 16 Occhipinti-Ambrogi, A. Global change and marine communities: Alien species and climate change. *Mar. Pollut. Bull.* 55, 342-352, <https://doi.org/10.1016/j.marpolbul.2006.11.014> (2007).
- 17 Hellmann, J. J., Byers, J. E., Bierwagen, B. G. & Dukes, J. S. Five Potential Consequences of Climate Change for Invasive Species. *Conserv. Biol.* 22, 534-543, <https://doi.org/10.1111/j.1523-1739.2008.00951.x> (2008).
- 18 Olyarnik, S. V. et al. in *Biological Invasions in Marine Ecosystems: Ecological, Management, and Geographic Perspectives* (eds Gil Rilov & Jeffrey A. Crooks) 215-238 (Springer Berlin Heidelberg, 2009).
- 19 Pyke, C. R. et al. Current Practices and Future Opportunities for Policy on Climate Change and Invasive Species. *Conserv. Biol.* 22, 585-592, <https://doi.org/10.1111/j.1523-1739.2008.00956.x> (2008).
- 20 Rahel, F. J. & Olden, J. D. Assessing the Effects of Climate Change on Aquatic Invasive Species. *Conserv. Biol.* 22, 521-533, <https://doi.org/10.1111/j.1523-1739.2008.00950.x> (2008).
- 21 Ng, C. A. & Gray, K. A. Forecasting the effects of global change scenarios on bioaccumulation patterns in great lakes species. *Global Change Biol.* 17, 720-733, <https://doi.org/10.1111/j.1365-2486.2010.02299.x> (2011).
- 22 Holopainen, R. et al. Impacts of changing climate on the non-indigenous invertebrates in the northern Baltic Sea by end of the twenty-first century. *Biol. Invasions* 18, 3015-3032, <https://doi.org/10.1007/s10530-016-1197-z> (2016).
- 23 Kornobis, S. Ecology of *Dreissena polymorpha* (Pal.) (*Dreissena: Bivalvia*) in lakes receiving heated water discharges. *Polskie Archiwum Hydrobiologii* 24, 531-545 (1977).
- 24 Stanczykowska, A. & Stoczkowski, R. Are the changes in *Dreissena polymorpha* [Pall.] distribution in the Great Masurian Lakes related to trophic state? *Polskie Archiwum Hydrobiologii* 44, 417-429 (1997).
- 25 Borchertding, J. The annual reproductive cycle of the freshwater mussel *Dreissena polymorpha* Pallas in lakes. *Oecologia* 87, 208-218, <https://doi.org/10.1007/BF00325258> (1991).
- 26 Sprung, M. Costs of reproduction : a study on metabolic requirements of the gonads and fecundity of the bivalve *Dreissena polymorpha*. *Malacologia* 33, 63-70 (1991).
- 27 Sprung, M. Ecological requirements of developing *Dreissena polymorpha* eggs. *Arch. Hydrobiol., Suppl.bd., Monogr. Beitr* 79, 69-86 (1987).
- 28 Claudi, R. & Mackie, G. L. (Lewis publishers, Boca Raton, Florida, 1994).

- 29 Fanslow, D. L., Nalepa, T. F. & Lang, G. A. Filtration Rates of the Zebra Mussel (*Dreissena polymorpha*) on Natural Seston from Saginaw Bay, Lake Huron. *J. Great Lakes Res.* 21, 489-500, [https://doi.org/10.1016/S0380-1330\(95\)71061-9](https://doi.org/10.1016/S0380-1330(95)71061-9) (1995).
- 30 Aldridge, D. W., Payne, B. S. & Miller, A. C. Oxygen consumption, nitrogenous excretion, and filtration rates of *Dreissena polymorpha* at acclimation temperatures between 20 and 32 °C. *Can. J. Fish. Aquat. Sci.* 52, 1761-1767, <https://doi.org/10.1139/f95-768> (1995).
- 31 HELCOM. HELCOM Baltic Sea Environment Fact Sheets, <http://www.HELCOM.fi/baltic-sea-trends/environment-fact-sheets/>, (last access: 19.07.2021).
- 32 Wimbush, J., Sloane, W., Farrell, J. & Nierzwicki-Bauer, S. A. in Conference proceeding: 13th international Conference on aquatic invasive species.
- 33 Lee, V. A. & Johnson, T. B. Development of a Bioenergetics Model for the Round Goby (*Neogobius melanostomus*). *J. Great Lakes Res.* 31, 125-134, [https://doi.org/10.1016/S0380-1330\(05\)70244-6](https://doi.org/10.1016/S0380-1330(05)70244-6) (2005).
- 34 Almqvist, G. Round goby *Neogobius melanostomus* in the Baltic Sea – Invasion Biology in practice, Systemekologiska institutionen, (2008).
- 35 Tomczak, M. T. & Sapota, M. R. The fecundity and gonad development cycle of the round goby (*Neogobius melanostomus* Pallas 1811) from the Gulf of Gdańsk. *Oceanol. Hydrobiol. Stud.* 35, 353-367 (2005).
- 36 Leppäkoski, E. & Olenin, S. Non-native Species and Rates of Spread: Lessons from the Brackish Baltic Sea. *Biol. Invasions* 2, 151-163, <https://doi.org/10.1023/A:1010052809567> (2000).
- 37 Orlova, M. I. et al. Effects of nonindigenous species on diversity and community functioning in the eastern Gulf of Finland (Baltic Sea). *Helgoland Marine Research* 60, 98-105, <https://doi.org/10.1007/s10152-006-0026-7> (2006).
- 38 Arend, K. K. et al. Seasonal and interannual effects of hypoxia on fish habitat quality in central Lake Erie. *Freshw. Biol.* 56, 366-383, <https://doi.org/10.1111/j.1365-2427.2010.02504.x> (2011).
- 39 Maximov, A. A. Large-scale invasion of *Marenzelleria* spp. (Polychaeta; Spionidae) in the eastern Gulf of Finland, Baltic Sea. *Russ. J. Biol. Invasions* 2, 11-19, <https://doi.org/10.1134/S2075111711010036> (2011).
- 40 Josefsson, S., Leonardsson, K., Gunnarsson, J. S. & Wiberg, K. Bioturbation-Driven Release of Buried PCBs and PBDEs from Different Depths in Contaminated Sediments. *Environ. Sci. Technol.* 44, 7456-7464, <https://doi.org/10.1021/es100615g> (2010).
- 41 Witte, S., Buschbaum, C., van Beusekom, J. E. E. & Reise, K. Does climatic warming explain why an introduced barnacle finally takes over after a lag of more than 50 years? *Biol. Invasions* 12, 3579-3589, <https://doi.org/10.1007/s10530-010-9752-5> (2010).
- 42 Pederson, J. et al. in ICES status report on climate change in the North Atlantic (eds P. C. Reid & L. Valdés) 262pp (ICES Cooperative Research Report No. 310, 2011).

Ecosystem function

- 1 Kahru, M., Elmgren, R. & Savchuk, O. P. Changing seasonality of the Baltic Sea. *Biogeosciences* 13, 1009-1018, <https://doi.org/10.5194/bg-13-1009-2016> (2016).

- 2 Hjerne, O., Hajdu, S., Larsson, U., Downing, A. S. & Winder, M. Climate Driven Changes in Timing, Composition and Magnitude of the Baltic Sea Phytoplankton Spring Bloom. *Front. Mar. Sci.* 6, <https://doi.org/10.3389/fmars.2019.00482> (2019).
- 3 Kahru, M., Elmgren, R., Kaiser, J., Wasmund, N. & Savchuk, O. Cyanobacterial blooms in the Baltic Sea: Correlations with environmental factors. *Harmful Algae* 92, 101739, <https://doi.org/10.1016/j.hal.2019.101739> (2020).
- 4 Kahru, M. & Elmgren, R. Multidecadal time series of satellite-detected accumulations of cyanobacteria in the Baltic Sea. *Biogeosciences* 11, 3619-3633, <https://doi.org/10.5194/bg-11-3619-2014> (2014).
- 5 Casini, M. et al. Trophic cascades promote threshold-like shifts in pelagic marine ecosystems. *Proceedings of the National Academy of Sciences* 106, 197, <https://doi.org/10.1073/pnas.0806649105> (2009).
- 6 Möllmann, C. et al. Reorganization of a large marine ecosystem due to atmospheric and anthropogenic pressure: a discontinuous regime shift in the Central Baltic Sea. *Global Change Biol.* 15, 1377-1393, <https://doi.org/10.1111/j.1365-2486.2008.01814.x> (2009).
- 7 Eriksson, B. K. et al. Declines in predatory fish promote bloom-forming macroalgae. *Ecol. Appl.* 19, 1975-1988, <https://doi.org/10.1890/08-0964.1> (2009).
- 8 Lindegren, M., Blenckner, T. & Stenseth, N. C. Nutrient reduction and climate change cause a potential shift from pelagic to benthic pathways in a eutrophic marine ecosystem. *Global Change Biol.* 18, 3491-3503, <https://doi.org/10.1111/j.1365-2486.2012.02799.x> (2012).
- 9 Suikkanen, S. et al. Climate Change and Eutrophication Induced Shifts in Northern Summer Plankton Communities. *PLoS ONE* 8, e66475, <https://doi.org/10.1371/journal.pone.0066475> (2013).
- 10 Carstensen, J., Andersen, J. H., Gustafsson, B. G. & Conley, D. J. Deoxygenation of the Baltic Sea during the last century. *P. Natl. Acad. Sci. USA* 111, 5628-5633, <https://doi.org/10.1073/pnas.1323156111> (2014).
- 11 Eriksson, B. K. et al. Effects of Altered Offshore Food Webs on Coastal Ecosystems Emphasize the Need for Cross-Ecosystem Management. *Ambio* 40, 786, <https://doi.org/10.1007/s13280-011-0158-0> (2011).
- 12 Östman, Ö. et al. Top-down control as important as nutrient enrichment for eutrophication effects in North Atlantic coastal ecosystems. *J. Appl. Ecol.* 53, 1138-1147, <https://doi.org/10.1111/1365-2664.12654> (2016).
- 13 Pekcan-Hekim, Z. et al. The role of climate and fisheries on the temporal changes in the Bothnian Bay foodweb. *ICES J. Mar. Sci.* 73, 1739-1749, <https://doi.org/10.1093/icesjms/fsw032> (2016).
- 14 Karlson, A. M. L. et al. Nitrogen fixation by cyanobacteria stimulates production in Baltic food webs. *Ambio* 44, 413-426, <https://doi.org/10.1007/s13280-015-0660-x> (2015).
- 15 Saraiva, S. et al. Baltic Sea ecosystem response to various nutrient load scenarios in present and future climates. *Clim. Dyn.* 52, 3369-3387, <https://doi.org/10.1007/s00382-018-4330-0> (2019).
- 16 Svensson, F. et al. In situ warming strengthens trophic cascades in a coastal food web. *Oikos* 126, 1150-1161, <https://doi.org/10.1111/oik.03773> (2017).
- 17 Pecuchet, L., Törnroos, A. & Lindegren, M. Patterns and drivers of fish community assembly in a large marine ecosystem.





- Mar. Ecol. Prog. Ser. 546, 239-248, <https://doi.org/10.3354/meps11613> (2016).
- 18 Niiranen, S. et al. Combined effects of global climate change and regional ecosystem drivers on an exploited marine food web. *Global Change Biol.* 19, 3327-3342, <https://doi.org/10.1111/gcb.12309> (2013).
- 19 Bauer, B. et al. Food web and fisheries in the future Baltic Sea. *Ambio*, <https://doi.org/10.1007/s13280-019-01229-3> (2019).
- 20 Olsson, J., Bergström, L. & Gårdmark, A. Abiotic drivers of coastal fish community change during four decades in the Baltic Sea. *ICES J. Mar. Sci.* 69, 961-970, <https://doi.org/10.1093/icesjms/fss072> (2012).
- 21 Helenius, L. K., Leskinen, E., Lehtonen, H. & Nurminen, L. Spatial patterns of littoral zooplankton assemblages along a salinity gradient in a brackish sea: A functional diversity perspective. *Estuar. Coast. Shelf Sci.* 198, 400-412, <https://doi.org/10.1016/j.ecss.2016.08.031> (2017).
- 22 Bartels, P., Ask, J., Andersson, A., Karlsson, J. & Giesler, R. Allochthonous Organic Matter Supports Benthic but Not Pelagic Food Webs in Shallow Coastal Ecosystems. *Ecosystems* 21, 1459-1470, <https://doi.org/10.1007/s10021-018-0233-5> (2018).
- 23 Sandberg, J. Cross-ecosystem analyses of pelagic food web structure and processes in the Baltic Sea. *Ecol. Model.* 201, 243-261, <https://doi.org/10.1016/j.ecolmodel.2006.09.023> (2007).
- 24 Eklöf, J. S. et al. Experimental climate change weakens the insurance effect of biodiversity. *Ecol. Lett.* 15, 864-872, <https://doi.org/10.1111/j.1461-0248.2012.01810.x> (2012).
- 25 Humborg, C. et al. High Emissions of Carbon Dioxide and Methane From the Coastal Baltic Sea at the End of a Summer Heat Wave. *Front. Mar. Sci.* 6, <https://doi.org/10.3389/fmars.2019.00493> (2019).
- 26 European Environment Agency. Maximum extent of ice cover in the Baltic Sea https://www.eea.europa.eu/data-and-maps/daviz/maximum-extent-of-ice-cover-3#tab-chart_1, (last access: 27.11.2020).
- concern. *Biol. Conserv.* 171, 177-185, <https://doi.org/10.1016/j.biocon.2014.01.033> (2014).
- 7 Clausen, K. K., Stjernholm, M. & Clausen, P. Grazing management can counteract the impacts of climate change-induced sea level rise on salt marsh-dependent waterbirds. *J. Appl. Ecol.* 50, 528-537, <https://doi.org/10.1111/1365-2664.12043> (2013).
- 8 Takolander, A., Leskinen, E. & Cabeza, M. Synergistic effects of extreme temperature and low salinity on foundational macroalga *Fucus vesiculosus* in the northern Baltic Sea. *J. Exp. Mar. Biol. Ecol.* 495, 110-118, <https://doi.org/10.1016/j.jembe.2017.07.001> (2017).
- 9 Westerborn, M., Mustonen, O., Jaatinen, K., Kilpi, M. & Norkko, A. Population Dynamics at the Range Margin: Implications of Climate Change on Sublittoral Blue Mussels (*Mytilus trossulus*). *Front. Mar. Sci.* 6, <https://doi.org/10.3389/fmars.2019.00292> (2019).
- 10 Dippner, J. W., Hänninen, J., Kuosa, H. & Vuorinen, I. The influence of climate variability on zooplankton abundance in the Northern Baltic Archipelago Sea (SW Finland). *ICES J. Mar. Sci.* 58, 569-578, <https://doi.org/10.1006/jmsc.2001.1048> (2001).
- 11 Kovtun, A. et al. Influence of abiotic environmental conditions on spatial distribution of charophytes in the coastal waters of West Estonian Archipelago, Baltic Sea. *J. Coastal Res.*, 412-416 (2011).
- 12 Ehlers, A., Worm, B. & Reusch, T. B. H. Importance of genetic diversity in eelgrass *Zostera marina* for its resilience to global warming. *Mar. Ecol. Prog. Ser.* 355, 1-7, <https://doi.org/10.3354/meps07369> (2008).
- 13 Jonsson, P. R. et al. High climate velocity and population fragmentation may constrain climate-driven range shift of the key habitat former *Fucus vesiculosus*. *Divers. Distrib.* 24, 892-905, <https://doi.org/10.1111/ddi.12733> (2018).
- 14 Kotta, J. et al. Integrating experimental and distribution data to predict future species patterns. *Sci. Rep.-UK* 9, 1821, <https://doi.org/10.1038/s41598-018-38416-3> (2019).
- 15 Jonsson, P. R., Moksnes, P.-O., Corell, H., Bonsdorff, E. & Nilsson Jacobi, M. Ecological coherence of Marine Protected Areas: New tools applied to the Baltic Sea network. *Aquat. Conserv.* 30, 743-760, <https://doi.org/10.1002/aqc.3286> (2020).
- 16 HELCOM. Ecological coherence assessment of the Marine Protected Area network in the Baltic. 74pp (Helsinki, Finland, 2016).
- 17 Khatiwala, S. et al. Global ocean storage of anthropogenic carbon. *Biogeosciences* 10, 2169-2191, <https://doi.org/10.5194/bg-10-2169-2013> (2013).

Marine protected areas

- 1 Laikre, L. et al. Lack of recognition of genetic biodiversity: International policy and its implementation in Baltic Sea marine protected areas. *Ambio* 45, 661-680, <https://doi.org/10.1007/s13280-016-0776-7> (2016).
- 2 Lewis, N. et al. Large-scale marine protected areas: guidelines for design and management. xxviii + 120pp (Best Practice Protected Area Guidelines Series, No. 26, IUCN, Gland, Switzerland, 2017).
- 3 Dureuil, M., Boerder, K., Burnett, K. A., Froese, R. & Worm, B. Elevated trawling inside protected areas undermines conservation outcomes in a global fishing hot spot. *Science* 362, 1403, <https://doi.org/10.1126/science.aau0561> (2018).
- 4 HELCOM. Implementation of the Baltic Sea Action Plan 2018 - Three years left to reach good environmental status. Background document to the 2018 HELCOM Ministerial Meeting. 91pp (Brussels, Belgium, 2018).
- 5 Reusch, T. B. H. et al. The Baltic Sea as a time machine for the future coastal ocean. *Sci. Adv.* 4, <https://doi.org/10.1126/sciadv.aar8195> (2018).
- 6 Clausen, K. K. & Clausen, P. Forecasting future drowning of coastal waterbird habitats reveals a major conservation

Nutrient concentrations and eutrophication

- 1 Schneider, B., Eilola, K., Lukkari, K., Müller-Karulis, B. & Neumann, T. in *Second Assessment of Climate Change for the Baltic Sea Basin* (ed BACC II Author Team) 337-361 (Springer International Publishing, 2015).
- 2 Gustafsson, E., Savchuk, O. P., Gustafsson, B. G. & Müller-Karulis, B. Key processes in the coupled carbon, nitrogen, and phosphorus cycling of the Baltic Sea. *Biogeochemistry* 134, 301-317, <https://doi.org/10.1007/s10533-017-0361-6> (2017).
- 3 Conley, D. J., Humborg, C., Rahm, L., Savchuk, O. P. & Wulff, F. Hypoxia in the Baltic Sea and basin-scale changes in phosphorus biogeochemistry. *Environ. Sci. Technol.* 36, 5315-5320,

- <https://doi.org/10.1021/es025763w> (2002).
- 4 Savchuk, O. P. Large-Scale Nutrient Dynamics in the Baltic Sea, 1970-2016. *Front. Mar. Sci.* 5, 95, <https://doi.org/10.3389/fmars.2018.00095> (2018).
- 5 HELCOM. HELCOM thematic assessment of eutrophication 2011-2016., 83pp (Helsinki, Finland, 2018).
- 6 Carstensen, J. & Conley, D. J. Baltic Sea Hypoxia Takes Many Shapes and Sizes. *Limnol. Oceanogr. Bull.* 28, 125-129, <https://doi.org/10.1002/lob.10350> (2019).
- 7 Carstensen, J., Henriksen, P. & Heiskanen, A.-S. Summer algal blooms in shallow estuaries: Definition, mechanisms, and link to eutrophication. *Limnol. Oceanogr.* 52, 370-384, <https://doi.org/10.4319/lo.2007.52.1.0370> (2007).
- 8 Wasmund, N. et al. Extension of the growing season of phytoplankton in the western Baltic Sea in response to climate change. *Mar. Ecol. Prog. Ser.* 622, 1-16, <https://doi.org/10.3354/meps12994> (2019).
- 9 Meier, H. E. M. et al. Assessment of Eutrophication Abatement Scenarios for the Baltic Sea by Multi-Model Ensemble Simulations. *Front. Mar. Sci.* 5, <https://doi.org/10.3389/fmars.2018.00440> (2018).
- 10 Meier, H. E. M. et al. Modeling the combined impact of changing climate and changing nutrient loads on the Baltic Sea environment in an ensemble of transient simulations for 1961-2099. *Clim. Dyn.* 39, 2421-2441, <https://doi.org/10.1007/s00382-012-1339-7> (2012).
- 11 Meier, H. E. M. et al. Impact of climate change on ecological quality indicators and biogeochemical fluxes in the Baltic Sea: a multi-model ensemble study. *Ambio* 41, 558-573, <https://doi.org/10.1007/s13280-012-0320-3> (2012).
- 12 Saraiva, S. et al. Uncertainties in Projections of the Baltic Sea Ecosystem Driven by an Ensemble of Global Climate Models. *Front. Earth Sci.* 6, <https://doi.org/10.3389/feart.2018.00244> (2019).
- 13 Neumann, T. et al. Extremes of temperature, oxygen and blooms in the Baltic Sea in a changing climate. *Ambio* 41, 574-585, <https://doi.org/10.1007/s13280-012-0321-2> (2012).
- 14 Saraiva, S. et al. Baltic Sea ecosystem response to various nutrient load scenarios in present and future climates. *Clim. Dyn.* 52, 3369-3387, <https://doi.org/10.1007/s00382-018-4330-0> (2019).
- 15 Meier, H. E. M. et al. Future projections of record-breaking sea surface temperature and cyanobacteria bloom events in the Baltic Sea. *Ambio* 48, 1362-1376, <https://doi.org/10.1007/s13280-019-01235-5> (2019).
- 16 Andersson, A. et al. Projected future climate change and Baltic Sea ecosystem management. *Ambio* 44, 345-356, <https://doi.org/10.1007/s13280-015-0654-8> (2015).
- 17 Meier, H. E. M. et al. Assessment of Uncertainties in Scenario Simulations of Biogeochemical Cycles in the Baltic Sea. *Front. Mar. Sci.* 6, <https://doi.org/10.3389/fmars.2019.00046> (2019).
- 18 Blomqvist, S., Gunnars, A. & Elmgren, R. Why the limiting nutrient differs between temperate coastal seas and freshwater lakes: A matter of salt. *Limnol. Oceanogr.* 49, 2236-2241, <https://doi.org/10.4319/lo.2004.49.6.2236> (2004).
- 19 Ehrnsten, E., Norkko, A., Müller-Karulis, B., Gustafsson, E. & Gustafsson, B. G. The meagre future of benthic fauna in a coastal sea—Benthic responses to recovery from eutrophication in a changing climate. *Global Change Biol.* 26, 2235-2250, <https://doi.org/10.1111/gcb.15014> (2020).
- 20 Carstensen, J. et al. Factors regulating the coastal nutrient

- filter in the Baltic Sea. *Ambio* 49, 1194-1210, <https://doi.org/10.1007/s13280-019-01282-y> (2020).
- 21 HELCOM. HELCOM Baltic Sea Action Plan. 101pp (Krakow, Poland, 2007).
- 22 European Commission. DIRECTIVE 2008/56/EC OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL of 17 June 2008, establishing a framework for community action in the field of marine environmental policy (Marine Strategy Framework Directive) (Text with EEA relevance). (2008).
- 23 European Commission. DIRECTIVE 2000/60/EC OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL of 23 October 2000 establishing a framework for Community action in the field of water policy. (2000).

Offshore wind farms

- 1 WindEurope. Offshore Wind in Europe - Key trends and statistics 2019. 40pp (Brussels, Belgium, 2020).
- 2 European Commission. Study on Baltic offshore wind energy cooperation under BEMIP - Final report. ENER/C1/2018-456, June 2019. 250pp (Directorate C - Renewables, Research and Innovation, Energy Efficiency, Brussels, Belgium, 2019).
- 3 Clark, S., Schroeder, F. & Baschek, B. The Influence of Large Offshore Wind Farms on the North Sea and Baltic Sea: A Comprehensive Literature Review. (Helmholtz-Zentrum Geesthacht, Zentrum für Material-und Küstenforschung, 2014).
- 4 Gutow, L. et al. in *Ecological Research at the Offshore Wind-farm alpha ventus: Challenges, Results and Perspectives* (eds Maritime Federal, Agency Hydrographic, Nature Conservation Federal Ministry for the Environment, & Safety Nuclear) 67-81 (Springer Fachmedien Wiesbaden, 2014).
- 5 Dierschke, V., Garthe, S. & Mendel, B. in *Offshore Wind Energy: Research on Environmental Impacts* (eds Julia Köller, Johann Köppel, & Wolfgang Peters) 121-143 (Springer Berlin Heidelberg, 2006).
- 6 Bergström, L. et al. Effects of offshore wind farms on marine wildlife—a generalized impact assessment. *Environ. Res. Lett.* 9, 034012, <https://doi.org/10.1088/1748-9326/9/3/034012> (2014).
- 7 Peschko, V., Mendel, B., Mercker, M., Dierschke, J. & Garthe, S. Northern gannets (*Morus bassanus*) are strongly affected by operating offshore wind farms during the breeding season. *J. Environ. Manag.* 279, 111509, <https://doi.org/10.1016/j.jenvman.2020.111509> (2021).
- 8 Peschko, V., Mercker, M. & Garthe, S. Telemetry reveals strong effects of offshore wind farms on behaviour and habitat use of common guillemots (*Uria aalge*) during the breeding season. *Mar. Biol.* 167, 118, <https://doi.org/10.1007/s00227-020-03735-5> (2020).
- 9 Peschko, V. et al. Effects of offshore windfarms on seabird abundance: Strong effects in spring and in the breeding season. *Mar. Environ. Res.* 162, 105157, <https://doi.org/10.1016/j.marenvres.2020.105157> (2020).
- 10 Rusu, E. An evaluation of the wind energy dynamics in the Baltic Sea, past and future projections. *Renew. Energy* 160, 350-362, <https://doi.org/10.1016/j.renene.2020.06.152> (2020).
- 11 European Commission. An EU Strategy to harness the potential of offshore renewable energy for a climate neutral future. (Brussels, Belgium, 2020).
- 12 European Commission. Guidance document on wind energy



- developments and EU nature legislation. 182pp (2020).
- 13 WindEurope. How offshore wind will help Europe go carbon-neutral 80pp (Brussels, Belgium, 2019).
- 14 Göke, C., Dahl, K. & Mohn, C. Maritime Spatial Planning supported by systematic site selection: Applying Marxan for offshore wind power in the western Baltic Sea. *PLoS ONE* 13, e0194362, <https://doi.org/10.1371/journal.pone.0194362> (2018).

Coastal protection

- 1 Harff, J. et al. in *Coastline Changes of the Baltic Sea from South to East: Past and Future Projection* (eds Jan Harff, Kazimierz Furmańczyk, & Hans von Storch) 15-35 (Springer International Publishing, 2017).
- 2 Liqueste, C., Zulian, G., Delgado, I., Stips, A. & Maes, J. Assessment of coastal protection as an ecosystem service in Europe. *Ecol. Indic.* 30, 205-217, <https://doi.org/10.1016/j.ecolind.2013.02.013> (2013).
- 3 Weisner, E. & Schernewski, G. Adaptation to climate change: A combined coastal protection and re-alignment scheme in a Baltic tourism region. *J. Coastal Res.*, 1963-1968, <https://doi.org/10.2112/si65-332.1> (2013).
- 4 Nordstrom, K. F. Living with shore protection structures: A review. *Estuar. Coast. Shelf Sci.* 150, 11-23, <https://doi.org/10.1016/j.ecss.2013.11.003> (2014).
- 5 Spalding, M. D. et al. The role of ecosystems in coastal protection: Adapting to climate change and coastal hazards. *Ocean Coast. Manage.* 90, 50-57, <https://doi.org/10.1016/j.ocecoaman.2013.09.007> (2014).
- 6 Schernewski, G., Schumacher, J., Weisner, E. & Donges, L. A combined coastal protection, realignment and wetland restoration scheme in the southern Baltic: planning process, public information and participation. *J. Coast. Conserv.* 22, 533-547, <https://doi.org/10.1007/s11852-017-0542-4> (2018).
- 7 Schernewski, G., Bartel, C., Kobarg, N. & Karnaускаite, D. Retrospective assessment of a managed coastal realignment and lagoon restoration measure: the Geltinger Birk, Germany. *J. Coast. Conserv.* 22, 157-167, <https://doi.org/10.1007/s11852-017-0496-6> (2018).
- 8 Schmidt, A., Striegnitz, M. & Kuhn, K. Integrating regional perceptions into climate change adaptation: a transdisciplinary case study from Germany's North Sea Coast. *Reg. Environ. Change* 14, 2105-2114, <https://doi.org/10.1007/s10113-012-0338-x> (2014).
- 9 Hoffmann, G. & Lampe, R. Sediment budget calculation to estimate Holocene coastal changes on the southwest Baltic Sea (Germany). *Mar. Geol.* 243, 143-156, <https://doi.org/10.1016/j.margeo.2007.04.014> (2007).
- 10 BACC II Author Team. Second Assessment of Climate Change for the Baltic Sea Basin. (Springer International Publishing, 2015).
- 11 Hopkins, T. S. et al. A Systems Approach Framework for the Transition to Sustainable Development: Potential Value Based on Coastal Experiments. *Ecol. Soc.* 17, <https://doi.org/10.5751/ES-05266-170339> (2012).
- 12 Baltranaitė, E., Povilanskas, R., Dučinskas, K., Ernšteins, R. & Tõnisson, H. Systems Approach to Eastern Baltic Coastal Zone Management. *Water* 12, <https://doi.org/10.3390/w12113102> (2020).

- 13 HELCOM. RECOMMENDATION 16/3 - HELSINKI COMMISSION. 2pp (1995).

Shipping

- 1 HELCOM. HELCOM Map and Data Service, <https://maps.helcom.fi/website/mapservice/>, (last access: 29.09.2020).
- 2 HELCOM. Shipping accidents in the Baltic Sea 2018. 33pp (2019).
- 3 Turner, D. R., Hassellöv, I.-M., Ytreberg, E. & Rutgersson, A. Shipping and the environment: Smokestack emissions, scrubbers and unregulated oceanic consequences. *Elementa-Sci. Anthropol.* 5, <https://doi.org/10.1525/elementa.167> (2017).
- 4 Haapala, J., Meier, H. E. M. & Rinne, J. Numerical investigations of future ice conditions in the Baltic Sea. *Ambio* 30, 237-244, <https://doi.org/10.1579/0044-7447-30.4.237> (2001).
- 5 Höglund, A., Pemberton, P., Hordoir, R. & Schimanke, S. Ice conditions for maritime traffic in the Baltic Sea in future climate. *Boreal Environ. Res.* 22, 245-265 (2017).
- 6 Rutgersson, A. et al. Natural Hazards and Extreme Events in the Baltic Sea region. *Earth Syst. Dynam. Discuss.*, 1-80, <https://doi.org/10.5194/esd-2021-13> (2021).
- 7 De Ceuster, G., van Herbruggen, B. & Logghe, S. TREMOVE - description of model and baseline version 2.41. Report for the European Commission. (Leuven, Belgium, 2006).
- 8 IMO. MEPC 72/17 - Report of the Marine Environment Protection Committee on its seventy-second session. 64 (International Maritime Organization, 2018).

Tourism

- 1 Jacobsen, P. B. State of the Tourism Industry in the Baltic Sea Region - 2018 Edition. 48pp (Baltic Sea Tourism Center (BSTC), Rostock, Germany, 2018).
- 2 Więckowski, M. & Cerić, D. Evolving tourism on the Baltic Sea coast: perspectives on change in the Polish maritime borderland. *Scand. J. Hosp. Tour.* 16, 98-111, <https://doi.org/10.1080/15022250.2016.1244598> (2016).
- 3 Port of Helsinki. Annual Report 2019, <https://vuosikertomus2019.portofhelsinki.fi/en/>, (last access: 30.11.2020).
- 4 Scott, D., Gössling, S. & Hall, C. M. International tourism and climate change. *WIREs Climate Change* 3, 213-232, <https://doi.org/10.1002/wcc.165> (2012).
- 5 Kahru, M., Elmgren, R. & Savchuk, O. P. Changing seasonality of the Baltic Sea. *Biogeosciences* 13, 1009-1018, <https://doi.org/10.5194/bg-13-1009-2016> (2016).
- 6 Ojaveer, H. et al. Twenty five years of invasion: management of the round goby *Neogobius melanostomus* in the Baltic Sea. *Manag. Biol. Invasions* 6, 329-339, <https://doi.org/10.3391/mbi.2015.6.4.02> (2015).
- 7 Hamilton, J. M., Maddison, D. J. & Tol, R. S. J. Climate change and international tourism: A simulation study. *Glob. Environ. Change* 15, 253-266, <https://doi.org/10.1016/j.gloenvcha.2004.12.009> (2005).
- 8 Tonazzini, D. et al. Blue tourism, Towards a Sustainable Coastal and Maritime Tourism in World Marine Regions. (eco-union Barcelona, 2019).

Fisheries

- 1 ICES. Baltic Sea Ecosystem - Fisheries Overview. Report of the ICES Advisory Committee ICES Advice 2019, section 4.2, 29pp, <https://doi.org/10.17895/ices.advice.5566> (2019).
- 2 HELCOM. Recreational fisheries in the Baltic Sea and availability of data. 9pp (Tallinn, Estonia, 2017).
- 3 Zeller, D. et al. The Baltic Sea: Estimates of total fisheries removals 1950–2007. *Fish. Res.* 108, 356-363, <https://doi.org/10.1016/j.fishres.2010.10.024> (2011).
- 4 Persson, L. in *Total marine fisheries extractions by country in the Baltic Sea: 1950-present* (eds P. Rossing, S. Booth, & D. Zeller) 225-263 (Fisheries Centre Research Reports 18 (1). Fisheries Centre, 2010).
- 5 Setälä, J. et al. Kalastuksen olosuhdekatsaus 2017. 46pp (Luonnonvarakeskus (Luke) 2018).
- 6 Pekcan-Hekim, Z. et al. Climate Warming and Pikeperch Year-Class Catches in the Baltic Sea. *Ambio* 40, 447-456, <https://doi.org/10.1007/s13280-011-0143-7> (2011).
- 7 Lappalainen, A. The effects of recent eutrophication on freshwater fish communities and fishery on the northern coast of the Gulf of Finland, Baltic Sea PhD thesis, University of Helsinki, Finland, (2002).
- 8 Bauer, B. et al. Food web and fisheries in the future Baltic Sea. *Ambio*, <https://doi.org/10.1007/s13280-019-01229-3> (2019).
- 9 Bauer, B. et al. Reducing eutrophication increases spatial extent of communities supporting commercial fisheries: a model case study. *ICES J. Mar. Sci.* 75, 1306-1317, <https://doi.org/10.1093/icesjms/fsy023> (2018).
- 10 Tunca, S., Lindegren, M., Ravn-Jonsen, L. & Lindroos, M. Co-operative Fisheries Outperform Non-cooperative Ones in the Baltic Sea Under Different Climate Scenarios. *Front. Mar. Sci.* 6, <https://doi.org/10.3389/fmars.2019.00622> (2019).
- 11 Niiranen, S. et al. Combined effects of global climate change and regional ecosystem drivers on an exploited marine food web. *Global Change Biol.* 19, 3327-3342, <https://doi.org/10.1111/gcb.12309> (2013).

Aquaculture

- 1 Asmala, E. & Saikku, L. Closing a Loop: Substance Flow Analysis of Nitrogen and Phosphorus in the Rainbow Trout Production and Domestic Consumption System in Finland. *Ambio* 39, 126-135, <https://doi.org/10.1007/s13280-010-0024-5> (2010).
- 2 Kankainen, M., Vielma, J., Koskela, J., Niukko, J. & Niskanen, L. Olosuhteiden vaikutus kirjoloehen kasvatuksen tehokkuuteen Suomen merialueilla. 34pp (Luonnonvarakeskus, Helsinki, Finland, 2020).
- 3 Huyben, D. et al. Dietary live yeast and increased water temperature influence the gut microbiota of rainbow trout. *J. Appl. Microbiol.* 124, 1377-1392, <https://doi.org/10.1111/jam.13738> (2018).
- 4 Kotta, J. et al. Cleaning up seas using blue growth initiatives: Mussel farming for eutrophication control in the Baltic Sea. *Sci. Total Environ.* 709, 136144, <https://doi.org/10.1016/j.scitotenv.2019.136144> (2020).
- 5 Visch, W. et al. Spatial differences in growth rate and nutrient mitigation of two co-cultivated, extractive species: The blue mussel (*Mytilus edulis*) and the kelp (*Saccharina latissima*). *Estuar. Coast. Shelf Sci.* 246, 107019, <https://doi.org/10.1016/j>

- ecss.2020.107019 (2020).
- 6 Benassai, G., Mariani, P., Stenberg, C. & Christoffersen, M. A Sustainability Index of potential co-location of offshore wind farms and open water aquaculture. *Ocean Coast. Manage.* 95, 213-218, <https://doi.org/10.1016/j.ocecoaman.2014.04.007> (2014).

Blue carbon storage capacity

- 1 Nellemann, C. et al. Blue carbon: the role of healthy oceans in binding carbon: a rapid response assessment. (UNEP/Earth-print, 2009).
- 2 Duarte, C. M., Losada, I. J., Hendriks, I. E., Mazarrasa, I. & Marbà, N. The role of coastal plant communities for climate change mitigation and adaptation. *Nat. Clim. Change* 3, 961-968, <https://doi.org/10.1038/nclimate1970> (2013).
- 3 Waycott, M. et al. Accelerating loss of seagrasses across the globe threatens coastal ecosystems. *Proceedings of the National Academy of Sciences* 106, 12377, <https://doi.org/10.1073/pnas.0905620106> (2009).
- 4 Hoegh-Guldberg, O. et al. in *The Ocean as a Solution to Climate Change: Five Opportunities for Action* 47-57 (World Resources Institute, 2019).
- 5 Pendleton, L. et al. Estimating Global “Blue Carbon” Emissions from Conversion and Degradation of Vegetated Coastal Ecosystems. *PLoS ONE* 7, e43542, <https://doi.org/10.1371/journal.pone.0043542> (2012).
- 6 Spivak, A. C., Sanderman, J., Bowen, J. L., Canuel, E. A. & Hopkinson, C. S. Global-change controls on soil-carbon accumulation and loss in coastal vegetated ecosystems. *Nat. Geosci.* 12, 685-692, <https://doi.org/10.1038/s41561-019-0435-2> (2019).
- 7 Gattuso, J.-P. et al. Ocean Solutions to Address Climate Change and Its Effects on Marine Ecosystems. *Front. Mar. Sci.* 5, <https://doi.org/10.3389/fmars.2018.00337> (2018).
- 8 United Nations Environment Programme. Out of the blue: The value of seagrasses to the environment and to people. (UNEP, Nairobi, Kenya, 2020).
- 9 Pessarrodona, A., Foggo, A. & Smale, D. A. Can ecosystem functioning be maintained despite climate-driven shifts in species composition? Insights from novel marine forests. *J. Ecol.* 107, 91-104, <https://doi.org/10.1111/1365-2745.13053> (2019).
- 10 Smale, D. A. et al. Marine heatwaves threaten global biodiversity and the provision of ecosystem services. *Nat. Clim. Change* 9, 306-312, <https://doi.org/10.1038/s41558-019-0412-1> (2019).
- 11 Krause-Jensen, D., Duarte, C. M., Sand-Jensen, K. & Carstensen, J. Century-long records reveal shifting challenges to seagrass recovery. *Global Change Biol.* 27, 563-575, <https://doi.org/10.1111/gcb.15440> (2021).
- 12 Takolander, A., Cabeza, M. & Leskinen, E. Climate change can cause complex responses in Baltic Sea macroalgae: A systematic review. *J. Sea Res.* 123, 16-29, <https://doi.org/10.1016/j.seares.2017.03.007> (2017).
- 13 Reusch, T. B. H. et al. The Baltic Sea as a time machine for the future coastal ocean. *Sci. Adv.* 4, <https://doi.org/10.1126/sciadv.aar8195> (2018).
- 14 Bobsien, I. C. et al. Modeling eelgrass spatial response to nutrient abatement measures in a changing climate. *Ambio* 50, 400-412, <https://doi.org/10.1007/s13280-020-01364-2> (2021).
- 15 Jonsson, P. R. et al. High climate velocity and population frag-





- mentation may constrain climate-driven range shift of the key habitat former *Fucus vesiculosus*. *Divers. Distrib.* 24, 892-905, <https://doi.org/10.1111/ddi.12733> (2018).
- 16 Clausen, K. K., Stjernholm, M. & Clausen, P. Grazing management can counteract the impacts of climate change-induced sea level rise on salt marsh-dependent waterbirds. *J. Appl. Ecol.* 50, 528-537, <https://doi.org/10.1111/1365-2664.12043> (2013).
 - 17 Duarte, C. M. et al. Rebuilding marine life. *Nature* 580, 39-51, <https://doi.org/10.1038/s41586-020-2146-7> (2020).
 - 18 Weisner, E. & Schernewski, G. Adaptation to climate change: A combined coastal protection and re-alignment scheme in a Baltic tourism region. *J. Coastal Res.*, 1963-1968, <https://doi.org/10.2112/si65-332.1> (2013).
 - 19 Leo, K. L., Gillies, C. L., Fitzsimons, J. A., Hale, L. Z. & Beck, M. W. Coastal habitat squeeze: A review of adaptation solutions for saltmarsh, mangrove and beach habitats. *Ocean Coast. Manage.* 175, 180-190, <https://doi.org/10.1016/j.ocecoaman.2019.03.019> (2019).
 - 20 Macreadie, P. I. et al. The future of Blue Carbon science. *Nat. Commun.* 10, 3998, <https://doi.org/10.1038/s41467-019-11693-w> (2019).
 - 21 Frigstad, H., Gundersen, H., Andersen, G. S., Borgersen, G., Kvile, K. Ø., Krause-Jensen, D., Boström, C., Bekkby, T., d'Auriac, M.A., Ruus, A., Thormar, J., Asdal, K., Hancke, K. Blue Carbon-climate adaptation, CO₂ uptake and sequestration of carbon in Nordic blue forests: Results from the Nordic Blue Carbon Project. *TemaNord* 2020:541. 139 pp. ISBN 978-92-893-6860-5 (PDF); ISBN 978-92-893-6861-2 (ONLINE). <http://dx.doi.org/10.6027/temanord2020-541> <https://www.norden.org/no/node/49407> (2021)
 - 22 The Blue Carbon Initiative. <https://www.thebluecarboninitiative.org/>, (last access: 22.07.2021).
 - 23 IUCN. Report: Blue Infrastructure Finance, where all win, <https://www.iucn.org/news/marine-and-polar/202003/report-blue-infrastructure-finance-where-all-win>, (last access: 22.07.2021).
 - 24 Nordic Co-operation. Nordic Ministerial Declaration on Oceans and Climate, <https://www.norden.org/en/declaration/nordic-ministerial-declaration-oceans-and-climate>, (last access: 22.07.2021).
 - 25 European Commission. Nature-based solutions, https://ec.europa.eu/info/research-and-innovation/research-area/environment/nature-based-solutions_en, (last access: 22.07.2021).
 - 26 FutureMARES. Climate Change and Future Marine Ecosystem Services and Biodiversity, <https://www.futuremares.eu/>, (last access: 22.07.2021).
 - Coastal and Marine Ecosystem Services. *Water* 11, 1518, <https://doi.org/10.3390/w11071518> (2019).
 - 5 Bauer, B. et al. Food web and fisheries in the future Baltic Sea. *Ambio*, <https://doi.org/10.1007/s13280-019-01229-3> (2019).
 - 6 Huyben, D. et al. Dietary live yeast and increased water temperature influence the gut microbiota of rainbow trout. *J. Appl. Microbiol.* 124, 1377-1392, <https://doi.org/10.1111/jam.13738> (2018).
 - 7 Visch, W. et al. Spatial differences in growth rate and nutrient mitigation of two co-cultivated, extractive species: The blue mussel (*Mytilus edulis*) and the kelp (*Saccharina latissima*). *Estuar. Coast. Shelf Sci.* 246, 107019, <https://doi.org/10.1016/j.ecss.2020.107019> (2020).
 - 8 Bobsien, I. C. et al. Modeling eelgrass spatial response to nutrient abatement measures in a changing climate. *Ambio* 50, 400-412, <https://doi.org/10.1007/s13280-020-01364-2> (2021).
 - 9 Jonsson, P. R. et al. High climate velocity and population fragmentation may constrain climate-driven range shift of the key habitat former *Fucus vesiculosus*. *Divers. Distrib.* 24, 892-905, <https://doi.org/10.1111/ddi.12733> (2018).
 - 10 Saraiva, S. et al. Baltic Sea ecosystem response to various nutrient load scenarios in present and future climates. *Clim. Dyn.* 52, 3369-3387, <https://doi.org/10.1007/s00382-018-4330-0> (2019).
 - 11 Svensson, F. et al. In situ warming strengthens trophic cascades in a coastal food web. *Oikos* 126, 1150-1161, <https://doi.org/10.1111/oik.03773> (2017).
 - 12 Olsson, J., Bergström, L. & Gårdmark, A. Abiotic drivers of coastal fish community change during four decades in the Baltic Sea. *ICES J. Mar. Sci.* 69, 961-970, <https://doi.org/10.1093/icesjms/fss072> (2012).
 - 13 Ivanauskas, E. & Razinkovas-Baziukas, A. Mapping and assessing commercial fisheries services in the Lithuanian part of the Curonian Lagoon. Submitted to *Acta Ichthyologica Et Piscatoria* (2021).
 - 14 Bartoli, M. et al. Drivers of Cyanobacterial Blooms in a Hypertrophic Lagoon. *Front. Mar. Sci.* 5, <https://doi.org/10.3389/fmars.2018.00434> (2018).
 - 15 Dam Lam, R., Gasparatos, A., Chakraborty, S., Rivera, H. & Stanley, T. Multiple values and knowledge integration in indigenous coastal and marine social-ecological systems research: A systematic review. *Ecosyst. Serv.* 37, 100910, <https://doi.org/10.1016/j.ecoser.2019.100910> (2019).

Marine and coastal ecosystem services

- 1 Culhane, F. E., Frid, C. L. J., Royo Gelabert, E., White, L. & Robinson, L. A. Linking marine ecosystems with the services they supply: what are the relevant service providing units? *Ecol. Appl.* 28, 1740-1751, <https://doi.org/10.1002/eap.1779> (2018).
- 2 Filies, C. & Schumacher, S. in *Climate Change Adaptation in Practice* 225-238 (2013).
- 3 Townsend, M. et al. The Challenge of Implementing the Marine Ecosystem Service Concept. *Front. Mar. Sci.* 5, <https://doi.org/10.3389/fmars.2018.00359> (2018).
- 4 Milon, J. W. & Alvarez, S. The Elusive Quest for Valuation of

