


Total phosphorus


Key Message

For total phosphorus, 12 open sea assessment units were evaluated for the period 2011-2015, of which good status has been achieved only in the Great Belt (total phosphorus concentration below defined threshold value). In Kattegat, Gdansk Bay and Bothnian Bay the concentrations were only slightly above the threshold value.

In the majority of coastal water assessment units, the threshold values set for total phosphorus were failed.



Key message figure 1: Status assessment results based on evaluation of the indicator 'Total phosphorus' (annual data). The assessment is carried out using Scale 4 HELCOM assessment units (defined in the [HELCOM Monitoring and Assessment Strategy Annex 4](#)). Please note that for some open sea areas threshold values still are under discussion, and that only those coastal areas are shown in which the assessment is based on annual data sets (same as in the open sea assessment).



Key message figure 2: Status assessment results based on evaluation of the indicator ‘Total phosphorus’ for coastal areas in which the summer concentrations (June-September) were used for assessment. The assessment is carried out using Scale 4 HELCOM assessment units (defined in the [HELCOM Monitoring and Assessment Strategy Annex 4](#)).

The indicator is applicable in all coastal and open sea areas. The indicator period and method of calculation varies between open sea and coastal areas, and thus the threshold value- or assessment concentrations are not directly comparable between the open sea and coast, nor between all coastal assessment units where nationally binding threshold values may have been set.

The indicator is applicable in the waters of all countries bordering the Baltic Sea, though not operational in all assessment units yet as for some open sea areas threshold values still need to be agreed upon.

Relevance of the core indicator

Eutrophication is caused by excessive inputs of nutrients (nitrogen and phosphorus) resulting from various human activities. High concentrations of nutrients and their ratios form the preconditions for huge algal blooms, reduced water clarity and increased oxygen consumption. Long-term nutrient data are key

parameters for quantifying the effects of anthropogenic activities and evaluating the success of measures undertaken.

Policy relevance of the core indicator

	BSAP Segment and Objectives	MSFD Descriptors and Criteria
Primary link	A Baltic Sea unaffected by eutrophication	D5 Human-induced eutrophication - D5C1 Nutrient concentrations are not at levels that indicate adverse eutrophication effects
Secondary link	A favourable conservation status of Baltic Sea biodiversity	D1 Biological diversity of species and habitats Theme: Pelagic habitats -D1C6 The condition of the habitat type, including its biotic and abiotic structure and its functions, is not adversely affected due to anthropogenic pressures. Theme: Benthic habitats -D6C5 The extent of adverse effects from anthropogenic pressures on the condition of the habitat type, including alteration to its biotic and abiotic structure and its functions, does not exceed a specified proportion of the natural extent of the benthic habitat type in the assessment area.
Other relevant legislation: Water Framework Directive, ecological status, QE4		

Cite this indicator

HELCOM (2017). Total phosphorus. HELCOM core indicator report. Online. [Date Viewed], [Web link].
ISSN 2343-2543


Download full indicator report

[HOLAS II component - Core indicator report – web-based version July 2017 \(pdf\)](#)


Results and Confidence

The assessment of total phosphorus in the open sea areas is made as the average of total phosphorus concentration in the upper (0-10 m) water layer throughout the year.

One assessment unit, namely the Great Belt, was found to achieve the threshold value during the assessment period 2011-2015. The remaining sub-basins were assessed as failing the threshold value or could not be assessed as threshold values still need to be agreed upon at HELCOM-wide level (Results figures 1 and 3, table 1). In Kattegat, Gdansk Bay and Bothnian Bay the concentrations were only slightly above the threshold value.




















Results figure 1. Detailed eutrophication status assessment with the Eutrophication ratio (ER) of total phosphorus (annual data) being split up into 5 classes to show a more differentiated picture than the 2-class division used in the key message figures. ER is calculated as the ratio of the average concentration during assessment period and the target (Fleming-Lehtinen et al. 2015). Please note that for some open sea areas threshold values still are under discussion, and that only those coastal areas are shown in which the assessment is based on annual data sets (same as in the open sea assessment). See Results figure 2 for other coastal areas.



Results figure 2. Detailed Eutrophication status assessment with the Eutrophication ratio (ER) of total phosphorus in coastal areas being split up into 5 classes to show a more differentiated picture than the 2-class division used in the key message figures. In coastal areas, the assessment is based on summer values (June-September). ER is calculated as the ratio of the average concentration during assessment period and the threshold value (Fleming-Lehtinen et al. 2015).

As becomes obvious from figure 1, some coastal areas in the southwestern Baltic are highly eutrophied. In the remaining coastal areas, seasonal instead of annual averages were used for assessment. Based on mean summer concentrations (June-September), some areas along the coasts of Sweden, Finland and Estonia are classified as achieving good status (figure 2), but highly eutrophied areas are found as well. If compared with the total nitrogen assessment, these coastal areas are somewhat stronger affected by phosphorus than by nitrogen.




Results figure 3. Average annual surface total phosphorus concentrations (black line; average for 2011-2015). The dashed red line displays the threshold value. For Kiel Bay, Bay of Mecklenburg, Arkona Basin, Bornholm Basin and Eastern Gotland Basin threshold values are still under discussion. Therefore, no dashed red line is shown in these cases, and these basins occur as “not assessed” in the maps above. The low concentration value in Gulf of Gdansk in 2011 are due to data handling problems. The issue is being investigated and is planned to be rectified for the next update of this indicator report.


Results table 1. Threshold values, present concentration (as average 2011-2015), eutrophication ratio (ER) and status of total nitrogen in the open sea basins. ER is a quantitative value for the level of eutrophication, calculated as the ratio between the threshold value and the present concentration – when $ER > 1$, threshold value has not been reached.

HELCOM ID	Assessment unit (open sea)	Threshold value ($\mu\text{mol l}^{-1}$)	Average 2011-2015 ($\mu\text{mol l}^{-1}$)	ER	Status (achieve/fail threshold value)
SEA-001	Kattegat	0.64	0.67	1.05	Fail
SEA-002	Great Belt	0.95	0.76	0.80	Achieve
SEA-003	The Sound	0.68	0.76	1.12	fail
SEA-004	Kiel Bay		0.72		Not assessed
SEA-005	Bay of Mecklenburg		0.73		Not assessed
SEA-006	Arkona Sea		0.84		Not assessed
SEA-007	Bornholm Basin		0.86		Not assessed
SEA-008	Gdansk Basin	0.6	0.67	1.12	Fail
SEA-009	Eastern Gotland Basin		0.73		Not assessed
SEA-010	Western Gotland Basin	0.45	0.78	1.72	Fail
SEA-011	Gulf of Riga	0.7	0.93	1.32	Fail
SEA-012	Northern Baltic Proper	0.38	0.70	1.85	Fail
SEA-013	Gulf of Finland	0.55	0.91	1.66	Fail
SEA-014	Åland Sea	0.28	0.50	1.80	Fail
SEA-015	Bothnian Sea	0.24	0.43	1.77	Fail
SEA-016	The Quark	0.24	0.32	1.32	Fail
SEA-017	Bothnian Bay	0.18	0.19	1.05	Fail

Additional information on temporal trends

Temporal trends provide additional information on the total nutrients in the Baltic Sea that supports the interpretation of the indicator results (Results figure 4). It should be noted that the temporal trends do not affect the indicator result, which is a status assessment where a concentration is compared to a threshold value. It should be further noted that the long-term temporal trends are not presented for the HELCOM assessment units, but are calculated for the BALTSEM basins.






Results 4. Long-term trends in annual surface total phosphorus concentrations (in $\mu\text{mol P l}^{-1}$), in the BALTSEM basins (HELCOM 2013) for 1970-2015. The data until 2012 is from TARGREV project. The spatial and seasonal patterns of historical are separated across the years, using a GLM-GAM model according to Carstensen et al. 2006. Data for 2013-2015 is based on data extraction from the HELCOM eutrophication assessment database and shows annual average concentrations for each sub-basin. Blue dashed lines indicate the 5-year moving average (starting from 1970) and error bars represent standard errors (SE).

Confidence of the indicator status evaluation

The data confidence of the total phosphorus indicator status evaluation for the open sea areas (Results figure 5) is high in all assessed sub-basins. It should be noted that the confidence is only based on data, not the target confidence since target confidence was not available for the indicator calculation.




Results figure 5. Indicator data confidence, determined combining information on data availability for the indicator when using observations from all months of the year. Low indicator confidence calls for increase in monitoring.

The indicator confidence was estimated only for the indicator data (ES-Score) due to absence of ET-Score, which describes the uncertainty of the threshold value setting procedure. The ES-Score is based on the number as well as spatial and temporal coverage of the observations for the assessment period 2011-2015. To estimate the overall indicator confidence, the ET-score should be defined and ET- and ES-Scores combined. See Andersen et al. 2010 and Fleming-Lehtinen et al. 2015 for further details.

As the indicator period and method of calculation varies between open sea and coastal areas, and thus the threshold or assessment concentrations are not directly comparable between the open sea and coast, nor between all coastal assessment units where nationally binding threshold values may have been set, only the confidence for the open sea areas are shown in Results figure 5.

Good Environmental Status

The threshold value of the ‘Total phosphorus’ core indicator is an assessment unit-specific concentration which is not to be exceeded in order for an assessment unit to be evaluated as having achieved the threshold value indicating good status (Good environmental status figure 1).



Good environmental status figure 1. Schematic representation of the threshold value for the core indicator ‘Total phosphorus’. Assessment unit-specific threshold values are used (see Good environmental status table 1).

Threshold values for the open-sea assessment units have been derived in HELCOM (Good environmental status table 1). For coastal assessment units, national boundaries used for estimating Good Environmental Status under WFD may be used.

Good environmental status table 1. Assessment unit-specific threshold values for total phosphorus.

HELCOM_ID	Assessment unit (open sea)	Threshold value [$\mu\text{mol l}^{-1}$]	Reference	Comments
SEA-001	Kattegat	0.64	HELCOM 38-2017	TARGREV value applied
SEA-002	Great Belt	0.95	HELCOM 38-2017	TARGREV value applied
SEA-003	The Sound	0.68	HELCOM 38-2017	TARGREV value applied
SEA-004	Kiel Bay			
SEA-005	Bay of Mecklenburg			
SEA-006	Arkona Basin			
SEA-007	Bornholm Basin			
SEA-008	Gdansk Basin	0.60	HELCOM 38-2017	
SEA-009	Eastern Gotland Basin			
SEA-010	Western Gotland Basin	0.45	HELCOM 38-2017	TARGREV value applied
SEA-011	Gulf of Riga	0.70	HELCOM 38-2017	New value (expert judgement)
SEA-012	Northern Baltic Proper	0.38	HELCOM 38-2017	TARGREV value applied
SEA-013	Gulf of Finland	0.55	HELCOM 38-2017	TARGREV value applied
SEA-014	Åland Sea	0.28	HELCOM 38-2017	TARGREV value applied
SEA-015	Bothnian Sea	0.24	HELCOM 38-2017	TARGREV value applied
SEA-016	The Quark	0.24	HELCOM 38-2017	TARGREV value applied
SEA-017	Bothnian Bay	0.18	HELCOM 38-2017	TARGREV value applied

Some of the open sea indicator threshold values were based on the results obtained in the TARGREV project (HELCOM 2013), taking also advantage of the work carried out during the EUTRO PRO process (HELCOM 2009) and national work for EU WFD implementation. The TARGREV values were derived as geometrical means, thus bearing close resemblance to median values (J. Carstensen, pers. comm.).

However, Total phosphorus (TP) was not simulated in the TARGREV modeling exercise, only upper limits of annual means of TP derived from estimates of the mean level during 1970-1975 are used as threshold values (see TARGREV report pages 84 and 134). These upper levels might already represent a eutrophied Baltic Sea in the early 1970s, and thus not be in agreement with the threshold value of the other eutrophication indicators with modelled threshold values (e.g. DIN, DIP) or threshold values based on extensive monitoring (e.g. Secchi depth). They are however expected to be in agreement with threshold values based on shorter term monitoring data (e.g. chlorophyll-*a*).

A new modeling approach has recently provided revised concentrations for German national threshold value of total nutrients in the Kiel Bay, Mecklenburg Bay, Arkona Basin and Bornholm Basin (Hirt et al. 2013; Schernewski et al. 2014; BLANO 2014) taking into account HELCOM, MSFD and WFD requirements for good status. The finally agreed BLANO threshold values represent median values and are included in the Federal Surface Water Ordinance (2016).

Break-point analysis was applied for setting Polish national threshold value in the Gdansk Basin. The results of these exercises were used as additional input in the threshold setting.

Assessment Protocol

The assessment of total phosphorus in open sea areas is made as the average of total phosphorus concentration in the upper (0-10 m) water layer throughout the year. In some coastal areas, annual averages are used as well (Key message figure 1), while in Sweden, Finland, Estonia, Lithuania and Poland the summer average is used to assess total phosphorus in coastal areas (Key message figure 2).

Assessment protocol table 1. Specifications of the indicator 'Total phosphorus'.

Indicator	Total phosphorus
Response to eutrophication	positive
Parameters	Total phosphorus concentration ($\mu\text{mol l}^{-1}$)
Data source	Monitoring data provided by the HELCOM Contracting Parties, and kept in the HELCOM COMBINE database, hosted by ICES (www.ices.dk)
Assessment period	2011-2015
Assessment season	Annual / Summer (June-September)
Depth	Surface = average in the 0-10 m layer
Removing outliers	No outliers removed
Removing close observations	No close observations removed, but Station 431 (Ven station) in The Sound has been included in the open sea area of The Sound, despite that it is located within the WFD baseline of the Ven island. However, due to the strong currents in The Sound this station is representative for the open waters in this assessment unit. Including this station will result in a much improved assessment for this assessment unit.
Indicator level (ES)	Average of annual/seasonal average values (mostly average = arithmetic mean, in some Contracting Parties the median is used instead to assess status versus threshold)
Indicator target (ET)	Agreed threshold values are mainly derived from TARGREV values as agreed by HOD 39-2012 with additions as agreed by HELCOM 38-2017. For some basins discussions on threshold values are still ongoing.
Eutrophication ratio (ER)	$ER = ES/ET$
Status confidence (ES-Score)	HIGH (=100%), if more than 15 spatially non-biased status observations are found each year. MODERATE (=50%), if more than 5 but no more than 15 status observations are found per year. LOW (=0%), if no more than 5 annual status observations are found during one or more years.
Indicator target confidence (ET-Score)	HIGH, if the target was based on numerous observations made earlier than the 1950's, possibly in combination with hindcast modelling. MODERATE, if the target was based on observations made earlier than the 1980's and/or hindcast modelling. LOW, if the target was set through expert judgement and/or information from reference sites and/or observations made during or after the 1980's.
Indicator confidence (I-Score)	Confidence (%) = average of ES-Score and ET-Score

Assessment unit

The indicator is assessed within the geographical HELCOM assessment unit scale 4: open sea sub-basin areas and coastal waters WFD coastal types and bodies.

The assessment units are defined in the [HELCOM Monitoring and Assessment Strategy Annex 4](#).

Relevance of the Indicator

Eutrophication assessment

The status of eutrophication is assessed using several core indicators. Each indicator focuses on one important aspect of the complex issue. In addition to providing an indicator-based evaluation of the total nutrients, this indicator also contributes to the overall eutrophication assessment along with the other biodiversity core indicators.

Policy relevance

Eutrophication is one of the four thematic segments of the HELCOM Baltic Sea Action Plan (BSAP) with the strategic goal of having a Baltic Sea unaffected by eutrophication (HELCOM 2007). Eutrophication is defined in the BSAP as a condition in an aquatic ecosystem where high nutrient concentrations stimulate the growth of algae which leads to imbalanced functioning of the system. The goal for eutrophication is broken down into five ecological objectives, of which one is "Concentrations of nutrients close to natural levels". Increase in nutrient concentrations can be assessed using measurements of all suspended and dissolved nutrients.

The EU Marine Strategy Framework Directive (Anonymous 2008) requires that "human-induced eutrophication is minimized, especially adverse effects thereof, such as losses in biodiversity, ecosystem degradation, harmful algal blooms and oxygen deficiency in bottom waters" (Descriptor 5). 'Total Phosphorus (TP)' is identified as an criteria element to be assessed using the criterion D5C1 'Nutrient concentrations are not at levels that indicate adverse eutrophication effects' in the Commission Decision on criteria and methodological standards on good environmental status of marine waters (Anonymous 2017).


The EU Water Framework Directive (Anonymous 2000) requires good ecological and chemical status in the European coastal waters. Good ecological status is defined in Annex V of the Water Framework Directive, in terms of the quality of the biological community including phytoplankton biomass (usually measured as chlorophyll-*a*), the hydromorphological/hydrological characteristics and the chemical characteristics. Nutrient concentrations, measured as total or inorganic nutrients, is one of the indicators listed in Annex V.

Role of total phosphorus in the ecosystem


Marine eutrophication is mainly caused by nutrient enrichment leading to increased production of organic matter in the Baltic Sea with subsequent effects on water transparency, phytoplankton communities, benthic fauna and vegetation as well as oxygen conditions. Phytoplankton and benthic vegetation need nutrients, mainly nitrate, ammonia and phosphorus, for growth.

Adding total nutrients alongside inorganic nutrients as core indicators strengthens the link from nutrient concentrations in the sea to nutrient enrichment. In particular these parameters allow to take account of climate change in the eutrophication assessment since higher temperatures will lead to year-round phytoplankton proliferation and / or possible changes in zooplankton communities. To illustrate this point, the concentration of the total and the dissolved inorganic fractions of nutrients have been compared, and diverging trends have been observed in some sub-basins. For example, an indication of decrease in winter DIP concentrations has been identified in the Arkona Basin during the last five years, but TP concentrations have remained somewhat unchanged (see figure below). A possible reason for this observation could be

that in winter more nutrients are bound in the phytoplankton due to the higher water temperatures. In such a situation, assessing only dissolved inorganic concentrations gives the wrong impression that nutrient concentrations seem to be declining, while, in fact, they are stable or increasing as can be seen when also assessing total concentrations. In conclusion, to get a good understanding of the trend in nutrient concentrations in the marine environment monitoring and assessing both, total and dissolved nutrients, is important.



Relevance figure 1. Time series of annual TP (black line and dots) and winter DIP (gray line and dots) in the Arkona Basin. The late (since 2008) decrease in winter DIP is not expressed by annual TP. The figure is modified from HELCOM 2013.



Relevance figure 2. Monthly values of total phosphorus concentration (as $\mu\text{mol l}^{-1}$) in the surface layer (0-10m) during 2007-2011.

Human pressures linked to the indicator

	General	MSFD Annex III, Table 2a
Strong link	Nutrient concentrations in the water column are affected by anthropogenic nutrient loads, both water- and airborne.	Substances, litter and energy - Input of nutrients – diffuse sources, point sources, atmospheric deposition - Input of organic matter – diffuse sources and point sources
Weak link		

Monitoring Requirements

Monitoring methodology

Monitoring of total phosphorus in the Contracting Parties of HELCOM is described on a general level in the **HELCOM Monitoring Manual in the [sub-programme: Nutrients](#)**

[Monitoring guidelines](#) specifying the sampling strategy are adopted and published.

Current monitoring

The monitoring activities relevant to the indicator that are currently carried out by HELCOM Contracting Parties are described in the HELCOM Monitoring Manual [sub-programme: Nutrients](#).

Description of optimal monitoring

For assessment purposes, at least 15 status observations should be conducted annually during the period January to December in each open sea assessment unit. The compilation of observations is expected to be distributed spatially within the assessment unit in a non-biased way. In coastal areas, at least monthly sampling of representative stations is desirable.

Data and updating

Access and use

The data and resulting data products (tables, figures and maps) available on the indicator web page can be used freely given that the source is cited. The indicator should be cited as following:

HELCOM (2017) Total nitrogen. HELCOM core indicator report. Online. [Date Viewed], [Web link].

ISSN 2343-2543

Metadata

[Result: Total phosphorus](#)

Data source: The average for 2011-2015 was estimated using monitoring data provided by the HELCOM Contracting Parties, and kept in the HELCOM COMBINE database, hosted by ICES (www.ices.dk). Nominated members of HELCOM STATE & CONSERVATION group were given the opportunity to review the data, and to supply any missing monitoring observations, in order to achieve a complete dataset.

Description of data: The data includes total phosphorus observations, determined as explained in the HELCOM COMBINE manual. Measurements made at the depth of 0 – 10 m from the surface were used in the assessment.

Temporal coverage: The raw data includes observations throughout the year, during the assessment period 2011-2015. For the summer average, observations taken during June-September were included only.

Data aggregation: The 2011-2015 averages for each sub-basin were produced as an inter-annual estimates using observations from all months / June-September.

Contributors and references

Contributors

Wera Leujak, Federal Environment Agency, Germany

Günther Nausch, Leibniz Institute for Baltic Sea Research Warnemünde (IOW), Germany

Marina Carstens, Ministry of Agriculture and Environment, Germany

Vivi Fleming-Lehtinen, HELCOM Secretariat and SYKE, Finland

Joni Kaitaranta, HELCOM Secretariat

Lena Avellan, HELCOM Secretariat

With contributions from other participants of HELCOM EUTRO-OPER and HELCOM IN-Eutrophication.

Archive

This version of the HELCOM core indicator report was published in July 2017:

[HOLAS II component - Core indicator report – web-based version July 2017](#) (pdf)

References

Andersen, J.H., Murray, C., Kaartokallio, H., Axe, P., Molvær, J., 2010. A simple method for confidence rating of eutrophication status assessments. *Mar. Pollut. Bull.* 60:919–924.

Anonymous (2000). Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 establishing a framework for community action in the field of water policy. *Official Journal of the European Communities L 327/1*.

Anonymous (2008). Directive 2008/56/EC of the European Parliament and of the Council. 17 June 2008. Establishing a Framework for Community Action in the Field of Marine Environmental Policy. *Marine Strategy Framework Directive*. 22p.

Anonymous (2017). COMMISSION DECISION (EU) 2017/848 of 17 May 2017 laying down criteria and methodological standards on good environmental status of marine waters and specifications and standardised methods for monitoring and assessment, and repealing Decision 2010/477/EU.

BLANO (2014). Harmonisierte Hintergrund- und Orientierungswerte für Nährstoffe und Chlorophyll-a in den deutschen Küstengewässern der Ostsee sowie Zielfrachten und Zielkonzentrationen für die Einträge über die Gewässer. Bund-Länder-Ausschuss Nord- und Ostsee (BLANO), Stand 6.10.2014, revised version 16.04.2015; <http://www.meeresschutz.info/sonstige-berichte.html>.

Federal Surface Water Ordinance (2016). Verordnung zum Schutz von Oberflächengewässern vom 20.6.2016. *Bundesgesetzblatt Jahrgang 2016, Teil 1, Nummer 28*, ausgegeben zu Bonn am 23. Juni 2016: 1373-1443.

https://www.bgbl.de/xaver/bgbl/start.xav?start=%2F%2F*%5B%40attr_id%3D%27bgbl116s0745.pdf%27%5D#_bgbl_%2F%2F*%5B%40attr_id%3D%27bgbl116s0745.pdf%27%5D_1496071661783.

Fleming-Lehtinen, V., Andersen, J.H., Carstensen, J., Łysiak-Pastuszek, E., Murray, C., Pyhälä, M. and Laamanen, M., 2015. Recent developments in assessment methodology reveal that the Baltic Sea eutrophication problem is expanding. *Ecological Indicators* 48:380-388.

HELCOM, 2007. Baltic Sea Action Plan. Baltic Sea Environment Protection Commission. 101p.

HELCOM, 2013. Approaches and methods for eutrophication target setting in the Baltic Sea region. In: Baltic Sea Environment Proceedings 133, 147 pp. Retrieved via:
<http://www.helcom.fi/Lists/Publications/BSEP133.pdf>

Hirt, U., Mahnkopf, J., Gadegast, M., Czudowski, L., Mischke, U., Heidecke, C., Schernewski, G. and Venhohr, M. 2013. Reference conditions for rivers of the German Baltic Sea catchment: reconstructing nutrient regimes using the model MONERIS. Reg Environ Change DOI 10.1007/s10113-013-0559-7.

Schernewski, G., Friedland, R., Carstens, M., Hirt, U., Leujak, W., Nausch, G., Neumann, T., Petenati, T., Sagert, S., Wasmund, N. and von Weber, M. 2015. Implementation of European marine policy: New water quality targets for German Baltic Waters. Marine Policy 51:305-321.

HELCOM core indicator report,
ISSN 2343-2543